

Balanis Quick Reference

Andrew Whelan

Contents

1	Time-Varying and Time-Harmonic Electromagnetic Fields	1
2	Electrical Properties of Matter	1
3	Wave Equation and its Solutions	2

1 Time-Varying and Time-Harmonic Electromagnetic Fields

1. Circuit theory is only valid when the physical dimensions of the circuit are small compared to the wavelength
2. Boundary conditions are derived by considering
 - (a) the integral curl equations on a tiny rectangle along the boundary and taking the limit as the height goes to zero, and
 - (b) the integral divergence equations on a cylinder along the boundary and taking the limit as the height goes to zero.
3. To get conservation of energy relations, consider
 - (a) a finite volume of space V ,
 - (b) scalar multiplying each curl equation by the other field,
 - (c) subtracting the equations, and
 - (d) use vector identity for divergence of cross product
4. Poynting vector \mathcal{S} and complex Poynting vector \mathbf{S} are given by

$$\mathcal{S} \stackrel{\text{def}}{=} \mathcal{E} \times \mathcal{H} \tag{1}$$

$$\begin{aligned} \mathbf{S} &\stackrel{\text{def}}{=} \mathcal{F}(\mathcal{S}) \stackrel{(1)}{=} \mathcal{F}(\mathcal{E} \times \mathcal{H}) \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} \mathbf{E}(\omega') \times \mathbf{H}(\omega - \omega') d\omega' \end{aligned} \tag{2}$$

$$\begin{aligned} \langle \mathcal{S} \rangle &\stackrel{\text{def}}{=} \lim_{T \rightarrow \infty} \frac{1}{T} \int_{-T/2}^{T/2} \mathcal{S}(t) dt = \mathbf{S}(\omega = 0) \\ &\stackrel{(2)}{=} \frac{1}{2\pi} \int_{\mathbb{R}} \mathbf{E}(\omega') \times \mathbf{H}^*(\omega') d\omega' \end{aligned} \tag{3}$$

2 Electrical Properties of Matter

1. Dielectric \implies no free charges \implies less resistance to electrical forces, and is quantified by ϵ . Think of bound charges storing elastic energy
2. Magnetic \implies less resistance to magnetic forces, and is quantified by μ .
3. Conductor \implies free charges and is quantified by σ . Think of free charges and dissipative energy
4. Media Properties:
 - (a) *Linear* : parameters are not functions of applied field (or are linearly mapped to them)

- (b) *Homogeneous* : parameters not function of position
- (c) *Isotropic* : parameters not function of field direction
- (d) *Dispersive* : parameters are function of frequency

3 Wave Equation and its Solutions

1. Assume a source-free, homogeneous medium, then we can derive the Helmholtz equations in the frequency domain:

$$(\nabla^2 - \gamma^2)E = 0 \tag{4a}$$

$$(\nabla^2 - \gamma^2)H = 0 \tag{4b}$$