



Indian Institute of Science, Bangalore
Department of Computational and Data Sciences (CDS)

DS284: Numerical Linear Algebra

Assignment 1 [Posted Aug 14, 2024]

Faculty Instructor: Dr. Phani Motamarri

TAs: Gourab Panigrahi, Srinibas Nandi, Nihar Shah, Rushikesh Pawar

Submission Deadline: NA

Max Points: NA

Notations: Vectors and matrices are denoted below by bold-faced lower case and upper case alphabets, respectively.

Problem 1

Let $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\mathbf{B} \in \mathbb{R}^{n \times p}$, $R(\cdot)$ denotes the range of the matrix, $N(\cdot)$ denotes the null space of a given matrix, $\dim(\cdot)$ denotes the dimension of a vector space, $\text{rank}(\cdot)$ denotes $\dim(R(\cdot))$, then prove the following:

- (a) $\dim[R(\mathbf{AB})] \leq \dim[R(\mathbf{A})]$
- (b) If the matrix \mathbf{B} is non-singular then $\dim[R(\mathbf{AB})] = \dim[R(\mathbf{A})]$
- (c) $\dim[N(\mathbf{AB})] \leq \dim[N(\mathbf{A})] + \dim[N(\mathbf{B})]$
- (d) $\dim[R(\mathbf{A})] + \dim[N(\mathbf{A})] = n$
- (e) $\text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}) - n \leq \text{rank}(\mathbf{AB}) \leq \min(\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B}))$
Hint: Use the result in (a)
- (f) Given a vector $\mathbf{u} \in \mathbb{R}^n$, $\text{rank}(\mathbf{uu}^T)$ is 1.
Hint: Use the result in (a)
- (g) Row rank always equals column rank.

Problem 2

Suppose there always exists a set of real coefficients $c_1, c_2, c_3, \dots, c_{10}$ for any set of real numbers $d_1, d_2, d_3, \dots, d_{10}$

$$\sum_{j=1}^{10} c_j f_j(i) = d_i \quad \text{for } i \in \{1, 2, \dots, 10\}$$

where $f_1, f_2, f_3, \dots, f_{10}$ are a set of functions defined on the interval $[1, 10]$

- (a) Use the concepts discussed in class to show that $d_1, d_2, d_3, \dots, d_{10}$ determine $c_1, c_2, c_3, \dots, c_{10}$ uniquely.
- (b) Let \mathbf{A} be a 10×10 matrix representing the linear mapping from data $d_1, d_2, d_3, \dots, d_{10}$ to coefficients $c_1, c_2, c_3, \dots, c_{10}$. What is the i, j th entry of \mathbf{A}^{-1} ?

Problem 3

A matrix \mathbf{S} is said to be symmetric if $\mathbf{S}^T = \mathbf{S}$ and skew-symmetric if $\mathbf{S}^T = -\mathbf{S}$. Now verify the following:

- (a) The matrix $\mathbf{Q} = (\mathbf{I} - \mathbf{S})^{-1}(\mathbf{I} + \mathbf{S})$ is an orthogonal matrix for any skew-symmetric matrix \mathbf{S} .
- (b) Note that a symmetric matrix $\mathbf{A} \in \mathbb{R}^{m \times m}$ can be decomposed as $\mathbf{Q}\mathbf{D}\mathbf{Q}^T$ where \mathbf{Q} is an orthogonal matrix and \mathbf{D} is a diagonal matrix. Using this result, show that $\mathbf{u}^T \mathbf{A} \mathbf{u} = 0 \forall \mathbf{u} \in \mathbb{R}^m$, if and only if $\mathbf{A} = 0$.
- (c) Show that “ $\mathbf{u}^T \mathbf{S} \mathbf{u} = 0 \forall \mathbf{u} \in \mathbb{R}^m$, if and only if \mathbf{S} is a skew-symmetric matrix.”

Problem 4

If $\mathbf{x} \in \mathbb{R}^m$ and $\mathbf{A} \in \mathbb{R}^{m \times n}$, then show the following.

- (a) $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2$
- (b) $\|\mathbf{x}\|_2 \leq \sqrt{m} \|\mathbf{x}\|_\infty$
- (c) $\|\mathbf{A}\|_\infty \leq \sqrt{n} \|\mathbf{A}\|_2$
- (d) $\|\mathbf{A}\|_2 \leq \sqrt{m} \|\mathbf{A}\|_\infty$
- (e) $\|\mathbf{A}\|_F = \sqrt{\text{tr}(\mathbf{A}^T \mathbf{A})}$
- (f) $\frac{1}{\sqrt{m}} \|\mathbf{A}\|_1 \leq \|\mathbf{A}\|_2 \leq \sqrt{n} \|\mathbf{A}\|_1$
- (g) $\|\mathbf{A}\|_2^2 \leq \|\mathbf{A}\|_1 \|\mathbf{A}\|_\infty$
Hint: $\|\mathbf{A}\|_2^2 = \text{maximum of absolute eigen values of } \mathbf{A}^T \mathbf{A}$

Problem 5

Induced matrix norm is defined as $\|\mathbf{A}\|^{(m,n)} = \max_{\mathbf{x}} \|\mathbf{Ax}\|^{(m)}$, where $\mathbf{x} \in \mathbb{R}^n$ and is a unit vector. $\|\cdot\|$ corresponds to p -norm ($1 \leq p < \infty$). For this exercise, let us consider p to be a natural number.

Using MATLAB/Octave/Python programming environment, create a matrix using “ $\mathbf{A} = \text{randn}(100, 2)$ ”. Subsequently, create random unit vectors \mathbf{x} using “ $\text{temp} = \text{randn}(2, 1)$ ” and normalize \mathbf{x} using “ $\mathbf{x} = \text{temp} / \text{norm}(\text{temp})$ ”. Check for multiple random vectors \mathbf{x} (use a loop, and check for about 1000 random vectors \mathbf{x}) using “ $\text{norm_of_Ax} = \text{norm}(\mathbf{Ax}, p)$ ” for $p = 1, 2, 3, 4, 5, 6, \infty$. What is the maximum value of p -norm for the vector \mathbf{Ax} ? Now calculate p -norm of \mathbf{A} using “ $\text{norm_of_A} = \text{norm}(\mathbf{A}, p)$ ” for $p = 1, 2, \infty$ within the same programming environment you used before. Verify the equality $\|\mathbf{A}\|^{(m,n)} = \max_{\mathbf{x}} \|\mathbf{Ax}\|^{(m)}$ for $p = 1, 2, \infty$. Note that this equality is true for other values of p as well but you are restricting to $p = 1, 2, \infty$ in this exercise.