Check for updates

Hydroboration


International Edition: DOI: 10.1002/anie.201915716
German Edition: DOI: 10.1002/ange.201915716

Cobalt-Catalyzed Regio- and Stereoselective Hydroboration of Allenes

Can Li[+], Zheng Yang[+], Lei Wang, Yinlong Guo, Zheng Huang,* and Shengming Ma**

Abstract: *An efficient pincer-ligand-based cobalt-complex-catalyzed allene hydroboration affording Z-allylic boronates is described. The reaction demonstrates an excellent regio- as well as Z-stereoselectivity and a wide substrate scope that tolerates many functional groups. Based on solvent-assisted electrospray ionization mass spectrometry (SAESI-MS) studies, a rationale for the cobalt-catalyzed hydroboration involving the highly selective insertion of an allene into the Co−H bond to form Z-allylic cobalt intermediates is proposed.*

Organic boronic acid derivatives are of low toxicity and decent stability, and thus serve as versatile intermediates for many synthetic transformations.[1,2] Of particular interest are the stereo-defined allylic boronic acid derivatives, which may provide a synthetically attractive allylic entity into organic skeletons. Compared with previous approaches to allylic boronic acid derivatives,[3] transition-metal-catalyzed hydroboration of readily available allenes with pinacolborane (HBpin) would be, in principle, a very efficient method featuring excellent atom economy. However, compared with the well-established catalytic hydroboration of alkenes,[4] alkynes,[5] and 1,3-dienes,[3b,6] the hydroboration of allenes has remained underdeveloped.[7,8] In 2009, a one-pot two-step reaction to form homoallylic alcohols involving non-catalytic hydroboration of alkyl-substituted terminal allenes with a chiral borane reagent was reported, however, the stereoselectivity is rather poor (Scheme 1 a).[9] In 2013, Tsuji and co-workers reported a copper-catalyzed hydroboration of mono-substituted terminal allenes to form linear *E*-allylic boronates as the major products, although the *E*/*Z* selectivity is subject to the steric effect of the R group (Scheme 1 b).[10] In 1999, Miyaura and co-workers reported the preparation of linear *Z*-allylic boronates as the major products via platinum-catalyzed hydroboration of 2,3-propadienyl *tert*-butyldimethylsilyl (TBS) ether or methyl ether with *Z*/*E* ratios ranging from 84:16 to 91:9 (Scheme 1 c).[11] Herein, we report a cobalt-catalyzed hydroboration of allenes with pinacolborane (HBpin) to provide *Z*-allylboronates with an excellent regio- and stereoselectivity (Scheme 1 d).




Scheme 1. Hydroboration of allenes affording alkenyl or allylic boronates.

Based on the pincer ligand-based iron and cobalt complexes developed in Huang's group,[12] initially, ($^{t\mathrm{Bu}}$P^{O}NN$^{i\mathrm{Pr}}$)FeCl$_2$ (**1**) was chosen as the precatalyst for the hydroboration of 1,2-undecadiene (**3a**) with HBpin in toluene. Unfortunately, only a complicated mixture was formed. Subsequently, a series of complexes of Fe and Co supported by various P^{C}NN ligands[12] were investigated. To our delight, by using ($^{t\mathrm{Bu}}$P^{C}NN$^{i\mathrm{Pr}}$)FeCl$_2$ (**2a**) as the precatalyst, the reaction afforded *Z*-**4a** in a high yield together with small amounts of regioisomers **5a** and **6a** as determined by ^{1}H NMR analysis of the crude reaction mixture (Table 1, entry 2). The difference between **1** and **2a** is largely due to the increased steric hindrance around the metal center. Interest-

[*] C. Li,[+] Dr. Z. Yang,[+] L. Wang, Prof. Dr. Y.-L. Guo, Prof. Dr. Z. Huang, Prof. Dr. S.-M. Ma
State Key Laboratory of Organometallic Chemistry
Shanghai Institute of Organic Chemistry
Chinese Academy of Sciences
345 Lingling Lu, Shanghai 200032 (P. R. China)
E-mail: ylguo@sioc.ac.cn
huangzh@sioc.ac.cn
masm@sioc.ac.cn
C. Li[+]
University of Chinese Academy of Sciences
Beijing 100049 (P. R. China)
Prof. Dr. S.-M. Ma
Department of Chemistry, Fudan University
220 Handan Lu, Shanghai 200433 (P. R. China)
[+] These authors contributed equally to this work.
Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under: https://doi.org/10.1002/anie.201915716.

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Table 1: Optimization of the reaction conditions.[a]



Entry	Precatalyst	Solvent	NMR yield [%][b] *Z*-**4a**:**5a**:**6a**	Recovery of **3a** [%]
1	**1**	toluene	–	–
2	**2a**	toluene	85:7:6	–
3	**2b**	toluene	57:0:0	40
4[c]	**2b**	toluene	90(87):0:0	–
5	**2c**	toluene	41:12:12	–
6	**2d**	toluene	40:0:0	45
7	**2d**	toluene	66:13:0	22
8	**2e**	toluene	38:0:0	45
9[c]	**2b**	Et_2O	88:0:0	–
10[c]	**2b**	hexane	87:0:0	–
11[c]	**2b**	THF	60:0:0	–
12[c]	–	toluene	–	89
13[c,d]	**2b**	toluene	–	93

[a] Reaction conditions: **3a** (0.2 mmol), HBpin (0.2 mmol), 2 μmol of precatalyst, and 4 μmol of $NaBHEt_3$ in toluene (0.5 mL) for 4 h. [b] Determined by ^1H NMR analysis of the crude product with CH_3NO_2 as the internal standard. The value in parentheses is the isolated yield of product *Z*-**4a**. [c] **3a** (0.2 mmol), HBpin (0.2 mmol), 3 mol% of precatalyst, and 6 mol% of $NaBHEt_3$ in solvent (1 mL) for 24 h. [d] $NaBHEt_3$ was not added.

ingly, by using the cobalt complex with the same pincer ligand **2b** (structure established by X-ray diffraction; Table 1),[13] the reaction afforded *Z*-**4a** exclusively, albeit with 40% of **3a** being recovered. Further screening of the precatalysts revealed that Fe complex **2c** with a less hindered $^{iPr}P^CNN^{iPr}$ ligand is more active than the Co variant **2d** with the same ligand, but the former is much less selective (Table 1, compare entry 5 with entry 6). The Co complex **2e** with the least sterically demanding ligand among the PNN ligands investigated here is less efficient than **2b**, albeit with an excellent selectivity (Table 1, entry 8 vs. entry 2). Further optimization led to the observation that increasing the loading of precatalyst **2b** from 1 to 3 mol% afforded the hydroboration product *Z*-**4a** in 90% yield without the detection of other isomers (Table 1, entry 4). The yield dropped when the reaction was conducted in ethyl ether, hexane, or THF, although the selectivity remained unaffected (Table 1, entries 9–11). No conversion was observed in the absence of the metal complex or the catalyst activator $NaBHEt_3$ (Table 1, entries 12 and 13). Therefore, the conditions used in entry 4 were chosen as the optimized reaction conditions for further study.

Next, we investigated the scope of cobalt-catalyzed allene hydroboration. Various functionalized allenes underwent hydroboration with HBpin in high yields and excellent regio- and stereoselectivity at ambient temperature (Table 2). Hydroboration of alkyl-substituted allenes proceeded smoothly under the standard reaction conditions (**4a**–**4e**). The gram-scale reaction of **3a** with HBpin (5 mmol) afforded **4a** in 83% yield. Synthetically useful functional groups such as ketal (**4f**), OTBS (**4g**), benzyloxy (**4h**), halogen (**4i** and **4j**), ester (**4k**), and N-heterocycles (**4n**, **4o**, and **4p**) could be tolerated. Even the very bulky silyl-substituted allene **3l** could also be accommodated, affording the corresponding *Z*-allylboronate **4l** containing both C–B and C–Si bonds with a 97:3 *Z*/*E* selectivity. Exclusive *Z*-selectivity was observed even for product **4m**, which showed a *Z*/*E* selectivity of ≈90:10 as reported in the literature.[11] The high selectivity for the formation of allylic boronate *Z*-**4c** from cyclohexylpropadiene is also noteworthy because it afforded vinylic boronates under the conditions reported by Tsuji and co-workers.[10]

Although cobalt complexes have been reported to catalyze the hydroboration of alkenes,[4e,g,h] under the optimized conditions, allenes exhibit a much higher reactivity than both terminal and internal alkenes as demonstrated by the formation of *Z*-**4q**, *Z*-**4r**, and *Z*-**4s** in high yields with exclusive chemoselectivity.


Hydroboration of 1,1-disubstituted allenes also worked well allowing for the construction of trisubstituted (*Z*)-allylic boronates **4t**–**4x**. The stereoselectivity is determined by the steric effect of these two substituents as shown by the *Z*/*E* ratios of **4u**, **4w**, and **4x**. Even estrone 3-methyl-ether-derived product *Z*-**4v** was formed and isolated in a high yield and an excellent *Z*/*E* selectivity. Double hydroboration of Si- or C-tethered bisallenes **3y** and **3z** afforded bisallylic boronates **4y** and **4z** in high yields and an excellent *Z*-selectivity from the reaction with 2.2 equiv of HBpin, without formation of the monohydroboration products.

To demonstrate the synthetic utility of this methodology, synthetic transformations of (*Z*)-**4a** were conducted (Scheme 2). Linear allylic alcohol **7** or branched allylic alcohol **8** could be obtained in high yields by using H_2O_2[14] or PhNO[15] as the oxidant, respectively. Suzuki–Miyaura coupling reaction of (*Z*)-**4a** and iodobenzene afforded the corresponding branched allylbenzene derivative (**9**). Moreover, (*Z*)-**4a** could react with benzaldehyde,[16] furnishing homoallylic alcohol **10** with a high diastereoselectivity.


Based on previous reports of cobalt-catalyzed hydroborylation reactions,[4e,18] a rationale for the cobalt-catalyzed hydroboration of allenes is proposed as shown in Scheme 3. The precatalyst **2b** would react with $NaBHEt_3$, affording the catalytically active Co^I hydride species **Int 1**, followed by the coordination with the terminal C=C double bond of substrate **3a** from the less hindered side to form **Int 2-A** (Scheme 3b shows that **Int 2-A** is sterically more favorable than **Int 2-B**). Following the insertion of a double bond into the Co–H bond, a *Z*-allylic cobalt intermediate **Int 3-A** is formed. Subsequent reaction with HBpin, by either σ-bond metathesis or oxidative addition and reductive elimination, would furnish the hydroboration product *Z*-**4a** and regenerate **Int 1** (Scheme 3a).

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Table 2: Z-selective hydroboration of various allenes.[a]



[a] The reaction was performed with 1 equiv of **3**, 1 equiv of HBpin, 3 mol% of **2b**, and 6 mol% of $NaBHEt_3$ in toluene (0.2 M) at room temperature for 24 h on 1.0 or 0.5 mmol scale. Yields of isolated products are given. [b] The reaction was carried out on 5 mmol scale to afford 1.16 g of *Z*-**4a**. [c] 5 mol% of **2b** and 10 mol% of $NaBHEt_3$ were used. [d] The reaction was run for 43 h. [e] The reaction was run for 36 h. [f] The reaction was run for 35 h. [g] The reaction was run for 54 h. [h] The reaction was run for 22 h. [i] The reaction was run for 38.5 h. [j] The reaction was carried out at 50 °C for 48 h. [k] 5 mol% of **2b** and 10 mol% of $NaBHEt_3$ were used. The reaction was carried out at 50 °C for 48 h.



Scheme 2. Synthetic applications.

Alternatively, the Co^{I} hydride species **Int 1** may react with HBpin to form cobalt(I)-Bpin intermediate **Int 1′**.[17] **Int 1′** then undergoes a coordination and insertion processes to yield the vinyl cobalt intermediate **Int 3′** through **Int 2′**. Subsequently, the reaction of **Int-3′** with HBpin affords the hydroboration product *Z*-**4a** and regenerates **Int-1′** (Scheme 3c).

To distinguish between these two pathways, we sought to identify the catalytic intermediates of the catalytic reaction by solvent-assisted electrospray ionization mass spectrometry (SAESI-MS) and SAESI-MS/MS analysis. Two species with molecular weights matching those of Co^{I}-H **Int-1** (*m*/*z* 497, $[M-H]^{+}$) and *Z*-allylic cobalt **Int 3-A** (*m*/*z* 650, $[M]^{\cdot+}$) were detected (Figure 1; see the Supporting Information for details),[18] whereas the species corresponding to Co^{I}-Bpin intermediate **Int-1′** (*m*/*z* 624) and vinyl cobalt intermediate

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim





Figure 1. SAESI-MS studies. a) SAESI-MS spectrum of the solution. b) Expanded SAESI-MS spectrum, showing the major signal of **Int 1** $[C_{28}H_{43}CoN_2P]^+$ at *m*/*z* 497 and **Int 3-A** $[C_{39}H_{64}CoN_2P]^{\bullet+}$ at *m*/*z* 650. c) SAESI-MS/MS spectrum of complex ion **Int 1** $[C_{28}H_{43}CoN_2P]^{\bullet+}$ at *m*/*z* 497. d) SAESI-MS/MS spectrum of complex ion **Int 3-A** $[C_{39}H_{64}CoN_2P]^{\bullet+}$ at *m*/*z* 650.



Scheme 3. Proposed mechanisms.

Int 3′ (*m*/*z* 776) were not observed. The structures of **Int 1** and **Int 3-A** were further confirmed by SAESI-MS/MS experiments (Figure 1c and d). Such SAESI-MS data provide evidence in support of the catalytic cycle involving the intermediacies of Co^I-H **Int 1** and *Z*-allyl-cobalt **Int 3-A**, as shown in Scheme 3a.

In summary, we have developed a highly efficient hydroboration of allenes catalyzed by P^CNN-cobalt complex **2b** and identified two of the catalytic intermediates by using SAESI-MS and SAESI-MS/MS analyses. This reaction features excellent regio- and *Z*-stereoselectivity, 100% atom economy, and a broad substrate scope with functionalized mono- and bisallenes. In addition, for the reaction of enallenes, the exclusive hydroboration of the allene moiety was observed, leaving the alkene moiety untouched. (*Z*)-Allylboronate products could be transformed into different allylic alcohols,

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

or underwent Suzuki–Miyaura coupling reactions to form C–C bonds.

Acknowledgements

Financial support from National Natural Science Foundation of China (21690063 for S. Ma; 21825109 for Z. Huang) is greatly appreciated. We thank Mr. Penglin Wu in this group for reproducing the results for (*Z*)-**4e**, (*Z*)-**4i**, and (*Z*)-**4q** as presented in this study.

Conflict of interest

The authors declare no conflict of interest.

Keywords: allenes · cobalt catalysis · hydroboration · regioselectivity · stereoselectivity

How to cite: *Angew. Chem. Int. Ed.* **2020**, *59*, 6278–6283
Angew. Chem. **2020**, *132*, 6337–6342

[1] a) H. C. Brown, *Organic Synthesis* via *Boranes*, Wiley Interscience, New York, **1975**; b) D. G. Hall, *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*, Wiley-VCH, Weinheim, **2005**; c) G. C. Fu, *Acc. Chem. Res.* **2008**, *41*, 1555–1564; d) M. Tobisu, N. Chatani, *Angew. Chem. Int. Ed.* **2009**, *48*, 3565–3568; *Angew. Chem.* **2009**, *121*, 3617–3620; e) A. Bonet, M. Odachowski, D. Leonori, S. Essafi, V. K. Aggarwal, *Nat. Chem.* **2014**, *6*, 584–589.

[2] For reviews of the Suzuki–Miyaura cross-coupling reaction, see: a) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457–2483; b) A. Suzuki, *J. Organomet. Chem.* **1999**, *576*, 147–168; c) R. Jana, T. P. Pathak, M. S. Sigman, *Chem. Rev.* **2011**, *111*, 1417–1492; d) A. Suzuki, *Angew. Chem. Int. Ed.* **2011**, *50*, 6722–6737; *Angew. Chem.* **2011**, *123*, 6854–6869; e) C. Amatore, G. L. Duc, A. Jutand, *Chem. Eur. J.* **2013**, *19*, 10082–10093.

[3] For selected examples of other methods for the synthesis of allylboronates, see: a) N. Selander, K. J. Szabó, *J. Org. Chem.* **2009**, *74*, 5695–5698; b) R. J. Ely, J. P. Morken, *J. Am. Chem. Soc.* **2010**, *132*, 2534–2535; c) P. Zhang, I. A. Roundtree, J. P. Morken, *Org. Lett.* **2012**, *14*, 1416–1419; d) L.-J. Mao, R. Bertermann, S. G. Rachor, K. J. Szabó, T. B. Marder, *Org. Lett.* **2017**, *19*, 6590–6593; e) For a report on the application of the cross-metathesis of 2-propenyl borate and terminal alkenes catalyzed by W or Co-based MAP complexes under 100 torr pressure, see: E. T. Kiesewetter, R. V. O'Brien, E. C. Yu, S. J. Meek, R. R. Schrock, A. H. Hoveyda, *J. Am. Chem. Soc.* **2013**, *135*, 6026–6029.

[4] For selected examples of hydroboration of alkenes, see: a) D. Männig, H. Nöth, *Angew. Chem. Int. Ed. Engl.* **1985**, *24*, 878–879; *Angew. Chem.* **1985**, *97*, 854–855; b) D. Noh, H. Chea, J. Ju, J. Yun, *Angew. Chem. Int. Ed.* **2009**, *48*, 6062–6064; *Angew. Chem.* **2009**, *121*, 6178–6180; c) L. Zhang, D.-J. Peng, X.-B. Leng, Z. Huang, *Angew. Chem. Int. Ed.* **2013**, *52*, 3676–3680; *Angew. Chem.* **2013**, *125*, 3764–3768; d) M. D. Greenhalgh, S. P. Thomas, *Chem. Commun.* **2013**, *49*, 11230–11232; e) J. V. Obligacion, P. J. Chirik, *J. Am. Chem. Soc.* **2013**, *135*, 19107–19110; f) J. V. Obligacion, P. J. Chirik, *Org. Lett.* **2013**, *15*, 2680–2683; g) L. Zhang, Z.-Q. Zuo, X.-B. Leng, Z. Huang, *Angew. Chem. Int. Ed.* **2014**, *53*, 2696–2700; *Angew. Chem.* **2014**, *126*, 2734–2738; h) J.-H. Chen, T. Xi, X. Ren, B. Cheng, J. Guo, Z. Lu, *Org. Chem. Front.* **2014**, *1*, 1306–1309; i) A. J. MacNair, C. R. P. Millet, G. S. Nichol, A. Ironmonger, S. P. Thomas, *ACS Catal.* **2016**, *6*, 7217–7221; j) N. W. J. Ang, C. S. Buettner, S. Docherty, A. Bismuto, J. R. Carney, J. H. Docherty, M. J. Cowley, S. P. Thomas, *Synthesis* **2018**, *50*, 803–808; k) A. Bismuto, M. J. Cowley, S. P. Thomas, *ACS Catal.* **2018**, *8*, 2001–2005; l) C. Xue, Y. Luo, H.-L. Teng, Y.-H. Ma, M. Nishiura, Z.-M. Hou, *ACS Catal.* **2018**, *8*, 5017–5022; m) N. Hu, G. Zhao, Y. Zhang, X. Liu, G. Li, W. Tang, *J. Am. Chem. Soc.* **2015**, *137*, 6746–6749.

[5] For selected examples of hydroboration of alkynes, see: a) H. Jang, A. R. Zhugralin, Y. Lee, A. H. Hoveyda, *J. Am. Chem. Soc.* **2011**, *133*, 7859–7871; b) M. Haberberger, S. Enthaler, *Chem. Asian J.* **2013**, *8*, 50–54; c) J. V. Obligacion, J. M. Neely, A. N. Yazdani, I. Pappas, P. J. Chirik, *J. Am. Chem. Soc.* **2015**, *137*, 5855–5858; d) K. Nakajima, T. Kato, Y. Nishibayashi, *Org. Lett.* **2017**, *19*, 4323–4326; e) J. R. Lawson, L. C. Wilkins, R. L. Melen, *Chem. Eur. J.* **2017**, *23*, 10997–11000; f) N. Gorgas, L. G. Alves, B. Stöger, A. M. Martins, L. F. Veiros, K. Kirchner, *J. Am. Chem. Soc.* **2017**, *139*, 8130–8133; g) K. Nagao, A. Yamazaki, H. Ohmiya, M. Sawamura, *Org. Lett.* **2018**, *20*, 1861–1865; h) L. Carreras, M. Serrano-Torné, P. W. N. M. van Leeuwen, A. Vidal-Ferran, *Chem. Sci.* **2018**, *9*, 3644–3648.

[6] For selected examples of hydroboration of 1,3-dienes, see: a) J. Y. Wu, B. Moreau, T. Ritter, *J. Am. Chem. Soc.* **2009**, *131*, 12915–12917; b) D. Noh, S. K. Yoon, J. Won, J. Y. Lee, J. Yun, *Chem. Asian J.* **2011**, *6*, 1967–1969; c) Y.-C. Cao, Y.-L. Zhang, L. Zhang, D. Zhang, X.-B. Leng, Z. Huang, *Org. Chem. Front.* **2014**, *1*, 1101–1106; d) D. Fiorito, C. Mazet, *ACS Catal.* **2018**, *8*, 9382–9387.

[7] For selected examples of Cu-catalyzed hydroborylation with diboron, see: a) S. B. Thorpe, X. Guo, W. L. Santos, *Chem. Commun.* **2011**, *47*, 424–426; b) W. Yuan, S. Ma, *Adv. Synth. Catal.* **2012**, *354*, 1867–1872; c) W. Yuan, X. Zhang, Y. Yu, S. Ma, *Chem. Eur. J.* **2013**, *19*, 7193–7202; d) B. Jung, A. H. Hoveyda, *J. Am. Chem. Soc.* **2012**, *134*, 1490–1493; e) F.-K. Meng, B. Jung, F. Haeffner, A. H. Hoveyda, *Org. Lett.* **2013**, *15*, 1414–1417.

[8] For example of Pd-catalyzed hydroborylation with $(BPin)_2$, see: C. Zhu, B. Yang, Y.-A. Qiu, J.-E. Bäckvall, *Chem. Eur. J.* **2016**, *22*, 2939–2943.

[9] J. Kister, A. C. DeBaillie, R. Lira, W. R. Roush, *J. Am. Chem. Soc.* **2009**, *131*, 14174–14175.

[10] K. Semba, M. Shinomiya, T. Fujihara, J. Terao, Y. Tsuji, *Chem. Eur. J.* **2013**, *19*, 7125–7132.

[11] Y. Yamamoto, R. Fujikawa, A. Yamada, N. Miyaura, *Chem. Lett.* **1999**, *28*, 1069–1070.

[12] a) D.-J. Peng, Y.-L. Zhang, X.-Y. Du, L. Zhang, X.-B. Leng, M. D. Walter, Z. Huang, *J. Am. Chem. Soc.* **2013**, *135*, 19154–19166; b) X.-Y. Du, Y.-L. Zhang, D.-J. Peng, Z. Huang, *Angew. Chem. Int. Ed.* **2016**, *55*, 6671–6675; *Angew. Chem.* **2016**, *128*, 6783–6787; c) X.-Y. Du, W.-J. Hou, Y.-L. Zhang, Z. Huang, *Org. Chem. Front.* **2017**, *4*, 1517–1521; d) Z. Yang, D. Peng, X. Du, Z. Huang, S. Ma, *Org. Chem. Front.* **2017**, *4*, 1829–1832.

[13] Crystal data for **2b**: $C_{28}H_{43}Cl_2CoN_2P$, MW = 568.44, monoclinic, space group *P*21/*c*, final *R indices* $[I > 2\sigma(I)]$, *R1* = 0.0774, *wR2* = 0.1648; *R indices* (all data), *R1* = 0.1186, *wR2* = 0.1849; *a* = 18.196(2), *b* = 10.2119(12), *c* = 15.6291(19) Å, $\alpha = 90$, $\beta = 92.762(4)$, $\gamma = g = 90°$, $V = 2900.8(6)$ Å^3, $T = 193(2)$ K, $Z = 4$, reflections collected/unique 24238/ 5094 ($R_{int} = 0.0888$). CCDC 1906720 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

[14] H. Y. Cho, Z.-Y. Yu, J. P. Morken, *Org. Lett.* **2011**, *13*, 5267–5269.

[15] B. Potter, A. A. Szymaniak, E. K. Edelstein, J. P. Morken, *J. Am. Chem. Soc.* **2014**, *136*, 17918–17921.

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

[16] C. Zhu, B. Yang, B. K. Mai, S. Palazzotto, Y.-A. Qiu, A. Gudmundsson, A. Ricke, F. Himo, J.-E. Bäckvall, *J. Am. Chem. Soc.* **2018**, *140*, 14324 – 14333.
[17] a) L. Zhang, Z. Huang, *J. Am. Chem. Soc.* **2015**, *137*, 15600 – 15603; b) H.-N. Wen, L. Zhang, S.-Z. Zhu, G.-X. Liu, Z. Huang, *ACS Catal.* **2017**, *7*, 6419 – 6425; c) S. Krautwald, M. J. Bezdek, P. J. Chirik, *J. Am. Chem. Soc.* **2017**, *139*, 3868 – 3875.
[18] J.-T. Zhang, H.-Y. Wang, W. Zhu, T.-T. Cai, Y.-L. Guo, *Anal. Chem.* **2014**, *86*, 8937 – 8942.

Manuscript received: December 9, 2019
Accepted manuscript online: January 8, 2020
Version of record online: February 18, 2020

 © 2020 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim