ORGANIC CHEMISTRY FRONTIERS

RESEARCH ARTICLE

Published on 18 September 2024
Licence and permissions

Check for updates

Cite this: *Org. Chem. Front.*, 2025, **12**, 2401

Photoredox copper-catalyzed regioselective syntheses of allenols *via* HAT-mediated propargyl radical generation†

Wenyao Li,[a] Hao Yu *[a] and Shengming Ma *[a,b]

Selectivity control of the reactions of propargylic and/or allenylic radicals remains particularly elusive. Here, a visible light-mediated Cu-catalyzed protocol for the highly selective syntheses of tri- and tetra-substituted allenols from alkynol derivatives has been developed. The synthetic potential of the products has been demonstrated. The control experiments, Stern–Volmer quenching, and cyclic voltammetry studies support a reaction pathway involving the generation of allenyl radicals *via* photo-mediated N–O bond scission, an intramolecular 1,5-HAT process, the resonance to propargylic radicals, and coupling. A non-common intramolecular 1,6-HAT process has also been observed.

Received 8th July 2024,
Accepted 7th September 2024

DOI: 10.1039/d4qo01245c

rsc.li/frontiers-organic

Introduction

Selectivity control is an everlasting topic in organic synthesis, advanced materials, and medicinal chemistry or life science.[1] For example, it is of particular interest to see the chemistry on the propargylic/allenylic species due to the versatile potential of alkynes and allenes. Recently, much attention has been paid to the generation and reactivity of propargylic and/or allenylic radicals affording alkynes[2] and/or allenes.[3–6] However, the efficient generation of propargylic radicals and the selectivity-controlled trapping still remain particularly elusive (Fig. 1a). 1,5-Hydrogen atom transfer (HAT)[7,8] mediated by alkoxyl radicals[9,10] has emerged as an important strategy for C(sp^3)–H bond functionalization (Fig. 1b). Given the high efficiency of radical formation with this specific method, we wonder whether such an alkoxyl radical-induced HAT of alkynes would efficiently generate propargylic radicals followed by transition metal-catalyzed functionalization, providing an efficient and selective platform for synthesizing multi-substituted alkynes or allenes (Fig. 1c). However, notable challenges for such a concept included (Fig. 1c): firstly, the tautomeric equilibrium of propargylic radicals to allenyl radicals and the resulting control of selectivity for trapping of these two radicals;[2–4] secondly, the introduction of the alkyne functional group and the formation of the allene functional group may lead to direct cyclic radical addition reactions;[11] and thirdly, the facile HAT of allenyl radicals could produce other non-functionalized allene by-products before the transition metal-




Fig. 1 Development of a photoredox/copper cocatalysis strategy for allenol syntheses based on 1,5-HAT.

[a]*Research Center for Molecular Recognition and Synthesis, Department of Chemistry, Fudan University, 220 Handan Lu, Shanghai 200433, P. R. China. E-mail: hao_yu@fudan.edu.cn*
[b]*State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Lu, Shanghai 200032, P. R. China. E-mail: masm@sioc.ac.cn*

† Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d4qo01245c


catalyzed functionalization occurs. In addition, it would be a platform reaction for the study on how far a HAT reaction may go. Here, we report our success on the exclusive synthesis of tri- and tetra-substituted allenes *via* visible-light photoredox and copper dual catalysis (Fig. 1d).

Results and discussion


Our initial attempt used *N*-alkynoxyphthalimide **1a**, which is easily obtained by the Mitsunobu reaction from the corresponding alkynol. We began investigating the coupling reaction of **1a** with trimethylsilyl cyanide by applying a catalytic system consisting of 4CzIPN and CuCl/dmbpy in CH_3CN under blue LED irradiation, and the desired allene product **2a′** was afforded in 38% yield as determined by ^1H NMR spectroscopy (Table 1, entry 1), without the formation of an alkyne product. The photocatalyst $MesAcr^+ClO_4^-$ resulted in a largely decreased yield of the allene product **2a′** (Table 1, entry 2). Further exploration of photocatalysts revealed that iridium complexes, such as $Ir[(dtbbpy)(ppy)_2]PF_6$ or $Ir(ppy)_3$, were more efficient (Table 1, entries 3 and 4). Other copper catalysts such as CuBr and CuI (Table 1, entries 5 and 6) were also screened and CuCl was still the best. Subsequently, we tested a series of other bipyridine ligands, and identified 1,10-phenanthroline as equally excellent as dmbpy (Table 1, entry 7). We chose 1,10-phen as the ligand of preference finally in consideration of the price. Other alkynoxylamide derivatives, such as **3a** and **4a**, exhibited a much lower reactivity (Table 1, entries 8 and 9). Control experiments demonstrated that the CuCl/1,10-phen system, photocatalyst and light are indispensable for enabling this transformation to proceed smoothly (entries 10–12).

With the optimized conditions in hand, we conducted further investigations on the substrate scope of this reaction (Fig. 2). We found that a wide range of terminal alkynols are suitable for this transformation, leading to the successful preparation of the corresponding tri-substituted allenols **2a–2t** smoothly. In terms of the alkyl groups R^1 at the propargyl position, non-functionalized linear primary alkyl chains of various lengths (**2a–2d**) as well as alkyl substituents bearing ethers (**2e** and **2f**), chloro (**2g**), acetal (**2h**), terminal olefin (**2i**), and aryls

Table 1 Optimization of the reaction conditions[a]



Entry	[Cu]	PC	Ligand	Yields of **2a′**/**2a**[b] (%)	RSM[b] (%)
1	CuCl	4CzIPN	dmbpy	38/0	50
2	CuCl	$MesAcr^+ClO_4^-$	dmbpy	7/0	78
3	CuCl	$Ir[(dtbbpy)(ppy)_2]PF_6$	dmbpy	80/0	0
4	CuCl	$Ir(ppy)_3$	dmbpy	81/0	0
5	CuBr	$Ir(ppy)_3$	dmbpy	79/0	0
6	CuI	$Ir(ppy)_3$	dmbpy	60/0	0
7	CuCl	$Ir(ppy)_3$	1,10-phen	81/0	0
8[c]	CuCl	$Ir(ppy)_3$	1,10-phen	0/0	79
9[d]	CuCl	$Ir(ppy)_3$	1,10-phen	24/0	75
10	—	$Ir(ppy)_3$	—	12/0	2
11	CuCl	—	1,10-phen	0/0	96
12[e]	CuCl	$Ir(ppy)_3$	1,10-phen	0/0	92



[a] The reaction was conducted with **1a** (0.2 mmol), PC (1 mol%), [Cu] (10 mol%), **L** (12 mol%), and TMSCN (0.6 mmol) in CH_3CN (1 mL) under an Ar atmosphere, with blue LED (50 W, 460 nm) irradiation for 24 h. [b] The yield was determined by ^1H NMR spectroscopy with CH_2Br_2 as the internal standard. [c] **3a** instead of **1a**. [d] **4a** instead of **1a**. [e] The reaction was conducted without light. dmbpy = 4,4′-dimethyl-2,2′-bipyridine. 1,10-phen = 1,10-phenanthroline.




Fig. 2 Substrate scope for the syntheses of tri- and tetra-substituted allenols.

This journal is © the Partner Organisations 2025

(**2j**–**2m**) were all found to be compatible under the standard conditions. R^1 may also be secondary alkyl groups (**2n**–**2p**) or even a tertiary alkyl group (**2q**), resulting in the corresponding allenols in modest to good yields. Furthermore, beyond the alkyl-substituted substrates, the reaction of the substrate with R^1 as phenyl yielded **2r** smoothly, achieving a yield of 66%. Furthermore, the synthesis of secondary allenols **2s** and **2t** was realized with yields of 76% and 60%, respectively. Finally, this protocol also worked for internal alkynols with R^2 being alkyl, phenylalkyl or phenyl groups, affording tetra-substituted allenols (**2u**–**2ae**).




Fig. 3 Investigation of the 1,4- and 1,6-HAT pathways.

Furthermore, various *N*-alkynoxyphthalimide precursors with diverse tether sizes were employed to investigate the potential of our protocol for 1,4- or 1,6-HAT (Fig. 3). When subjected to the standard conditions, compound **1af** failed to yield the corresponding beta-allenol **2af**, with the complete consumption of the starting material observed (Fig. 3a). Conversely, the reaction of substrate **1ag** with TMSCN resulted in the formation of delta-allenol **2ag** in 45% yield, involving 1,6-HAT. Furthermore, endeavors to synthesize the initial *N*-alkynoxyphthalimide precursor for alpha-allenols proved unsuccessful (see Fig. S4 in the ESI†).

To demonstrate the synthetic potential of this reaction, a gram-scale reaction was conducted using 7 mmol of **1a**, resulting in the formation of allenol **2a** with an isolated yield of 73% (1.1 g) (Fig. 4a). These allenols are versatile synthetic building blocks as shown in Fig. 4b. In the presence of Cs_2CO_3, allenol **2a** underwent oxycyclization to give the dihydropyran product **5**, which may further react with 4-chlorobenzyl bromide[12] to produce the dihydropyran product **6** in a one-pot reaction. The






Fig. 4 Gram-scale reaction and synthetic applications. [a] Reaction conditions: (a) **2a** (0.2 mmol), Cs_2CO_3 (1.0 equiv.), CH_3CN, rt, 12 h; (b) **2a** (0.2 mmol), 4-chlorobenzyl bromide (1.1 equiv.), TBAI (5 mol%), NaH (2.0 equiv.), THF, 0 °C to rt, Ar, 22 h; (c) **2a** (0.2 mmol), NBS (1.2 equiv.), DCM/THF = 4/1, rt, 12 h; (d) **2a** (0.2 mmol), $Fe(NO_3)_3{\cdot}9H_2O$ (20 mol%), TEMPO (20 mol%), NaCl (20 mol%), DCE (0.05 M), O_2, rt, 11 h; (e) (i) **2a** (0.2 mmol), $Fe(NO_3)_3{\cdot}9H_2O$ (20 mol%), TEMPO (20 mol%), KCl (20 mol%), DCE (0.2 M), O_2, rt, 30 h; (ii) $TMSCHN_2$ (1.5 equiv.), Et_2O/MeOH = 4/1, 0 °C to rt, 15 h; (f) **2a** (0.2 mmol), PPh_3 (1.5 equiv.), CBr_4 (1.5 equiv.), DCM, 0 °C to rt, 24 h; (g) **2a** (0.2 mmol), I_2 (1.2 equiv.), PPh_3 (1.2 equiv.), imidazole (1.2 equiv.), DCM, rt, 0.5 h; (h) **2a** (1.15 equiv.), *N*-(but-2-yn-1-yl)-4-methylbenzenesulfonamide (0.2 mmol), PPh_3 (1.35 equiv.), DIAD (1.35 equiv.), THF, 0 °C to rt, Ar, 23 h.

a) Radical inhibiting experiments




Radical inhibitor	NMR yield of **2a**	Recovery of **1a**	NMR yield of **S1a**
/	81%	0	/
TEMPO (2.0 equiv)	30%	24%	/
TEMPO (5.0 equiv)	2%	75%	/
BHT (2.0 equiv)	38%	13%	18% (18%)[a]
BHT (5.0 equiv)	18%	44%	29%

b) Control experiment



c) Stern-Volmer quenching experiments of Ir(ppy)$_3$



d) Cyclic voltammograms




Fig. 5 Mechanistic studies. [a] Isolated yield.

This journal is © the Partner Organisations 2025




Fig. 6 Proposed mechanism.

tetrahydrofuran product **7** was obtained through bromocyclization[13] with NBS in 84% yield. In addition, aerobic oxidation of **2a** afforded allenal **8** under the catalysis of 20 mol% each of Fe$(NO_3)_3{\cdot}9H_2O$, TEMPO, and NaCl in DCE.[14] Meanwhile, under the catalytic system of Fe$(NO_3)_3{\cdot}9H_2O$, TEMPO, and KCl, **2a** was converted into carboxylic acid, which was further esterified with $TMSCHN_2$ in one pot to provide allenoate **9** in 66% yield over two steps.[15] Upon treatment with CBr_4 and I_2, the allenol product **2a** could be easily transformed into the synthetically versatile allenyl bromide **10** and allenyl iodide **11**, respectively.[16,17] Finally, **2a** may readily undergo a Mitsunobu reaction[18] to afford allenyne **12** in a decent yield.

To gain mechanistic insights into the reaction, firstly, we conducted the reaction of **1a** in the presence of radical trapping agents such as TEMPO and BHT. As shown in Fig. 5a, the yield of the desired allenols dropped dramatically with the increase in the amount of the radical inhibitor. In addition, when BHT was used, the alkynol product **S1a**, resulting from intermolecular HAT of the initially formed alkoxyl radical, was detected. When the internal alkyne **1ah** substituted with the triisopropylsilyl group was subjected to the standard conditions, in addition to the normal allene product **2a**, the isolation of the tetrahydrofuranyl triisopropylsilyl acetylene product **13ah** indicated the involvement of a propargylic cation, which may be formed *via* the oxidation of a propargylic radical intermediate (Fig. 5b). Moreover, Stern–Volmer quenching experiments (Fig. 5c) demonstrated that the CuCl/1,10-phen catalyst, rather than the substrate **1a**, quenched the excited-state photocatalyst Ir$(ppy)_3$, suggesting a reductive quenching pathway. The half peak potential of the catalyst CuCl/1,10-phen $E^{red}_{p/2}$ [Cu^{II}/Cu^{I}] was determined by Cyclic Voltammetry (CV) experiments to be +0.26 V *vs.* SCE in CH_3CN and the half peak potential of substrate **1a** $E^{red}_{p/2}$ [**1a**/**1a**$^{\bullet-}$] was determined to be −1.27 V *vs.* SCE in CH_3CN (Fig. 5d).

On the basis of these experimental results, a plausible reaction mechanism was proposed (Fig. 6). Initially, the excitation of the photocatalyst Ir$(ppy)_3$ leads to its excited state Ir*$(ppy)_3$ ($E^{red}_{1/2}$ [Ir^{III*}/Ir^{II}] = +0.31 V *vs.* SCE in CH_3CN),[19] which can be reductively quenched with $LCu^{I}CN$ ($E^{red}_{p/2}$ [Cu^{II}/Cu^{I}] = +0.26 V *vs.* SCE in CH_3CN) to generate $LCu^{II}(CN)$ and [Ir$(ppy)_3]^-$ species. [Ir$(ppy)_3]^-$ ($E^{red}_{1/2}$ [Ir^{III}/Ir^{II}] = −2.19 V *vs.* SCE in CH_3CN)[19] would reduce substrate **1** ($E^{red}_{p/2}$ [**1a**/**1a**$^{\bullet-}$] = −1.27 V *vs.* SCE), producing radical anion intermediate **I** and regenerating the photocatalyst Ir$(ppy)_3$. Subsequently, alkoxy radical intermediate **II** would be generated *via* the cleavage of the N–O bond of intermediate **I**. The subsequent intramolecular 1,5-HAT would generate radical intermediate **III**. Mulliken atomic spin density analysis suggests that intermediate **III** is a delocalized radical, with the spin density distributed across both ends of the system (0.763 at C1 and 0.487 at C2, with R^1 being *n*-hexyl and R^2 being methyl). Finally, intermediate **III** can be trapped by the $LCu^{II}(CN)_2$ species, leading to the formation of the allene product **2**.

Conclusions

In summary, we have developed a unified platform for the highly selective preparation of tri- and tetra-substituted allenols. This approach leverages dual copper and photoredox catalysis to enable the generation and functionalization of allenyl radicals. This mild reaction demonstrates a broad substrate scope, and the post-reaction transformations of the allenol products highlight the significant synthetic potential of this methodology.

Author contributions

S. M., H. Y. and W. L. designed the experiments. W. L. performed the experiments. W. L., H. Y. and S. M. wrote the manuscript. W. L., H. Y. and S. M. checked the experimental data.

Data availability

All detailed procedures, characterization data, computational details, and spectra are available in the ESI.†

Conflicts of interest

The authors declare no competing interests.

Acknowledgements

Financial support from the National Key R&D Program of China (Grant No. 2022YFA1503200 for S. Ma and H. Yu) and the Shanghai Sailing Program (20YF1402100 for H. Yu) is greatly appreciated. We thank Mr Yaqi Shi in this group for reproducing the results of **2g** and **2l** in Fig. 2. We thank Mr Dong Huang and Dr Xue Zhang in this group for the Mulliken atomic spin density analysis and their helpful discussion.

References

1 (*a*) Z. Huang and G. Dong, Site-selectivity control in organic reactions: A quest to differentiate reactivity among the same kind of functional groups, *Acc. Chem. Res.*, 2017, **50**, 465; (*b*) L. M. Stateman, K. M. Nakafuku and D. A. Nagib, Remote C–H functionalization via selective hydrogen atom transfer, *Synthesis*, 2018, 1569.

2 For a recent review on the synthesis of alkynes *via* propargylic and allenylic radicals, see: (*a*) F. Wang, P. Chen and G. Liu, Copper-catalysed asymmetric radical cyanation, *Nat. Synth.*, 2022, **1**, 107. For selected recent examples, see: (*b*) F. Lu, D. Liu, L. Zhu, L. Lu, Q. Yang, Q. Zhou, Y. Wei, Y. Lan and W. Xiao, Asymmetric propargylic radical cyanation enabled by dual organophotoredox and copper Catalysis, *J. Am. Chem. Soc.*, 2019, **141**, 6167; (*c*) Y. Deng, R. Lu, P. Chen and G. Liu, Enantioselective cyanation of propargylic C–H bonds via cooperative photoredox and copper catalysis, *Chem. Commun.*, 2023, **59**, 4656; (*d*) J. Xia, X. Zhu and H. Bao, Copper-catalyzed enantioselective radical esterification of propargylic C–H bonds, *Synthesis*, 2024, 1393; (*e*) S. Lu, R. Agata, S. Nomura, H. Matsuda, K. Isozaki and M. Nakamura, Regioselective propargylic Suzuki–Miyaura coupling by SciPROP-Iron catalyst, *J. Org. Chem.*, 2024, **89**, 8385.

3 For reviews, see: (*a*) L. Fu, S. Greßies, P. Chen and G. Liu, Recent advances and perspectives in transition metal-catalyzed 1,4-functionalizations of unactivated 1,3-enynes for the synthesis of allenes, *Chin. J. Chem.*, 2020, **38**, 91; (*b*) Y. Li and H. Bao, Radical transformations for allene synthesis, *Chem. Sci.*, 2022, **13**, 8491.

4 For a selection of recent reports on the formation of allenyl radicals through radical addition to 1,3-enynes, see: (*a*) F. Wang, D. Wang, Y. Zhou, L. Liang, H. Lu, P. Chen, Z. Lin and G. Liu, Divergent synthesis of CF_3-substituted allenyl nitriles by ligand-controlled radical 1,2- and 1,4-addition to 1,3-enynes, *Angew. Chem., Int. Ed.*, 2018, **57**, 7140; (*b*) X. Zhu, W. Deng, M.-F. Chiou, C. Ye, W. Jian, Y. Zeng, Y. Jiao, L. Ge, Y. Li, X. Zhang and H. Bao, Copper-catalyzed radical 1,4-difunctionalization of 1,3-enynes with alkyl diacyl peroxides and *N*–fluorobenzenesulfonimide, *J. Am. Chem. Soc.*, 2019, **141**, 548; (*c*) K. Zhang, K. Bian, C. Li, J. Sheng, Y. Li and X. Wang, Nickel-catalyzed carbofluoroalkylation of 1,3-enynes to access structurally diverse fluoroalkylated allenes, *Angew. Chem., Int. Ed.*, 2019, **58**, 5069; (*d*) C. Ye, Y. Li, X. Zhu, S. Hu, D. Yuan and H. Bao, Copper-catalyzed 1,4-alkylarylation of 1,3-enynes with masked alkyl electrophiles, *Chem. Sci.*, 2019, **10**, 3632; (*e*) Y. Zeng, M.-F. Chiou, X. Zhu, J. Cao, D. Lv, W. Jian, Y. Li, X. Zhang and H. Bao, Copper-catalyzed enantioselective radical 1,4-difunctionalization of 1,3-enynes, *J. Am. Chem. Soc.*, 2020, **142**, 18014; (*f*) X. Dong, T. Zhan, S. Jiang, X. Liu, L. Ye, Z. Li, Q. Gu and X. Liu, Copper-catalyzed asymmetric coupling of allenyl radicals with terminal alkynes to access tetrasubstituted allenes, *Angew. Chem., Int. Ed.*, 2021, **60**, 2160; (*g*) Y. Chen, J. Wang and Y. Lu, Decarboxylative 1,4-carbocyanation of 1,3-enynes to access tetra-substituted allenes *via* copper/photoredox dual catalysis, *Chem. Sci.*, 2021, **12**, 11316; (*h*) L. Wang, R. Ma, J. Sun, G. Zheng and Q. Zhang, NHC and visible light-mediated photoredox co-catalyzed 1,4-sulfonylacylation of 1,3-enynes for tetrasubstituted allenyl ketones, *Chem. Sci.*, 2022, **13**, 3169; (*i*) H. Chen, C. Zhu, H. Yue and M. Rueping, Group 14 elements hetero-difunctionalizations *via* nickel-catalyzed electroreductive cross-coupling, *Angew. Chem., Int. Ed.*, 2023, **62**, e202306498; (*j*) C. Kuai, B. Teng and X. Wu, Palladium-catalyzed carbonylative multicomponent fluoroalkylation of 1,3-enynes: Concise construction of diverse cyclic compounds, *Angew. Chem., Int. Ed.*, 2024, **63**, e202318257; (*k*) Q. Shan, Y. Zhao, S. Wang, H. Liu, X. Duan and L. Guo, Nickel-catalyzed modular four-component 1,4-alkylcarbonylation of 1,3-enynes to tetra-substituted CF_3–allenyl ketones, *ACS Catal.*, 2024, **14**, 2144; (*l*) Y. Pu, S. Ding, H. Zhao, Q. Xue, H. Zhang, X. Xie, Y. Shang and J. Wang, Three-component synthesis of multiple functionalized allenes *via* copper/photoredox dual catalyzed 1,4-alkylcyanation of 1,3-enynes, *Org. Lett.*, 2024, **26**, 1834; (*m*) Y. Chen, Q. Wang, K. Peng, L. Cheng, Z. She and G. Deng, Copper/photoredox-catalyzed regioselective 1,4-addition to 1,3-enynes: A convenient access to perfluoroalkylated allenes, *Chin. J. Chem.*, 2024, **42**, 2023; (*n*) J. Zhang, J. Xiong, L. Fei, F. Cui, S. Li, H. Tang, Y. Xie and Y. Pan, Electrochemically mediated synthesis of trifluoromethylallenes, *Org. Chem. Front.*, 2024, **11**, 3485; (*o*) J. H. Jeon, G. H. Kim, H. S. Lee, D. H. Kim, S. Lee, W. Choe, B. Jung, J. Rohde and S. Y. Hong, Chemo- and regioselective nickel-catalyzed reductive 1,4-alkylarylation of 1,3-enynes through an L_2NiAr intermediate, *ACS Catal.*, 2024, **14**, 8996.

5 For a selection of recent reports on the formation of allenyl radicals from alkynes, see: (*a*) R. Soler-Yanes, I. Arribas-

 This journal is © the Partner Organisations 2025

Álvarez, M. Guisán-Ceinos, E. Buñuel and D. J. Cárdenas, NiI catalyzes the regioselective cross-coupling of alkylzinc halides and propargyl bromides to allenes, *Chem. – Eur. J.*, 2017, **23**, 1584; (*b*) P. Domingo-Legarda, R. Soler-Yanes, M. T. Quirós-López, E. Buñuel and D. J. Cárdenas, Iron-catalyzed coupling of propargyl bromides and alkyl grignard reagents, *Eur. J. Org. Chem.*, 2018, 4900; (*c*) F. W. Friese and A. Studer, Deoxygenative borylation of secondary and tertiary alcohols, *Angew. Chem., Int. Ed.*, 2019, **58**, 9561; (*d*) R. Lu, T. Yang, X. Chen, W. Fan, P. Chen, Z. Lin and G. Liu, Enantioselective copper-catalyzed radical cyanation of propargylic C−H bonds: easy access to chiral allenyl nitriles, *J. Am. Chem. Soc.*, 2021, **143**, 14451; (*e*) W. Zhang, L. Lu, W. Zhang, Y. Wang, S. D. Ware, J. Mondragon, J. Rein, N. Strotman, D. Lehnherr, K. A. See and S. Lin, Electrochemically driven cross-electrophile coupling of alkyl halides, *Nature*, 2022, **604**, 292; (*f*) Q. Liu, J. Zheng, X. Zhang and S. Ma, Photo and copper dual catalysis for allene syntheses from propargylic derivatives via one-electron process, *Nat. Commun.*, 2022, **13**, 3302; (*g*) F. Lu, Z. Shu, G. He, J. Bai, L. Lu and W. Xiao, Photoinduced C–O bond cleavage for copper-catalyzed allenyl radical cyanation, *Org. Chem. Front.*, 2022, **9**, 5259; (*h*) D. Zhang, J. Fan, Y. Shi, Y. Huang, C. Fu, X. Wu and S. Ma, Copper-catalyzed propargylic C–H functionalization for allene syntheses, *Chem. Sci.*, 2023, **14**, 9191; (*i*) J. Qin, Z. Xiong, C. Cheng, M. Hu and J. Li, Electroreductive carboxylation of propargylic acetates with CO_2: access to tetrasubstituted 2,3-allenoates, *Org. Lett.*, 2023, **25**, 9176.

6 For a selection of recent reports on the formation of allenyl radicals from allenes, see: (*a*) Y. Wei, X. Wu, Y. Chen and C. Zhu, Radical-polar crossover allenylation of alkenes: divergent synthesis of homoallenic alcohols and amides, *Chem. Catal.*, 2023, **3**, 100551; (*b*) Z. Cheng, T. Yang, C. Li, Y. Deng, F. Zhang, P. Chen, Z. Lin, S. Ma and G. Liu, Site-selective sp^2 C−H cyanation of allenes via copper-catalyzed radical relay, *J. Am. Chem. Soc.*, 2023, **145**, 25995; (*c*) Y. Wei, Z. Wang, K. Wang, J. Qiu, Z. Wang, H. Li, X. Duan, K. Guo, X. Bao and X. Wu, Copper-catalyzed radical allene C(sp^2)−H cyanation, *Angew. Chem., Int. Ed.*, 2024, **63**, e202317132.

7 For selected reviews on 1,5-HAT, see: (*a*) Ž. Čeković, Reactions of carbon radicals generated by 1,5-transposition of reactive centers, *J. Serb. Chem. Soc.*, 2005, **70**, 287; (*b*) S. Chiba and H. Chen, Sp^3 C–H oxidation by remote H-radical shift with oxygen- and nitrogen-radicals: a recent update, *Org. Biomol. Chem.*, 2014, **12**, 4051; (*c*) L. M. Stateman, K. M. Nakafuku and D. A. Nagib, Remote C–H functionalization *via* selective hydrogen atom transfer, *Synthesis*, 2018, 1569; (*d*) N. Goswami and D. Maiti, An update on distal C(sp^3)-H functionalization involving 1,5-HAT emerging from nitrogen radicals, *Isr. J. Chem.*, 2020, **60**, 303; (*e*) X. Wu and C. Zhu, Radical functionalization of remote C(sp^3)–H bonds mediated by unprotected alcohols and amides, *CCS Chem.*, 2020, **2**, 813; (*f*) H. Chen and S. Yu, Remote C–C bond formation via visible light photoredox-catalyzed intramolecular hydrogen atom transfer, *Org. Biomol. Chem.*, 2020, **18**, 4519; (*g*) S. Sarkar, K. P. S. Cheung and V. Gevorgyan, C–H functionalization reactions enabled by hydrogen atom transfer to carbon-centered radicals, *Chem. Sci.*, 2020, **11**, 12974; (*h*) W. Li and C. Zhu, *Radically initiated distal C(sp^3)–H functionalization*, Wiley-VCH, Weinheim, 2021, pp. 315–341; (*i*) W. Guo, Q. Zhang and J. Zhu, Visible light photoredox-catalysed remote C–H functionalisation enabled by 1,5-hydrogen atom transfer (1,5-HAT), *Chem. Soc. Rev.*, 2021, **50**, 7359; (*j*) H. Liu, Q. Jin and X. Wu, Intramolecular hydrogen atom transfer from unactivated C(sp^3)-H bond to alkyl radical, *Eur. J. Org. Chem.*, 2024, e202301280.

8 For some seminal reports on 1,5-HAT, see: (*a*) A. W. Hofmann, Ueber die Einwirkung des Broms in alkalischer Lösung auf die Amine, *Ber. Dtsch. Chem. Ges.*, 1883, **16**, 558; (*b*) D. H. R. Barton, The use of photochemical reactions in organic sunthesis, *Pure Appl. Chem.*, 1968, **16**, 1. For selected recent reports on 1,5-HAT reactions, see: (*c*) H. Jiang and A. Studer, α-Aminoxy-acid-auxiliary-enabled intermolecular radical γ-C(sp^3)-H functionalization of ketones, *Angew. Chem., Int. Ed.*, 2018, **57**, 1692; (*d*) J. Liu, Y. Wei and M. Shi, Direct activation of a remote C(sp^3)–H bond enabled by a visible-light photosensitized allene moiety, *Angew. Chem., Int. Ed.*, 2021, **60**, 12053; (*e*) K. M. Nakafuku, Z. Zhang, E. A. Wappes, L. M. Stateman, A. D. Chen and D. A. Nagib, Enantioselective radical C–H amination for the synthesis of β-amino alcohols, *Nat. Chem.*, 2020, **12**, 697; (*f*) L. Zhong, Z. Xiong, X. Ouyang, Y. Li, R. Song, Q. Sun, X. Lu and J. Li, Intermolecular 1,2-difunctionalization of alkenes enabled by fluoroamide-directed remote benzyl C(sp^3)−H functionalization, *J. Am. Chem. Soc.*, 2022, **144**, 339; (*g*) Y. Zhou, Y. Qin, Q. Wang, Z. Zhang and G. Zhu, Photocatalytic sulfonylcarbocyclization of alkynes using SEt as a traceless directing group: Access to cyclopentenes and indenes, *Angew. Chem., Int. Ed.*, 2022, **61**, e202110864; (*h*) H. Liu, X. Kong, X. Gong, Y. Li, Z. Niu, X. Gou, X. Li, Y. Wang, W. Shi, Y. Huang, X. Liu and Y. Liang, Site-selective coupling of remote C(sp^3)–H/meta-C(sp^2)–H bonds enabled by Ru/photoredox dual catalysis and mechanistic studies, *Chem. Sci.*, 2022, **13**, 5382; (*i*) R. Guo, H. Xiao, S. Li, Y. Luo, J. Bai, M. Zhang, Y. Guo, X. Qi and G. Zhang, Photoinduced copper-catalyzed asymmetric C(sp^3)-H alkynylation of cyclic amines by intramolecular 1,5-hydrogen atom transfer, *Angew. Chem., Int. Ed.*, 2022, **61**, e202208232; (*j*) S. Chen, B. Yuan, Y. Wang and L. Ackermann, Ruthenium-catalyzed remote difunctionalization of nonactivated alkenes for double meta-C(sp^2)–H/C-6(sp^3)–H functionalization, *Angew. Chem., Int. Ed.*, 2023, **62**, e202301168; (*k*) W. Li, B. Sun, L. Zhang and F. Mo, Photocarboxylation of remote C–H bonds through nitrogen-centred radical 1,5-hydrogen atom transfer, *Green Chem.*, 2023, **25**, 5030; (*l*) J. Wang, Y. Zhang, L. Zhu, X. Xue and C. Li, Photoinduced remote C(sp^3)–H phosphonylation

of amides, *Angew. Chem., Int. Ed.*, 2024, **63**, e202400856; (*m*) Y. Zhou, K. He, W. Zheng, J. Lv, S. He, N. Yu, Y. Yang, L. Liu, K. Jiang and Y. Wei, 1,5-Hydrogen atom transfer of α-iminyl radical cations: A new platform for relay annulation for pyridine derivatives and axially chiral heterobiaryls, *Chem. Sci.*, 2024, **15**, 7502.

9 For selected reviews on alkoxyl radicals, see: (*a*) K. Jia and Y. Chen, Visible-light-induced alkoxyl radical generation for inert chemical bond cleavage/functionalization, *Chem. Commun.*, 2018, **54**, 6105; (*b*) E. Tsui, H. Wang and R. R. Knowles, Catalytic generation of alkoxy radicals from unfunctionalized alcohols, *Chem. Sci.*, 2020, **11**, 11124; (*c*) L. Chang, Q. An, L. Duan, K. Feng and Z. Zuo, Alkoxy radicals see the light: New paradigms of photochemical synthesis, *Chem. Rev.*, 2022, **122**, 2429; (*d*) A. S. Budnikov, I. B. Krylov, A. V. Lastovko, B. Yu and A. O. Terent'ev, Copper/photoredox dual catalyzed syntheses of allenenitriles via 1,5-HAT, *Asian J. Org. Chem.*, 2022, **11**, e202200262; (*e*) J. Liu, J. Jin, K. Zhang and L. Cai, Alkoxy radical induced transformations from *N*-alkoxyphthalimides under the photoredox catalysis, *Asian J. Org. Chem.*, 2023, **12**, e202300160; (*f*) Y. Zhu, H. Wang, P. Shu, K. Zhang and Q. Wang, Recent advances on alkoxy radicals-mediated C(sp^3)–H bond functionalization via 1,5-hydrogen atom transfer, *Chin. J. Org. Chem.*, 2024, **44**, 1.

10 For selected recent reports on 1,5-HAT reactions mediated by alkoxyl radicals, see: (*a*) J. Zhang, Y. Li, F. Zhang, C. Hu and Y. Chen, Generation of alkoxyl radicals by photoredox catalysis enables selective C(sp^3)–H functionalization under mild reaction conditions, *Angew. Chem., Int. Ed.*, 2016, **55**, 1872; (*b*) A. Hu, J. Guo, H. Pan, H. Tang, Z. Gao and Z. Zuo, *δ*–Selective functionalization of alkanols enabled by visible-light-induced ligand-to-metal charge transfer, *J. Am. Chem. Soc.*, 2018, **140**, 1612; (*c*) Y. Zhu, K. Huang, J. Pan, X. Qiu, X. Luo, Q. Qin, J. Wei, X. Wen, L. Zhang and N. Jiao, Silver-catalyzed remote Csp^3-H functionalization of aliphatic alcohols, *Nat. Commun.*, 2018, **9**, 2625; (*d*) X. Wu, H. Zhang, N. Tang, Z. Wu, D. Wang, M. Ji, Y. Xu, M. Wang and C. Zhu, Metal-free alcohol-directed regioselective heteroarylation of remote unactivated C(sp^3)–H bonds, *Nat. Commun.*, 2018, **9**, 3343; (*e*) H. Guan, S. Sun, Y. Mao, L. Chen, R. Lu, Ji. Huang and L. Liu, Iron(II)-catalyzed site-selective functionalization of unactivated C(sp^3)-H Bonds guided by alkoxyl radicals, *Angew. Chem., Int. Ed.*, 2018, **57**, 11413; (*f*) Y. Li, Y. Bi, W. Ke, C. Wang, D. Hu, Y. Li and X. Wang, Copper/photoredox dual catalyzed syntheses of allenenitriles via 1,5-HAT, *Asian J. Org. Chem.*, 2023, **12**, e202300234; (*g*) G. Li, X. Hu, G. He and G. Chen, Photoredox-mediated remote C(sp^3)–H heteroarylation of free alcohols, *Chem. Sci.*, 2019, **10**, 688; (*h*) X. Bao, Q. Wang and J. Zhu, Dual photoredox/copper catalysis for the remote C(sp^3)-H functionalization of alcohols and alkyl halides by *N*-alkoxypyridinium salts, *Angew. Chem., Int. Ed.*, 2019, **58**, 2139; (*i*) A. N. Herron, D. Liu, G. Xia and J. Yu, *δ*–C–H mono- and dihalogenation of alcohols, *J. Am. Chem. Soc.*, 2020, **142**, 2766; (*j*) N. Xiong, Y. Li and R. Zeng, Iron-catalyzed photoinduced remote C(sp^3)–H amination of free alcohols, *Org. Lett.*, 2021, **23**, 8968; (*k*) Z. Jue, Y. Huang, J. Qian and P. Hu, Visible light-induced unactivated δ-C(sp^3)-H amination of alcohols catalyzed by iron, *ChemSusChem*, 2022, **15**, e202201241; (*l*) L. Huang, J. Sun, B. Sun, S. Song and J. Li, Regioselective synthesis of isoquinolinonediones through remote unactivated C(sp^3)–H bonds, *Chem. Commun.*, 2024, **60**, 4818; (*m*) D. S. Finis and D. A. Nicewicz, Alkoxy radical generation mediated by sulfoxide cation radicals for alcohol-directed aliphatic C–H functionalization, *J. Am. Chem. Soc.*, 2024, **146**, 16830.

11 For reviews on radical additions to allenes or alkynes, see: (*a*) L. Liu, R. M. Ward and J. M. Schomaker, Mechanistic aspects and synthetic applications of radical additions to allenes, *Chem. Rev.*, 2019, **119**, 12422; (*b*) G. Qiu, J. Zhang, K. Zhou and J. Wu, Recent advances in the functionalization of allenes via radical process, *Tetrahedron*, 2018, **74**, 7290; (*c*) C. Hu, J. Mena and I. V. Alabugin, Design principles of the use of alkynes in radical cascades, *Nat. Rev. Chem.*, 2023, **7**, 405; (*d*) M. Huang, W. Hao, G. Li, S. Tu and B. Jiang, Recent advances in radical transformations of internal alkynes, *Chem. Commun.*, 2018, **54**, 10791.

12 C. Li, Z. Yang, L. Wang, Y. Guo, Z. Huang and S. Ma, Cobalt-catalyzed regio- and stereoselective hydroboration of allenes, *Angew. Chem., Int. Ed.*, 2020, **59**, 6278.

13 J. Li, C. Fu, G. Chen, G. Chai and S. Ma, Studies on electrophilic interaction of 2,3-allenols with electrophilic halogen reagents: Selective synthesis of 2,5-dihydrofurans, 3-halo-3-alkenals, or 2-halo-2-alkenyl ketones, *Adv. Synth. Catal.*, 2008, **350**, 1376.

14 S. Ma, J. Liu, S. Li, B. Chen, J. Cheng, J. Kuang, Y. Liu, B. Wan, Y. Wang, J. Ye, Q. Yu, W. Yuan and S. Yu, Development of a general and practical iron nitrate/TEMPO-catalyzed aerobic oxidation of alcohols to aldehydes/ketones: Catalysis with table salt, *Adv. Synth Catal.*, 2011, **353**, 1005.

15 X. Jiang, J. Zhang and S. Ma, Iron catalysis for room-temperature aerobic oxidation of alcohols to carboxylic acids, *J. Am. Chem. Soc.*, 2016, **138**, 8344.

16 N. Xiong, Y. Li and R. Zeng, Iron-catalyzed photoinduced remote C(sp^3)–H amination of free alcohols, *Org. Lett.*, 2021, **23**, 8968.

17 D. Ma, X. Duan, C. Fu, X. Huang and S. Ma, Dimethylprolinol versus diphenylprolinol in $CuBr_2$-catalyzed enantioselective allenylation of terminal alkynols, *Synthesis*, 2018, 2533.

18 Y. Zhang, W. Xu, T. Gao, M. Guo, C. Yang, H. Xie, X. Kong, Z. Yang and J. Chang, Pd-catalyzed borylsilylative cyclization of 1,6-allenynes, *Org. Lett.*, 2022, **24**, 7021.

19 J.-H. Shon, D. Kim, M. D. Rathnayake, S. Sittel, J. Weaver and T. S. Teets, Photoredox catalysis on unactivated substrates with strongly reducing iridium photosensitizers, *Chem. Sci.*, 2021, **12**, 4069.

 This journal is © the Partner Organisations 2025