CJC Chinese Journal of Chemistry 中国化学 · An International Journal

For submission: https://wiley.atyponrex.com/journal/CJOC
For published articles: https://onlinelibrary.wiley.com/journal/16147065

Concise Report

Cite this paper: *Chin. J. Chem.* **2025**, *43*, 1181—1189. **DOI: 10.1002/cjoc.202401301**

A Divergent Synthesis of Spiroindenes via Ligand-Controlled [3+2]/[4+2] Cycloadditions of Zwitterionic π-Propargyl Palladium Species with Benzofulvenes


Yongjie Long, Xianhua Zhong, Min Shi,* and Yin Wei*

State Key Laboratory of Organometallic Chemistry, Center for Excellence in Molecular Synthesis, University of Chinese Academy of Sciences, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China

Keywords

Zwitterions | Palladium catalyzed | Divergent synthesis | Ligand control | Spiro compounds | π-Propargyl palladium species | Annulation

Comprehensive Summary



A divergent synthesis of spiroindenes through a palladium catalyzed cycloaddition between zwitterionic π-propargyl palladium species and benzofulvenes in moderate to good yields has been disclosed along with good functional group compatibility and a broad substrate universality. This protocol features a highly regioselective switchable process between [3+2] and [4+2] cycloadditions controlled by phosphine ligands with different bite angles. The reaction mechanism has been clarified by mechanistic studies and DFT calculations, rendering that the coordination modes of the ligands with the substrates and the bite angle of the ligands play critical roles in the product regioselectivity.

*E-mail: mshi@mail.sioc.ac.cn; weiyin@sioc.ac.cn

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

Background and Originality Content

Spirocyclic compounds occupy an important position in the field of chemical biology and pharmaceutical science.[1] In the vast family of spirocylic compounds, spiroindenes, widespread likewise spirocyclohexadienones or spiroindolones, are gradually revealing their significant potential in natural product chemistry,[2] drug synthesis[3] and asymmetric catalysis[4] (Figure 1). However, comparing with their immense potential, the construction of spiroindene skeleton is still running in traditional pathways, which need either long-winded procedures or complicated materials, such as sequential cyclization of bisphenols or 1,5-diphenyl-3-pentanone derivatives promoted by Brønsted or Lewis acid,[5] cascade dearomatization between alkynes and 1-aryl-2-naphthols[6] or intramolecular cyclization from complex substrates such as indene-substituted cyclobutanols and 2-phenethyl cyclic ketones.[7] Hence, more advanced synthetic strategies should be unearthed to realize various types of spiroindenes efficiently and concisely.





Figure 1 Natural products, bioactive molecules and ligands containing the spiroindene or spiroindane core.

As a kind of synthon having semi-aromatic characteristics, benzofulvenes have been mainly employed in the synthesis of polymeric compounds and transition-metal-complexes.[8] Based on the same indene structure, benzofulvenes could be transformed to spiroindenes in certain ways. In 2016, Jørgensen's group utilized benzofulvenes initiatively to form enantiomerically enriched polysubstituted spiro[5,6]indenes or spiro[5,3]indenes induced by chiral diphenylprolinol silylether[9] or chiral cinchona alkaloid catalysts,[10] which marked the milestone of spiroindene skeleton being formed tersely from benzofulvenes in one step. Nevertheless, although Wang's group combined azomethine ylides and benzofulvenes together to construct heterospiro[5,5]indenes in 2023 (Scheme 1, a),[11] the development of benzofulvenes to spiroindene's synthetic protocol has nearly stagnated. Thus far, each method only has the ability to build up one type of spiroindenes. Therefore, there is still lack of a diversified synthetic method in this field.

On the other hand, zwitterionic π-propargyl palladium species, similar as zwitterionic π-allyl palladium species, have already emerged its synthetic capacity in various cycloadditions.[12] Depending on the adjustment of bidentate ligand with different bite angles, the electrophilic point of π-propargyl palladium part was shifted between *3*-terminal carbon and 2-internal carbon (Scheme 1, b).[12] In continuation of our research interest in zwitterionic π-propargyl palladium species related chemistry, we supposed that the indene part of benzofulvene was stabilized by the aromatic cyclopentadienyl anion after the nucleophilic attack from zwitterionic π-propargyl palladium species, providing a driving force for the subsequent cyclization. Thus, we anticipate that a divergent synthesis of spiroindenes could be achieved upon using bidentate phosphine ligands having different bite angles as the key controlling point (Scheme 1, c).

Scheme 1 [a] Previous synthesis of spiroindenes via benzofulvenes. [b] Reaction mode of zwitterionic π-palladium species adjusted by ligands. [c] Design of this work.




Results and Discussion

We first utilized cyan substituted benzofulvene **1a** (1.0 eq) and vinylidenecyclopropane-diester (VDCP-diester) **2a** (2.0 eq) as the model substrates to perform the initial investigation and subsequently optimize the reaction conditions of [3+2] cycloaddition (Table 1). As a result, we found that the desired product **3a** was obtained in 69% NMR yield by using Pd_2dba_3 as a palladium catalyst, DPEphos as a ligand, $Yb(OTf)_3$ as the cocatalyst, Cs_2CO_3 as a base and upon heating at 55 °C in THF for 4.0 h with 4 Å MS (100 mg) (Table 1, entry 1). However, the polarity of **3a** is too close to the byproduct dibenzylideneacetone derived from Pd_2dba_3, which makes it difficult to separate these two compounds. Hence, the source of palladium catalyst was switched to $Pd(PPh_3)_4$, affording **3a** in 71% yield (Table 1, entry 2). The influence of solvent was investigated, discovering that using toluene, DCE, or DMF as solvent decreased the yield of **3a** to 41%, 52% and 48%, respectively, and nearly trace in 1,4-dioxane (Table 1, entries 3—6). The effect of different phosphine ligands was examined afterward. When other ligands with a large bite angle, such as dtpbf, Xantphos and BINAP were used to replace DPEphos, the yield of **3a** decreased to 43%, 30% and nearly zero (Table 1, entries 7, 9, 10).

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Table 1 Optimization of the reaction conditions for the synthesis of **3a**




entry[a]	ligand	solvent	base	*T*/°C	*t*/h	yield/%
1[b]	DPEphos	THF	Cs_2CO_3	55	4	69
2	DPEphos	THF	Cs_2CO_3	55	4	71
3	DPEphos	toluene	Cs_2CO_3	55	4	41
4	DPEphos	DCE	Cs_2CO_3	55	4	52
5	DPEphos	1,4-dioxane	Cs_2CO_3	55	4	trace
6	DPEphos	DMF	Cs_2CO_3	55	4	48
7	BINAP	THF	Cs_2CO_3	55	4	trace
8	dppe	THF	Cs_2CO_3	55	4	trace
9	dtpbf	THF	Cs_2CO_3	55	4	43
10	Xantphos	THF	Cs_2CO_3	55	4	30
11	DPEphos	THF	TEA	55	4	38
12	DPEphos	THF	DBU	55	4	trace
13	DPEphos	THF	K_3PO_4	55	4	N.R.
14	DPEphos	THF	CsOAc	55	4	65
15	**DPEphos**	**THF**	**Cs_2CO_3**	**55**	**8**	**78 (77[c])**
16	DPEphos	THF	Cs_2CO_3	55	12	65
17	DPEphos	THF	Cs_2CO_3	60	4	69
18	DPEphos	THF	Cs_2CO_3	60	8	70
19	DPEphos	THF	Cs_2CO_3	45	8	49
20[d]	DPEphos	THF	Cs_2CO_3	55	8	65
21[e]	DPEphos	THF	Cs_2CO_3	55	8	73

[a] Reactions were carried out with **1a** (0.1 mmol), **2a** (0.2 mmol), [Pd] (5 mol%), ligand (7.5 mol%), $Yb(OTf)_3$ (10 mol%), 4 Å MS (100 mg), Cs_2CO_3 (0.1 mmol) in solvents (1.0 mL). Yields were determined by NMR spectroscopy using CH_2Br_2 as an internal standard. [b] Pd_2dba_3 instead of $Pd(PPh_3)_4$. [c] Isolated yield. [d] 0.15 mmol **2a**. [e] 0.25 mmol **2a**.

Significantly, using dppe as a traditional bidentate ligand with a small bite angle only afforded trace of **3a** also without the formation of [4+2] cycloadduct (Table 1, entry 8). The exploration of base effect revealed that both organic bases such as TEA or DBU and inorganic bases such as K_3PO_4 or CsOAc did not give better performance than Cs_2CO_3 in this reaction (Table 1, entries 11—14). Next, we further examined the reaction temperature and time, identifying that the reaction should be carried out at 55 °C for 8.0 h, giving **3a** in 77% isolated yield. This result was better than those at 45 °C for 8.0 h or at 60 °C for 8.0 h, respectively, even if the yield of **3a** at 60 °C (70%) seems to be higher than that at 60 °C for 4.0 h (69%) (Table 1, entries 15—19). Finally, we also confirmed that decreasing or increasing the amount of VDCP-diester **2a** to 1.5 or 2.5 equivalents did not improve the yield of **3a** under otherwise identical conditions (Table 1, entries 20 & 21). It should be noted that the diastereoselectivity of this [3+2] cycloaddition is higher than 20 : 1, probably due to the steric hindrance derived from benzofulvene skeleton. Furthermore, a series of common Lewis acids including $SnCl_4$, $BF_3{\cdot}Et_2O$, $In(OTf)_3$, $Zn(OTf)_2$, $Y(OTf)_3$, $Sc(OTf)_3$, $Yb(OTf)_3$ were screened for this reaction; the $Yb(OTf)_3$ particularly demonstrated superior catalytic efficacy in mediating the [3+2] cycloaddition reaction when compared to other Lewis acid additives (see part **E** at the page S4 of the Supporting Information).

Table 2 Optimization of the reaction conditions for the synthesis of **4a**



entry[a]	ligand	solvent	base	*T*/°C	*t*/h	yield/%
1	(*R*)-Ph-PHOX	THF	Cs_2CO_3	45	6	trace
2	(*R*)-Ph-PHOX	THF	Cs_2CO_3	50	6	12
3	(*R*)-Ph-PHOX	THF	Cs_2CO_3	55	8	45
4	(*R*)-Ph-PHOX	THF	Cs_2CO_3	60	8	55
5	**(*R*)-Ph-PHOX**	**THF**	**Cs_2CO_3**	**65**	**8**	**78 (76[b])**
6	(*R*)-Ph-PHOX	THF	Cs_2CO_3	65	12	64
7	(*R*)-Ph-PHOX	THF	Cs_2CO_3	70	8	52
8	(*S,R*)-Duanphos	THF	Cs_2CO_3	65	8	59[c]
9	2Me-PHOX	THF	Cs_2CO_3	65	8	62[d]
10[e]	(*R*)-Ph-PHOX	THF	Cs_2CO_3	65	8	40
11[f]	(*R*)-Ph-PHOX	THF	Cs_2CO_3	65	8	64
12	(*R*)-Ph-PHOX	DCE	Cs_2CO_3	65	8	trace
13	(*R*)-Ph-PHOX	1,4-dioxane	Cs_2CO_3	65	8	trace
14	(*R*)-Ph-PHOX	toluene	Cs_2CO_3	65	8	13
15	(*R*)-Ph-PHOX	DMF	Cs_2CO_3	65	8	24

[a] Reactions were carried out with **1a** (0.1 mmol), **2a** (0.2 mmol), [Pd] (5 mol%), ligand (7.5 mol%), $Yb(OTf)_3$ (10 mol %), 4 Å MS (100 mg), Cs_2CO_3 (0.1 mmol) in solvents (1.0 mL). Yields were determined by NMR spectroscopy using CH_2Br_2 as an internal standard. [b] Isolated yield. [c] 5-Membered ring : 6-membered ring = 1 : 1. [d] 5-Membered ring : 6-membered ring = 0.7 : 1. [e] 0.15 mmol **2a**. [f] 0.25 mmol **2a**.

Based on the assumption of ligand-controlled regioselectivity, we attempted to find a ligand which can switch the reaction mode from [3+2] to [4+2] cycloaddition. Fortunately, (*R*)-Ph-PHOX was found out to realize the [4+2] cycloaddition mode. It should be mentioned here that although dppe as a traditional bidentate ligand having a small bite angle[12a] was a preferred bidentate phosphine ligand in the palladium-catalyzed [4+2] cycloaddition of **2a** with methyleneindolinone, it did not perform as a useful ligand in this reaction (Table 1, entry 8). The reaction temperature and time were investigated in the first place to optimize the reaction conditions of this [4+2] cycloaddition, revealing that the reaction outcome was better at 65 °C for 8.0 h rather than those at lower or higher temperatures for shorter or longer time, affording **4a** in 78% NMR yield along with 76% isolated yield (Table 2, entries 1—7). The use of other ligands having a small bite angle such as Duanphos or PHOX substituted with two methyl groups actually affected the regioselectivity of this reaction as well as the ratio of [4+2]/[3+2] products, and delivered **4a** in lower yields (Table 2, entries 8 & 9). In addition, using 1.5 or 2.5 equivalents of VDCP-diester **2a** did not improve the yield of **4a** under otherwise identical conditions as well (Table 2, entries 10 & 11). Moreover, the influence of solvent effect was explored, indicating that replacing THF with DCE or 1,4-dioxane inhibited the reaction, and altering to toluene or DMF decreased the yield of **4a** to 13% and 24%, respectively (Table 2, entries 12—15). The diastereomeric ratio of [4+2] cycloaddition product **4a** was also higher than 20 : 1

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH


16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

due to the steric effect of the employed substrates.

The exploration of substrate scopes on spiro[5,5]indenes **3** and spiro[5,6]indenes **4** was subsequently carried out based on their optimized reaction conditions. The results of [3+2] cycloadditions are displayed in Scheme 2; the ORTEP drawing of **3a** unambiguously identified by X-ray diffraction is also shown in Scheme 2. Introducing both electron-donating groups such as methyl, *tert*-butyl or methoxy and electron-withdrawing groups such as halogen atom, trifluoromethyl or nitryl at the *2,4,5,6,7*-position of indene ring afforded the corresponding products **3a**—**3j** in good to excellent yields ranging from 67% to 99%, suggesting that the electronic property of **1** did not have significant impact on the reaction outcome. The diastereoselectivity of **3j** was dropped to 8 : 1 on account of the steric effect from *2*-methyl group which may neutralize the steric effect from benzofulvene skeleton. Replacing the cyan group with carbomethoxy or formyl group also furnished the corresponding products **3k** and **3l** in the yields of 82% and 80%, while the substrates having the methyl group substitution for cyan group or without substitution did not undergo the reaction to afford the desired products **3m** and **3n**. In addition, condensed, polycyclic or even polysubstituted indene skeletons were also compatible in this reaction, producing the desired cycloadducts **3o**, **3p** and **3q** in good yields ranging from 74% to 80%. Besides, the shift from phenyl group to cinnamyl or heteroaryl group, except for pyridyl group, did not restrain the reaction, giving the desired products **3r**—**3v** in 72%—75% yields, while the diastereoselectivity of **3r** and **3s** decreased to 5 : 1 and 10 : 1 presumably due to the similar reason as that of **3j**. The introduction of *para*-methoxy, *para*-carbomethoxy, *para*-trifluoromethyl, *para*-bromo and *meta*/*ortho*-methyl group into the phenyl ring generated the corresponding products **3w**—**3ab** in good yields ranging from 74% to 77%. Furthermore, the [3+2] cycloaddition proceeded smoothly utilizing benzofulvenes with a carboethoxy group or a *tert*-butyl substituent, affording the desired products **3ac** and **3ad** in the yields of 74% and 36%, respectively. Moreover, we also examined the reaction outcome with VDCP-diesters **2** bearing different ester groups. The variation of diester's moiety slightly influenced the reaction results, delivering the corresponding products **3ae**—**3ag** in the yields of 67%—78%, while inserting an ethyl group to the terminal position of allene provided the desired product **3ah** in the yield of 76% with the diastereomeric ratio of 1.7 : 1, due to the introduction of an allenyl axial chirality.

The results of the study on the substrate universality of [4+2] cycloaddition are generally similar to the [3+2] cycloaddition mode, which are exhibited in Scheme 3 along with the ORTEP drawing of **4b** unequivocally determined by X-ray diffraction. When the DPEphos ligand with a large bite angle was replaced by the (*R*)-Ph-PHOX ligand with a small bite angle, the excellent diastereoselectivity was inherited and the regioselectivity on spiro[5,6]indene to spiro[5,5]indene ratio can be kept over 16 : 1 in most cases. Analogously, the introduction of electron-donating groups as well as electron-withdrawing group at the *4,5,6,7*-position of indene ring or varying diverse electron-withdrawing group at *1*-position of indene ring afforded the corresponding products **4a**—**4i** in moderate to good yields ranging from 44% to 76%. In the case of naphthobenzofulvene, the ratio of [4+2] cycloadduct **4h** and [3+2] cycloadduct as **3o** in Scheme 2 decreased to 10 : 1. Heteroaryl groups are compatible with this [4+2] cycloaddition, giving the desired products **4j** and **4k** with 2-furanyl or 2-thienyl moiety in 64% and 62% yields, although their diastereomeric ratios decreased to 10 : 1 and 11 : 1, respectively. In addition, the functional group tolerance of the exocyclic-olefinic moiety in benzofulvenes exhibited semblable peculiarity as those of spiro[5,5]indenes described in Scheme 2, affording spiro[5,6]indenes **4l**—**4r** in 52% to 70% yields. However, an exocyclic carboethoxy instead of a phenyl group gave the desired product **4s** in 19% yield and 5 : 1 regioselectivity. Besides, the reactivity of different VDCP-diesters **2** was also tested in this [4+2] cycloaddition, affording the corresponding products **4t**—**4w** in 56% to 68% yields. Interestingly, the diastereoselectivity for **4u** (> 20 : 1) and **4v** (14 : 1) seems to be reversed as compared with **3ad** (> 20 : 1) and **3ae** (10 : 1) in the [3+2] cycloaddition shown in Scheme 2, perhaps due to the mutual steric effect between ligand and the substrates. VDCP diester with an ethyl unit at the terminal position of allene furnished the desired product **4w** in 4 : 1 regioselectivity along with high diastereoselectivity.

Scheme 2 Substrate scope for the spiro[5,5]indenes **3**




[a] Reaction condition: **1** (0.1 mmol), **2** (0.2 mmol), $Pd(PPh_3)_4$ (5 mol%), Ligand (7.5 mol%), $Yb(OTf)_3$ (10 mol%), Cs_2CO_3 (0.1 mmol), 4 Å MS (0.1 g), THF (1.0 mL), heated at 55 °C for 8.0 h. [b] Scale-up reaction: **1a** (229 mg, 1.0 mmol), **2a** (0.35 mL, 1.5 mmol), $Pd(PPh_3)_4$ (58 mg, 5 mol%), DPEphos (40 mg, 7.5 mol%), $Yb(OTf)_3$ (62 mg, 10 mol%), Cs_2CO_3 (325 mg, 1.0 mmol), 4 Å MS (1.0 g), THF (10.0 mL), heated at 55 °C for 8.0 h. [c] d.r. = 8:1. [d] d.r. = 12:1. [e] d.r. = 5:1. [f] d.r. = 10:1. [g] d.r. = 10:1.

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH


16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Scheme 3 Substrate scope for the spiro[5,6]indenes **4**



[a] Reaction condition: **1** (0.1 mmol), **2** (0.2 mmol), Pd_2dba_3 (2.5 mol%), (*R*)-Ph-PHOX (7.5 mol%), $Yb(OTf)_3$ (10 mol%), Cs_2CO_3 (0.1 mmol), 4 Å MS (0.1 g), THF (1.0 mL), heated at 65 °C for 8.0 h. [b] Scale-up reaction: **1a** (229 mg, 1.0 mmol), **2a** (0.35 mL, 1.5 mmol), Pd_2dba_3 (23 mg, 2.5 mol%), (*R*)-Ph-PHOX (30 mg, 7.5 mol%), $Yb(OTf)_3$ (62 mg, 10 mol%), Cs_2CO_3 (325 mg, 1.0 mmol), 4 Å MS (1.0 g), THF (10.0 mL), heated at 65 °C for 8.0 h. [c] d.r. = 10:1. [d] d.r. = 11:1. [e] d.r. = 14:1.

Scheme 4 [a] Large-scale synthesis. [b] Asymmetric investigation. [c] Further transformations of **3a** and **4a**.




On the basis of above investigations, the potential synthetic practicability of this method on a large-scale was next assessed. As shown in Scheme 4, a, the desired products **3a** and **4a** were acquired in 85% yield (0.374 g) and 59% yield (0.259 g) in the reactions of **1a** (0.229 g) with **2a** (0.355 mL) under respective optimal reaction conditions. Furthermore, the feasibility of asymmetric catalysis was also investigated on both cycloaddition modes (Scheme 4, b). Regrettably, the spiro[5,5]indene **3a** could be only obtained in 58% yield along with 34% enantiomeric excess using spiral ligand (*S*)-PMP-SiSDP as a chiral ligand (for more information on the asymmetric investigation on [3+2] cycloaddition, see Table S1 in the Supporting Information). On the other hand, the [4+2]

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Scheme 5 Control experiments



cycloaddition product **4a** was formed with 11% enantiomeric excess upon using the original (*R*)-Ph-PHOX as a chiral ligand.


Moreover, the further conversions of **3a** and **4a** were also conducted as shown in Scheme 4, c. Aminolysis of the carboethoxy groups upon treating **3a** with NaOMe in formamide at 75 °C produced **3a-I** in 67% yield, which was derived from a further Michael addition on the vinylcyanide moiety after the aminolysis reaction. Moreover, a difunctionalization process of terminal allene with SIMesCuCl and $(BPin)_2$ in the presence of KO^{t}Bu and EtI developed by Ito's group[13] was also used in the transformation of **3a**, generating **3a-II** in 87% yield in DMF at room temperature; **3a** probably went through an insertion of alkenyl-copper species to conjugated cyanide due to the inductive effect brought from the conjugated cyanide similar as that of aminolysis reaction. The structures of **3a-I** and **3a-II** have been unambiguously identified by X-ray diffraction and their ORTEP drawings are shown in Scheme 4. On the other hand, **4a** could be transformed to the corresponding epoxide **4a-I** in 74% yield along with 1.5 : 1 diastereomeric ratio through an epoxidation upon treating with *m*-CPBA and Na_2CO_3 in THF.

In order to clarify the mechanism of this reaction, a series of control experiments were performed to probe the reaction pathways (Scheme 5). Neither [3+2] nor [4+2] cycloaddition occurred in the absence of palladium resources, while removing $Yb(OTf)_3$ led to the lower yields for both two types of cycloadditions. Hence, palladium as the basic catalyst and $Yb(OTf)_3$ as the cocatalyst were confirmed. In addition, using $Pd(PPh_3)_4$ as the palladium source gave the [3+2] cycloaddition product **3a** in 28% yield in absence of DPEphos and none of the [4+2] cycloaddition product **4a** was formed in the absence of (*R*)-Ph-PHOX ligand, rendering that the bidentate ligand plays a key role in these cycloaddition reactions.

On the basis of the above experimental results and our previous investigations on the reaction pathways,[12,14] a plausible reaction mechanism was proposed in Scheme 6. The zwitterionic π-propargyl palladium species **I**, generated from an oxidative addition between ligand-coordinated Pd(0) and Yb^{3+}-coordinated VDCP-diester **2a**, launches a nucleophilic attack to the exocyclic position of benzofulvene **1a**, forming an intermediate **II** stabilized by the aromaticity of cyclopentadienyl anion. When palladium catalyst coordinates with DPEphos, a bidentate ligand with a large bite angle, the cyclopentadienyl anion tends to attack onto the carbon atom at *3*-position of the π-propargyl palladium species to provide [3+2] cycloaddition intermediate **III-3-pre**, and subsequently produces spiro[5,5]indene **3a** through a ligand exchange process with **2a** along with the regeneration of the zwitterionic π-propargyl palladium species **I** to complete the catalytic cycle. Whereas, the cyclopentadienyl anion is more likely to approach onto the carbon atom at *2*-position of π-propargyl palladium species if palladium catalyst is coordinated by (*R*)-Ph-PHOX, a special bidentate ligand with a small bite angle, affording palladacyclobutene species **III**,[15] which can be formed in a similar way as that of palladacyclobutane derived from π-allyl palladium species.[15] Then, the palladium intermediate **III** shall undergo a base-induced protonation to form a π-allyl palladium species **IV**, which undergoes a β-H elimination to furnish the [4+2] cycloaddition intermediate **V-4-pre**. A ligand exchange process between **V-4-pre** and **2a** affords spiro[5,6]indene **4a** and regenerates the zwitterionic π-propargyl palladium species **I** to complete the catalytic cycle.

As shown in Scheme 6, the selectivity originates from the cyclopentadienyl anion moiety of the intermediate **II** attacking different carbon of π-propargyl palladium moiety. Thus, density functional theory (DFT) calculation was used to investigate the selectivity-determining step (Figure 2). When DPEphos is utilized as the ligand, the cycloaddition can undergo two pathways originated from η^1/η^3 type intermediates **INT A-1** or **INT A-1'** to form [3+2] product or [4+2] product. Although the energy of intermediate **INT A-1'** is lower than that of **INT A-1** by 13.7 kcal/mol, the

Scheme 6 A plausible mechanism



 © 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




Figure 2 DFT calculation studies of ligand-controlled selectivity.

[4+2] pathway has to overcome a much higher energy barrier which was 19.0 kcal/mol rather than 12.9 kcal/mol as [3+2] pathway. On another side, the energy of [4+2] product **INT A-2'** was slightly higher than that of **INT A-1'** by 1.4 kcal/mol, which means that [4+2] pathway is a reversible process; in contrast, the [3+2] pathway is an exothermic process (ΔG = −13.8 kcal/mol). Therefore, the [3+2] product is thermodynamically and kinetically favorable which is in line with the experimental result. In the case of (*R*)-Ph-PHOX used as the ligand, [4+2] pathway and [3+2] pathway shared the same intermediate **INT B-1**. The [4+2] pathway is easier to occur both thermodynamically and kinetically, which showed an energy gap of 15.1 kcal/mol between two transition states and an energy gap of 4.1 kcal/mol between the two products. We speculate that DPEphos with a large bite angle, which is about 104°—108° upon coordinating with palladium source, would produce more steric hindrance, thereby restraining the formation of **TS A'** from **INT A-1'** and raising the energy barrier of [4+2] type reaction; (*R*)-Ph-PHOX with a small bite angle, that is about 81°—90° upon coordinating with palladium source, results in much smaller energy barrier for generation of the [4+2] product due to the steric effect.

Conclusions

In summary, a ligand-controlled divergent [3+2]/[4+2] cycloaddition between benzofulvenes and zwitterionic π-propargyl palladium species generated from VDCP-diesters has been established for the rapid construction of substituted spiro[5,5]indenes and spiro[5,6]indenes in moderate to excellent yields with good functional group toleration and a broad substrate scope. The plausible mechanistic paradigm was proposed based on the control experiments and our previous work, rendering that the reaction proceeds through a zwitterionic π-propargyl palladium species, a key intermediate in these cycloaddition reactions. The large-scale synthesis and the further conversion of the obtained cyclized products indicated that this synthetic method can easily access multi-substituted endocyclic compounds, polycyclic compounds or epoxide substances. The DFT calculations reveal the origin of product selectivity. The further applications of this methodology in the preparing bioactive substances or drug-like molecules are also undergoing in our laboratory.

Experimental

General procedure for the [3+2] products. Tetrakis(triphenylphosphine)palladium (0.005 mmol, 5 mol%) and DPEPhos (0.007 mmol, 7.5 mol%) were added to an oven dried 10 mL sealed tube equipped with a magnetic stirring bar under argon atmosphere. The resulting catalyst system was stirred in THF (degassed) (1.0 mL) with 4 Å MS (100 mg) for 30 min. After that, **1** (0.1 mmol), **2** (0.2 mmol), $Yb(OTf)_3$ (0.01 mmol, 10 mol%) and Cs_2CO_3 (0.1 mmol, 1.0 eq) were added under argon atmosphere. The reaction mixture was stirred at 55 °C for 8.0 h. The reaction solution was filtered through a pad of celite to remove insoluble solids. Then, the solvent was evaporated under vacuum and the residue was purified by a silica gel column chromatography to afford the desired product **3**.

General procedure for the [4+2] products. Tris(dibenzylideneacetone)dipalladium (0.002 mmol, 2.5 mol%) and (*R*)-Ph-PHOX (0.007 mmol, 7.5 mol%) were added to an oven dried 10 mL sealed tube equipped with a magnetic stirring bar under argon atmosphere. The resulting catalyst system was stirred in THF (degassed) (1.0 mL) with 4 Å MS (100 mg) for 30 min. After that, **1** (0.1 mmol), **2** (0.2 mmol), $Yb(OTf)_3$ (0.01 mmol, 10 mol%) and Cs_2CO_3 (0.1 mmol, 1.0 eq) were added under argon atmosphere. The reaction mixture was stirred at 65 °C for 8.0 h. The reaction solution was filtered through a pad of celite to remove insoluble solids. Then, the solvent was evaporated under vacuum and the residue was purified by a silica gel column chromatography to afford the desired product **4**.

Supporting Information

The supporting information for this article is available on the WWW under https://doi.org/10.1002/cjoc.202401301.

Acknowledgement

We are grateful for the financial support from the National Key R & D Program of China (2023YFA1506700), the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB0590000), and the National Natural Science Foundation of China (21372250, 21121062, 21302203, 21772037, 21772226, 21861132014, 91956115 and 22171078).

References

[1] (a) Garthwaite, S. M.; McMahon, E. G. The Evolution of Aldosterone

 © 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Antagonists. *Mol. Cell. Endocrinol.* **2004**, *217*, 27–31; (b) Cella, J. A.; Brown, E. A.; Burtner, R. R. Steroidal Aldosterone Blockers. I. *J. Org. Chem.* **1959**, *24*, 743–748; (c) Cella, J. A.; Tweit, R. C. Steroidal Aldosterone Blockers. II[1]. *J. Org. Chem.* **1959**, *24*, 1109–1110; (d) Quintavalla, A. Spirolactones: Recent Advances in Natural Products, Bioactive Compounds and Synthetic Strategies. *Curr. Med. Chem.* **2018**, *25*, 917–962; (e) Elger, W.; Beier, S.; Pollow, K.; Garfield, R.; Shi, S.-Q.; Hillisch, A. Conception and Pharmacodynamic Profile of Drospirenone. *Steroids* **2003**, *68*, 891–905.

[2] Smith, L. K.; Baxendale, I. R. Total syntheses of natural products containing spirocarbocycles. *Org. Biomol. Chem.* **2015**, *13*, 9907–9933.

[3] (a) Marson, C. M. New and unusual scaffolds in medicinal chemistry. *Chem. Soc. Rev*. **2011**, *40*, 5514–5533; (b) Zheng, Y.; Tice, C. M.; Singh, S. B. The use of spirocyclic scaffolds in drug discovery. *Bioorg. Med. Chem. Lett*. **2014**, *24*, 3673–3682; (c) Hiesinger, K.; Dar'in, D.; Proschak, E.; Krasavin, M. Spirocyclic Scaffolds in Medicinal Chemistry. *J. Med. Chem*. **2021**, *64*, 150–183.

[4] (a) Zhu, S.-F.; Zhou, Q.-L. Transition-Metal-Catalyzed Enantioselective Heteroatom–Hydrogen Bond Insertion Reactions. *Acc. Chem. Res.* **2012**, *45*, 1365–1377; (b) Zhu, S.-F.; Zhou, Q.-L. Iridium-Catalyzed Asymmetric Hydrogenation of Unsaturated Carboxylic Acids. *Acc. Chem. Res*. **2017**, *50*, 988–1001; (c) Yang, F.; Xie, J.-H.; Zhou, Q.-L. Highly Efficient Asymmetric Hydrogenation Catalyzed by Iridium Complexes with Tridentate Chiral Spiro Aminophosphine Ligands. *Acc. Chem. Res*. **2023**, *56*, 332–349; (d) Li, Y.-P.; Zhu, S.-F.; Zhou, Q.-L. Chiral Spiro Phosphoramide-Catalyzed Sulfa-Michael Addition/Enantioselective Protonation of Exocyclic Enones. *Org. Lett*. **2019**, *21*, 9391–9395; (e) Wang, B.; Zhang, T.-Z.; Xiao, L.-J.; Zhou, Q.-L. Nickel-Catalyzed Intermolecular Reductive Coupling of Styrene and Its Derivatives with Imines Using Methanol as a Reductant. *ACS Catal*. **2023**, *13*, 8692–8699; (f) Han, A.-C.; Xiao, L.-J.; Zhou, Q.-L. Construction of Ge-Stereogenic Center by Desymmetric Carbene Insertion of Dihydrogermanes. *J. Am. Chem. Soc*. **2024**, *146*, 5643–5649; (g) Xu, B.; Ji, D.-T.; Wu, L.-Z.; Zhou, L.-J.; Liu, Y.; Zhang, Z.-M.; Zhang, J.-L. Palladium/Xu-Phos-Catalyzed Enantioselective Cascade Heck/Remote C(sp^2)–H Alkylation Reaction. *Chem* **2022**, *8*, 836–849; (h) Chu, H.-K.; Cheng, J.; Yang, J.-F.; Guo, Y.-L.; Zhang, J.-L. Asymmetric Dearomatization of Indole via Palladium/PC-Phos-Catalyzed Dynamic Kinetic Transformation. *Angew. Chem. Int. Ed.* **2020**, *59*, 21991–21996.

[5] Zhao, Q.-H.; Tang, L.-C.; Jiao, P. Construction of Spiro Skeletons in 2,2',3,3'-Tetrahydro-1,1'-spirobi[1*H*-indene]-7,7'-diol (SPINOL) and Analogues. *Chin. J. Org. Chem*. **2021**, *41*, 3400–3413.

[6] (a) Zheng, J.; Wang, S.-B.; Zheng, C.; You, S.-L. Asymmetric Dearomatization of Naphthols via a Rh-Catalyzed C(sp^2)–H Functionalization/Annulation Reaction. *J. Am. Chem. Soc*. **2015**, *137*, 4880–4883; (b) Yang, L.; Zheng, H.-Y.; Luo, L.; Nan, J.; Liu, J.-J.; Wang, Y.-Y.; Luan, X.-J. Palladium-Catalyzed Dynamic Kinetic Asymmetric Transformation of Racemic Biaryls: Axial-to-Central Chirality Transfer. *J. Am. Chem. Soc*. **2015**, *137*, 4876–4879.

[7] (a) Chai, Z.; Rainey, T. J. Pd(II)/Brønsted Acid Catalyzed Enantioselective Allylic C–H Activation for the Synthesis of Spirocyclic Rings. *J. Am. Chem. Soc*. **2012**, *134*, 3615–3618; (b) Fan, L.-L.; Takizawa, S.; Takeuchi, Y.; Takenaka, K.; Sasai, H. Pd-catalyzed enantioselective intramolecular α-arylation of α-substituted cyclic ketones: facile synthesis of functionalized chiral spirobicycles. *Org. Biomol. Chem*. **2015**, *13*, 4837–4840.

[8] (a) Diekmann, M.; Bockstiegel, G.; Lützen, A.; Friedemann, M. Saak, W.; Haase, D.; Beckhaus, R. Chiral Bis(η^5:η^1-pentafulvene)titanium Complexes. *Organometallics* **2006**, *25*, 339–348; (b) Yan, J.-C.; Cheng, X.; Zhou, Q.-L.; Pei, J. Chiral Polyfluorene Derivatives: Synthesis, Chiroptical Properties, and Investigation of the Structure–Property Relationships. *Macromolecules* **2007**, *40*, 832–839; (c) Cappelli, A.; Galeazzi, S.; Zanardi, I.; Travagli, V.; Anzini, M.; Mendichi, R.; Petralito, S.; Memoli, A.; Paccagnini, E.; Peris, W.; Giordani, A.; Makovec, F.; Fresta, M.; Vomero, S. A nanocomposite material formed by benzofulvene polymer nanoparticles loaded with a potent 5-HT_3 receptor antagonist (CR3124). *J. Nanopart. Res.* **2010**, *12*, 895–903; (d) Kosaka, Y.; Kitazawa, K.; Inomata, S.; Ishizone, T. Living Anionic Polymerization of Benzofulvene: Highly Reactive Fixed Transoid 1,3-Diene. *ACS Macro Lett.* **2013**, *2*, 164–167; (e) Kosaka, Y.; Kawauchi, S.; Goseki, R.; Ishizone, T. High Anionic Polymerizability of Benzofulvene: New Exo-Methylene Hydrocarbon Monomer. *Macromolecules* **2015**, *48*, 4421–4430.

[9] Donslund, B. S.; Nielsen, R. P.; Mønsted, S. M. N.; Jørgensen, K. A. Benzofulvenes in Trienamine Catalysis: Stereoselective Spiroindene Synthesis. *Angew. Chem. Int. Ed*. **2016**, *55*, 11124–11128.

[10] Donslund, B. S.; Jessen, N. I.; Jakobsen, J. B.; Monleo'n, A.; Nielsen, R. P.; Jørgensen, K. A. Enantioselective formation of cyclopropane spiroindenes from benzofulvenes by phase transfer catalysis. *Chem. Commun.* **2016**, *52*, 12474–12477.

[11] Chang, X.; Liu, X.-T.; Li, F.-F.; Yang, Y.-H.; Chung, L. W.; Wang, C-J. Electron-rich benzofulvenes as effective dipolarophiles in copper(I)-catalyzed asymmetric 1,3-dipolar cycloaddition of azomethine ylides. *Chem. Sci.* **2023**, *14*, 5460–5469.

[12] (a) Niu, B.; Wei, Y.; Shi, M. Palladium catalyzed divergent cycloadditions of vinylidenecyclopropane-diesters with methyleneindolinones enabled by zwitterionic π-propargyl palladium species. *Chem. Commun.* **2021**, *57*, 4783–4786; (b) Shen, J.-H.; Long, Y.-J.; Shi, M.; Wei, Y. Palladium-catalyzed and ligand-controlled divergent cycloadditions of vinylidenecyclopropane-diesters with *para*-quinone methides enabled by zwitterionic π-propargyl palladium species. *Org. Chem. Front.* **2024**, *11*, 3214–3224; (c) Long, Y.-J.; Shen, J.-H.; Wei, Y.; Shi, M. Substrate-Controlled [8+3] Cycloaddition of Tropsulfimides and Tropones with Zwitterionic Allenyl Palladium Species Derived from Vinylidenecyclopropane-diesters. *J. Org. Chem.* **2024**, *89*, 14831–14850; (d) Long, Y.-J.; Shen, J.-H.; Shi, M.; Wei, Y. Theoretical Studies on the Reaction Mechanism for the Cycloaddition of Zwitterionic π-Allenyl Palladium Species: Substrate-Controlled Isomerization. *Molecules* **2025**, *30*, 103–113.

[13] Ozawa, Y.; Endo, K.; Ito, H. Regio- and Stereoselective Synthesis of Multi-Alkylated Allylic Boronates through Three-Component Coupling Reactions between Allenes, Alkyl Halides, and a Diboron Reagent. *J. Am. Chem. Soc*. **2021**, *143*, 13865–13877.

[14] Zhong, X.-H.; Long, Y.-J.; Wei, Y.; Shi, M. Palladium-Catalyzed $Sc(OTf)_3/Cs_2CO_3$-Assisted C–O Cross-Coupling of Vinylidenecyclopropane-Diesters with Phenolic Compounds. *Adv. Syn. Catal.* **2024**, *366*, 3387–3396.

[15] (a) Montgomery, T. D.; Rawal, V. H. Palladium-Catalyzed Modular Synthesis of Substituted Piperazines and Related Nitrogen Heterocycles. *Org. Lett*. **2016**, *18*, 740–743; (b) O'Broin, C. Q.; Guiry, P. J. Synthesis of 2‑Amino-1,3-dienes from Propargyl Carbonates via Palladium-Catalyzed Carbon–Nitrogen Bond Formation. *Org. Lett*. **2020**, *22*, 879–883; (c) Mondal, D.; Pal, G.; Chowdhury, C. Palladium(0)-catalysed regioselective cyclisations of 2-amino(tosyl) benzamides/sulphonamides: the stereoselective synthesis of 3-ylidene-[1,4]benzodiazepin-5-ones/benzo[*f*][1,2,5]thiadiazepine-1,1-dioxides. *Chem. Commun.* **2021**, *57*, 5462–5465; (d) Dai, M.-F.; Sun, Z.-M.; Chen, L.-A. Palladium-Catalyzed Regiodivergent Synthesis of 1,3-Dienyl and Allyl Esters from Propargyl Esters. *Angew. Chem. Int. Ed*. **2022**, *61*, e202203835.


Manuscript received: December 22, 2024
Manuscript revised: February 10, 2025
Manuscript accepted: February 11, 2025
Version of record online: March 5, 2025

 © 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

16147065, 2025, 10, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cjoc.202401301 by Zhejiang University, Wiley Online Library on [01/02/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

The Authors



Left to Right: Yongjie Long, Xianhua Zhong, Min Shi, Yin Wei

© 2025 SIOC, CAS, Shanghai, & WILEY-VCH GmbH