ORGANIC CHEMISTRY FRONTIERS

RESEARCH ARTICLE

View Article Online
View Journal | View Issue

Check for updates

Cite this: *Org. Chem. Front.*, 2025, **12**, 3960

Received 27th December 2024,
Accepted 21st March 2025

DOI: 10.1039/d4qo02426e

rsc.li/frontiers-organic

Published on 22 March 2025. Downloaded by Zhejiang University on 1/30/2026 3:53:45 AM.

Bromotrifluoromethoxylation of allenes: expedient access to allylic trifluoromethoxy derivatives†

Rui Liu,[a] Yangdong Hou *[b] and Pingping Tang*[a,b]

A mild and efficient method for the bromotrifluoromethoxylation of allenes has been explored using trifluoromethyl arylsulfonate (TFMS) as the trifluoromethoxy source. A series of allylic trifluoromethoxy derivatives were obtained with high yields and good regioselectivity by terminal selectivity of the allenes. This transition metal-free process expands the scope of the application of trifluoromethoxy reagents.


Introduction

The introduction of fluorinated functional groups into organic molecules has a significant impact on their physical and chemical properties.[1] Thus, more and more fluorinated drug molecules are being synthesized.[2] In particular, the trifluoromethoxy group (OCF_3) brings good metabolic stability and high lipophilicity to molecules, earning it the nickname of superhalogen.[3] The unique physical and chemical properties of trifluoromethoxy functional groups have attracted attention in the fields of materials science and pharmaceuticals. However, the number of marketed drugs and agricultural chemicals containing OCF_3 remains low, and the study of OCF_3 is largely behind the development of other fluorinated functional groups. Pioneering works have focused on the construction of the O–CF_3 bond from an already installed OH group, but this indirect method usually requires harsh conditions or toxic reagents (*e.g.*, HF and SF_4).[4] Directly introducing an OCF_3 functional group is an effective alternative method for obtaining trifluoromethoxylated compounds.[5] However, this is limited by the reversible decomposition of trifluoromethoxy and trifluoromethoxylation reagents.

In 2017, our research group developed trifluoromethyl benzenesulfonate (TFMS) as a trifluoromethoxy reagent, which can be synthesized in one step from Togni reagent. It has been used to achieve the first asymmetric trifluoromethoxylation reaction, realizing internal trifluoromethoxylation selectivity of alkenes (Scheme 1a).[6] Allenes exist in many natural products and pharmaceuticals.[7] The high activity of allenes makes them widely used as synthons for cyclization,[8] hydrogenheteroatom reactions[9] and multifunctionalization.[10] With our continuous efforts on the trifluoromethoxylation reactions,[11] we speculated that if allenes were subjected to our reaction system, OCF_3 would simultaneously be easily introduced into the products. However, the reaction of allenes presented more complex reactivity than that of alkenes. The versatile reactivities of allenes allow for the selective modification of different positions, enabling the synthesis of different regioselectivity functionalized compounds,[12] which could deliver more possible trifluoromethoxylation products. This interesting but questionable selectivity motivated us to investigate this chemistry. Conversely, due to the toxic reagents and the cost of transition metal catalysts or metallic reagents, transition metal-free processes with readily available OCF_3 sources have always attracted the attention of chemists. Herein, we report our efforts to develop a transition metal-free process for bromotrifluoromethoxylation of allenes with our trifluoromethoxy reagent (Scheme 1b).

Results and discussion

To validate our hypothesis, the initial study focused on the reaction of substrate **1a**, which was treated with the previous



Scheme 1 Intermolecular bromotrifluoromethoxylations.

[a]*College of Chemistry and Environmental Science, Kashi University, Xinjiang 844008, China. E-mail: ptang@nankai.edu.cn, liurui1237890@163.com*
[b]*State Key Laboratory and Institute of Elemento-Organic Chemistry, College of Chemistry Nankai University, Tianjin 300071, China. E-mail: hyangdong2015@163.com*

†Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d4qo02426e


This journal is © the Partner Organisations 2025

Published on 22 March 2025. Downloaded by Zhejiang University on 1/30/2026 3:53:45 AM.

olefin reaction conditions. We were delighted to find the bromosubstituted allyl trifluoromethoxy product **3a** in 24% yield with excellent regioselectivity but poor stereoselectivity (see Table S1 in the ESI†). We did not detect the benzyl trifluoromethoxylation product, similar to the olefin reactions in our previous work. Meanwhile, **3a** was obtained in 11% yield without the addition of silver fluoride and the ligand (Table 1, entry 1). NBS delivered the product in 16% yield and resulted in 56 : 44 *E*/*Z* selectivity. Other bromination reagents, such as TBCA, PHBP, TBCO and *N*-bromosaccharin afforded the desired product in less than 5% yield (Table 1, entries 3–6). Further optimization of the bromination reagents revealed that TBATB provided the best results, giving the corresponding product **3a** in 86% yield with 84 : 16 *E*/*Z* selectivity (Table 1, entry 7). The product **3a** was obtained with 76/24 *E*/*Z* selectivity when using 2 equivalents of TBATB (Table 1, entry 8). **3a** cannot be obtained when using NaF or CaF_2, perhaps due to the poor solubility of these two fluoride salts (Table 1, entries 9 and 11). Otherwise, KF can afford **3a** in 63% yield (Table 1, entry 10). However, the yield significantly decreased to 27% using silver fluoride (Table 1, entry 12). When the reaction was performed at 40 °C, the conversion after 16 h was slightly reduced (Table 1, entry 13). Meanwhile, upon heating the reaction mixture to 60 °C, compound **3a** was isolated in 78% yield (Table 1, entry 14).


With the optimized conditions in hand (entry 7 in Table 1), we investigated the scope of the present transformation of allenes (Scheme 2). Generally, electron-donating groups

Table 1 Optimization of the reaction conditions[a]



Entry	[Br$^+$]	[F$^-$]	*T*/°C	Yield/%	*E*/*Z*
1	DBDMH	CsF	50	11	63/37
2	NBS	CsF	50	16	56/44
3	TBCA	CsF	50	1	ND
4	PHBP	CsF	50	0	—
5	TBCO	CsF	50	0	—
6	*N*-Bromosaccharin	CsF	50	2	50/50
7	TBATB	CsF	50	86	84/16
8[b]	TBATB	CsF	50	80	76/24
9	TBATB	NaF	50	0	—
10	TBATB	KF	50	63	78/22
11	TBATB	CaF_2	50	0	—
12	TBATB	AgF	50	27	52/48
13	TBATB	CsF	40	74	80/20
14	TBATB	CsF	60	78	77/23

[a] Reaction conditions: **1a** (0.1 mmol), TFMS (3 equiv.), [Br$^+$] (1 equiv.), [F$^-$] (3.0 equiv.), CH_3CN (0.25 mL), 16 h. Yields were determined by ^{19}FNMR spectroscopy with benzotrifluoride as a standard. [b] TBATB (2 equiv.). TBATB = tetrabutylammonium tribromide. DBDMH = 1,3-dibromo-5,5-dimethylhydantoin. NBS = *N*-bromosuccinimide. TBCA = 1,3,5-tribroMo-1,3,5-triazinane-2,4,6-trione. PHBP = pyridinium bromide perbromide. TBCO = 2,4,4,6-tetrabromo-2,5-cyclohexadien-1-one.



Scheme 2 Substrate scope of trifluoromethoxylation. Reaction conditions: **1** (0.2 mmol), TFMS (3.0 equiv.), TBATB (1.0 equiv.), CsF (3.0 equiv.), CH_3CN (0.5 mL), 50 °C, 16 h. Yields of isolated products are given, unless otherwise noted.


including phenyl, Me, ^{t}Bu, and OMe were well tolerated, providing the corresponding products in 62–86% yields (**3a–3f**). Naphthylallene was also reactive, giving a higher yield (**3g**, 82%, *E*/*Z* = 83 : 17). Substrates that contain a halogen can be applied to this system, which allowed further decoration (**3h** and **3i**, 80–82%, *E*/*Z* = 90 : 10 and 89 : 11). Allene substrates with an ester group were transformed into products **3j–3l** with good *E*-isomer selectivity, probably due to the electron-with-

Published on 22 March 2025. Downloaded by Zhejiang University on 1/30/2026 3:53:45 AM.


drawing induction effect. Moreover, alkyne can survive in these conditions (**3m**, 62%, *E*/*Z* = 89 : 11). The reaction of sulfur-containing heterocyclic allene **1n** delivered the product with a lower stereoselectivity of the C=C configuration (**3n**, 65%, *E*/*Z* = 75 : 25). In contrast, substrate **1o** with an oxygen-containing heterocyclic structure afforded the product in a better stereoselectivity (**3o**, 58%, *E*/*Z* = 86 : 14). Heteroaromatic allenes, such as thiopheneallene, were also suitable for this reaction (**3p**, 56%, *E*/*Z* = 70 : 30). Notably, a substrate containing a ternary ring was also compatible with this catalytic system (**3q**, 65%, *E*/*Z* = 86 : 14). The desired products **3r** and **3s** were obtained in moderate yields, which highlighted the good potential post-modification of this methodology for natural complex products. The product **3t** was observed using alkylallene as a substrate; moreover, the small steric hindrance led to a decrease in stereoselectivity. Internal allene **1u** and disubstituted allene **1v** afforded a small amount of the products, possibly due to the low activity of the substrates. Similarly, no product was obtained for the diarylallene, and the alkylallene gave the product in 20% yield (see Scheme S1 in the ESI†).

To evaluate the practicability and efficiency of this reaction, a gram-scale synthesis of **3a** was conducted under the standard reaction conditions in 73% yield (Scheme 3a). Further transformation of the bromoallylic trifluoromethoxy products for the construction of a series of OCF_3-containing compounds was investigated (Scheme 3b). Under hydrogenation conditions, alkyl trifluoromethoxy **5a** can be obtained with a yield of 61%. Propargyl trifluoromethoxy compound **5b** was produced with a yield of 46% by eliminating hydrogen bromide.

To gain more insight into the reaction mechanism, we carried out some control experiments (see Schemes S2 and S3 in the ESI†). Monitoring of the reaction between **1a** and TBATB by ^{1}H NMR spectrum showed that the characteristic hydrogen of the allene has undergone a shift. The HRMS-ESI (*m*/*z*) results indicated that alkene bromide ions were generated in the reaction (Scheme 4). This indicates the appearance of bromonium active intermediates. Meanwhile, we also conducted D-NOESY NMR spectra of **3a** and **3d** to confirm the configuration (see Schemes S4–S6 in the ESI†).[13]



Scheme 3 Gram-Scale synthesis and synthetic transformations.



Scheme 4 Mechanistic study.

Conclusions

We have presented the first example of bromotrifluoromethoxylation of allenes using TFMS as an OCF_3 source. By the combination of TBATB with CsF, a transition metal-free process with a readily available OCF_3 source provided a range of allylic trifluoromethoxy derivatives in up to 86% yield and with complete regiocontrol, which are difficult to access by other methods. Due to the easy decomposition of the OCF_3 group, carefully selecting the appropriate bromine source, reaction temperature, solvent, and base were very critical, as all the factors jointly guaranteed the smooth transformation. This discovery also expanded the scope of the application of the trifluoromethoxy reagent.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by NSFC (21925105, 92156001), and the Program for Innovative Research Team in Kashi University.

References

1 (*a*) K. Müller, C. Faeh and F. Diederich, Fluorine in pharmaceuticals: looking beyond intuition, *Science*, 2007, **317**, 1881–1886; (*b*) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, Fluorine in medicinal chemistry, *Chem. Soc. Rev.*, 2008, **37**, 320–330; (*c*) O. A. Tomashenko and V. V. Grushin, Aromatic Trifluoromethylation with Metal Complexes, *Chem. Rev.*, 2011, **111**, 4475–4521; (*d*) L. E. Zimmer, C. Sparr and R. Gilmour, Fluorine Conformational Effects in Organocatalysis: An Emerging Strategy for Molecular Design, *Angew. Chem., Int. Ed.*, 2011, **50**, 11860–11871; (*e*) T. Liang, C. N. Neumann and T. Ritter, Introduction of Fluorine and Fluorine-Containing Functional Groups, *Angew. Chem., Int. Ed.*, 2013, **52**, 8214–

 This journal is © the Partner Organisations 2025

Published on 22 March 2025. Downloaded by Zhejiang University on 1/30/2026 3:53:45 AM.

8264; (*f*) P. A. Champagne, J. Desroches, J.-D. Hamel, M. Vandamme and J.-F. Paquin, Monofluorination of Organic Compounds: 10 Years of Innovation, *Chem. Rev.*, 2015, **115**, 9073–9174; (*g*) Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Aceña, V. A. Soloshonok, K. Izawa and H. Liu, Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II–III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas, *Chem. Rev.*, 2016, **116**, 422–518; (*h*) B. M. Johnson, Y.-Z. Shu, X. L. Zhuo and N. A. Meanwell, Metabolic and Pharmaceutical Aspects of Fluorinated Compounds, *J. Med. Chem.*, 2020, **63**, 6315–6386.

2 (*a*) N. G. Myers, J. K. Barbay and B. Y. Zhong, Asymmetric Synthesis of Chiral Organofluorine Compounds: Use of Nonracemic Fluoroiodoacetic Acid as a Practical Electrophile and Its Application to the Synthesis of Monofluoro Hydroxyethylene Dipeptide Isosteres within a Novel Series of HIV Protease Inhibitors, *J. Am. Chem. Soc.*, 2001, **123**, 7207–7219; (*b*) G. Hughes, P. N. Devine, J. R. Naber, P. D. O'Shea, B. S. Foster, D. J. McKay and R. P. Volante, Diastereoselective Reductive Amination of Aryl Trifluoromethyl Ketones and α-Amino Esters, *Angew. Chem., Int. Ed.*, 2007, **46**, 1839–1842; (*c*) K. Müller, C. Faeh and F. Diederich, Fluorine in Pharmaceuticals: Looking Beyond Intuition, *Science*, 2007, **317**, 1881–1886; (*d*) A. Cavaliere, K. C. Probst, A. D. Westwell and M. Slusarczyk, Fluorinated Nucleosides as an Important Class of Anticancer and Antiviral Agents, *Future Med. Chem.*, 2017, **9**, 1809–1833.

3 (*a*) M. Shimizu and T. Hiyama, Modern Synthetic Methods for Fluorine-Substituted Target Molecules, *Angew. Chem., Int. Ed.*, 2005, **44**, 214–231; (*b*) F. Leroux, P. Jeschke and M. Schlosser, *α*-Fluorinated Ethers, Thioethers, and Amines: Anomerically Biased Species, *Chem. Rev.*, 2005, **105**, 827–856; (*c*) P. Jeschke, E. Baston and F. R. Leroux, *α*-Fluorinated Ethers as "Exotic" Entity in Medicinal Chemistry, *Mini-Rev. Med. Chem.*, 2007, **7**, 1027–1034; (*d*) B. Manteau, S. Pazenok, J. P. Vors and F. R. Leroux, New trends in the chemistry of *α*-fluorinated ethers, thioethers, amines and phosphines, *J. Fluorine Chem.*, 2010, **131**, 140–158; (*e*) G. Landelle, A. Panossian and F. R. Leroux, *Curr. Top. Med. Chem.*, 2014, **14**, 941–951.

4 (*a*) L. M. Yagupolskii, Synthesis of derivatives of phenyl trifluoromethyl ether, *Dokl. Nauk SSSR*, 1955, **105**, 100–102; (*b*) N. N. Yarovenko and A. S. Vasileva, A new method for the introduction of trihalomethyl groups into organic molecules, *Zh. Obshch. Khim.*, 1958, **28**, 2502–2504; (*c*) W. A. Sheppard, α-Fluorinated Ethers. I. Aryl Fluoroalkyl Ethers, *J. Org. Chem.*, 1964, **29**, 1–11; (*d*) M. Kuroboshi, K. Suzuki and T. Hiyama, Oxidative desulfurization-fluorination of xanthates. A convenient synthesis of trifluoromethyl ethers and difluoro(methylthio)methyl ethers, *Tetrahedron Lett.*, 1992, **33**, 4173–4176; (*e*) K. Kanie, Y. Tanaka and T. Hiyama, A convenient synthesis of trifluoromethyl ethers by oxidative desulfurization-fluorination of dithiocarbonates, *Bull. Chem. Soc. Jpn.*, 2000, **73**, 471–484; (*f*) T. Umemoto, K. Adachi and S. Ishihara, CF_3 Oxonium Salts, O-(Trifluoromethyl) dibenzofuranium Salts: In Situ Synthesis, Properties, and Application as a Real CF_3^+ Species Reagent, *J. Org. Chem.*, 2007, **72**, 6905–6917; (*g*) R. Koller, K. Stanek, D. Stolz, R. Aardoom, K. Niedermann and A. Togni, Zinc-Mediated Formation of Trifluoromethyl Ethers from Alcohols and Hypervalent Iodine Trifluoromethylation Reagents, *Angew. Chem., Int. Ed.*, 2009, **48**, 4332–4336; (*h*) J. B. Liu, C. Chen, L. L. Chu, Z. H. Chen, X. H. Xu and F. L. Qing, Silver-Mediated Oxidative Trifluoromethylation of Phenols: Direct Synthesis of Aryl Trifluoromethyl Ethers, *Angew. Chem., Int. Ed.*, 2015, **54**, 11839–11842; (*i*) A. Tlili, F. Toulgoat and T. Billard, Synthetic Approaches to Trifluoromethoxy-Substituted Compounds, *Angew. Chem., Int. Ed.*, 2016, **55**, 11726–11735.

5 (*a*) A. A. Kolomeitsev, M. Vorobyev and H. Gillandt, Versatile application of trifluoromethyl triflate, *Tetrahedron Lett.*, 2008, **49**, 449–454; (*b*) O. Marrec, T. Billard, J. P. Vors, S. Pazenok and B. R. Langlois, A deeper insight into direct trifluoromethoxylation with trifluoromethyl triflate, *J. Fluorine Chem.*, 2010, **131**, 200–207; (*c*) O. Marrec, T. Billard, J. P. Vors, S. Pazenok and B. R. Langlois, A New and Direct Trifluoromethoxylation of Aliphatic Substrates with 2,4-Dinitro(trifluoromethoxy)benzene, *Adv. Synth. Catal.*, 2010, **352**, 2831–2837; (*d*) C. H. Huang, T. Liang, S. Harada, E. Lee and T. Ritter, Silver-Mediated Trifluoromethoxylation of Aryl Stannanes and Arylboronic Acids, *J. Am. Chem. Soc.*, 2011, **133**, 13308–13310; (*e*) K. N. Hojczyk, P. J. Feng, C. B. Zhan and M. Y. Ngai, Trifluoromethoxylation of Arenes: Synthesis of ortho-Trifluoromethoxylated Aniline Derivatives by OCF_3 Migration, *Angew. Chem., Int. Ed.*, 2014, **53**, 14559–14563; (*f*) C. H. Chen, P. H. Chen and G. S. Liu, Palladium-Catalyzed Intramolecular Aminotrifluoromethoxylation of Alkenes, *J. Am. Chem. Soc.*, 2015, **137**, 15648–15651; (*g*) W. J. Zheng, C. A. Morales-Rivera, J. W. Lee, P. Liu and M. Y. Ngai, Catalytic C-H Trifluoromethoxylation of Arenes and Heteroarenes, *Angew. Chem., Int. Ed.*, 2018, **57**, 9645–9649; (*h*) W. J. Zheng, J. W. Lee, C. A. Morales-Rivera, P. Liu and M. Y. Ngai, Redox-Active Reagents for Photocatalytic Generation of the OCF_3 Radical and (Hetero)Aryl C-H Trifluoromethoxylation, *Angew. Chem., Int. Ed.*, 2018, **57**, 13795–13799; (*i*) B. J. Jelier, P. F. Tripet, E. Pietrasiak, I. Franzoni, G. Jeschke and A. Togni, Radical Trifluoromethoxylation of Arenes Triggered by a Visible-Light-Mediated N-O Bond Redox Fragmentation, *Angew. Chem., Int. Ed.*, 2018, **57**, 13784–13789; (*j*) M. Zhou, C. F. Ni, Y. W. Zeng and J. B. Hu, Trifluoromethyl Benzoate: A Versatile Trifluoromethoxylation Reagent, *J. Am. Chem. Soc.*, 2018, **140**, 6801–6805; (*k*) M. Jakubczyk, S. Mkrtchyan, M. Shkoor, S. Lanka, Š. Budzák, M. Iliaš, M. Skoršepa and V. O. Iaroshenko, Mechanochemical Conversion of Aromatic Amines to Aryl Trifluoromethyl Ethers, *J. Am. Chem. Soc.*, 2022, **144**, 10438–10445.

Published on 22 March 2025. Downloaded by Zhejiang University on 1/30/2026 3:53:45 AM.

6 S. Guo, F. Cong, R. Guo, L. Wang and P. P. Tang, Asymmetric silver-catalysed intermolecular bromotrifluoromethoxylation of alkenes with a new trifluoromethoxylation reagent, *Nat. Chem.*, 2017, **9**, 546–551.

7 A. Hoffmann-Rçder and N. Krause, Synthesis and Properties of Allenic Natural Products and Pharmaceuticals, *Angew. Chem., Int. Ed.*, 2004, **43**, 1196–1216.

8 (*a*) Y. Kawaguchi, K. Yabushita and C. Mukai, Rhodium(I)-Catalyzed Cycloisomerization of Homopropargylallene-Alkynes through C(sp^3)-C(sp) Bond Activation, *Angew. Chem., Int. Ed.*, 2018, **57**, 4707–4711; (*b*) C. Zhu, J. Liu, B. K. Mai, F. Himo and J. E. Bäckvall, Efficient Stereoselective Carbocyclization to cis-1,4-Disubstituted Heterocycles Enabled by Dual Pd/Electron Transfer Mediator (ETM) Catalysis, *J. Am. Chem. Soc.*, 2020, **142**, 5751–5759; (*c*) M.-B. Li, D. Posevins, A. Geoffroy, C. Zhu and J.-E. Bäckvall, Efficient Heterogeneous Palladium-Catalyzed Oxidative Cascade Reactions of Enallenols to Furan and Oxaborole Derivatives, *Angew. Chem., Int. Ed.*, 2020, **59**, 1992–1996.

9 (*a*) R. Zimmer, C. U. Dinesh, E. Nandanan and F. A. Khan, Palladium-Catalyzed Reactions of Allenes, *Chem. Rev.*, 2000, **100**, 3067–3126; (*b*) P. Koschker, P. A. Lumbroso and B. Breit, Enantioselective Synthesis of Branched Allylic Esters via Rhodium-Catalyzed Coupling of Allenes with Carboxylic Acids, *J. Am. Chem. Soc.*, 2011, **133**, 20746–20749; (*c*) K. L. Butler, M. Tragni and R. A. Widenhoefer, Gold(I)-Catalyzed Stereoconvergent, Intermolecular Enantioselective Hydroamination of Allenes, *Angew. Chem., Int. Ed.*, 2012, **51**, 5175–5178; (*d*) A. B. Pritzius and B. Breit, Asymmetric Rhodium-Catalyzed Addition of Thiols to Allenes: Synthesis of Branched Allylic Thioethers and Sulfones, *Angew. Chem., Int. Ed.*, 2015, **54**, 3121–3125; (*e*) R. Blieck, J. Bahri, M. Taillefer and F. Monnier, Copper-Catalyzed Hydroamination of Terminal Allenes, *Org. Lett.*, 2016, **18**, 1482–1485; (*f*) Z. P. Yang and J. Wang, Enantioselective Palladium-Catalyzed Hydrophosphinylation of Allenes with Phosphine Oxides: Access to Chiral Allylic Phosphine Oxides, *Angew. Chem., Int. Ed.*, 2021, **60**, 27288–27292.

10 (*a*) F. Meng, H. Jang, B. Jung and A. H. Hoveyda, Cu-Catalyzed Chemoselective Preparation of 2-(Pinacolato) boron-Substituted Allylcopper Complexes and their In Situ Site-, Diastereo-, and Enantioselective Additions to Aldehydes and Ketones, *Angew. Chem., Int. Ed.*, 2013, **52**, 5046–5051; (*b*) Y. Tani, T. Fujihara, J. Terao and Y. Tsuji, Copper-Catalyzed Regiodivergent Silacarboxylation of Allenes with Carbon Dioxide and a Silylborane, *J. Am. Chem. Soc.*, 2014, **136**, 17706–17709; (*c*) W. X. Zhao and J. Montgomery, Cascade Copper-Catalyzed 1,2,3-Trifunctionalization of Terminal Allenes, *J. Am. Chem. Soc.*, 2016, **138**, 9763–9766.

11 (*a*) X. H. Jiang, Z. J. Deng and P. P. Tang, Direct Dehydroxytrifluoromethoxylation of Alcohols, *Angew. Chem., Int. Ed.*, 2018, **57**, 292–295; (*b*) H. D. Yang, F. Wang, X. H. Jiang, Y. Zhou, X. F. Xu and P. P. Tang, Silver-Promoted Oxidative Benzylic C-H Trifluoromethoxylation, *Angew. Chem., Int. Ed.*, 2018, **57**, 13266–13277; (*c*) S. Q. Yang, M. Chen and P. P. Tang, Visible-Light Photoredox-Catalyzed and Copper-Promoted Trifluoromethoxylation of Arenediazonium Tetrafluoroborates, *Angew. Chem., Int. Ed.*, 2019, **58**, 7840–7844; (*d*) Y. Li, Y. Yang, J. R. Xin and P. P. Tang, Nucleophilic trifluoromethoxylation of alkyl halides without silver, *Nat. Commun.*, 2020, **11**, 755–761; (*e*) Z. J. Deng, M. X. Zhao, F. Wang and P. P. Tang, Selective C-H trifluoromethoxylation of (hetero)arenes as limiting reagent, *Nat. Commun.*, 2020, **11**, 2569–2577; (*f*) J. R. Xin, X. Y. Deng and P. P. Tang, Silver-Catalyzed Trifluoromethoxylation of Aziridines, *Org. Lett.*, 2022, **24**, 881–885; (*g*) Y. D. Hou, Z. Zhang, X. Y. Sun, Z. Yang, Y. X. Luan and P. P. Tang, Copper-Catalyzed Enantioselective Trifluoromethoxylation of Propargyl Sulfonates, *Angew. Chem., Int. Ed.*, 2023, **62**, e202218919; (*h*) Z. J. Deng, L. D. Meng, X. Bing, S. X. Niu, X. F. Zhang, J. Q. Peng, Y. X. Luan, L. Chen and P. P. Tang, Silver-Enabled Dearomative Trifluoromethoxylation of Indoles, *J. Am. Chem. Soc.*, 2024, **146**, 2325–2332.

12 (*a*) S. Ma, Some Typical Advances in the Synthetic Applications of Allenes, *Chem. Rev.*, 2005, **105**, 2829–2872; (*b*) S. Yu and S. Ma, Allenes in Catalytic Asymmetric Synthesis and Natural Product Syntheses, *Angew. Chem., Int. Ed.*, 2012, **51**, 3074–3112.

13 (*a*) D. Biswas, D. R. Kishore and J. Naveen, ^{t}BuOK Assisted Regio- and Stereo-Selective Alkoxy-Isomerization of Ortho-Alkynyl Allylic Alcohols: Synthesis of Alkylidene Dihydroisobenzofurans and Indenones, *ChemistrySelect*, 2024, **9**, e202400224; (*b*) F. Kayser, M. Biesemans, H. Pan, M. Gielen and R. Willem, Structural and conformational characterization of (*Z*,)-17-[2-(triphenylstannyl) vinyl]-4-estren-17β-ol using 2D proton detected ^{1}H-^{13}C relay and 2D NOESY spectroscopy, *Magn. Reson. Chem.*, 1992, **30**, 877–882.

 This journal is © the Partner Organisations 2025