



Aalto University
School of Electrical
Engineering

Stochastic (partial) differential equations and Gaussian processes

Simo Särkkä

Aalto University, Finland

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Kernel vs. SPDE representations of GPs

GP model $\mathbf{x} \in \mathbb{R}^d, t \in \mathbb{R}$	Equivalent S(P)DE model
Spatial $k(\mathbf{x}, \mathbf{x}')$	SPDE model (\mathcal{L} is an operator) $\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$
Temporal $k(t, t')$	State-space/SDE model $\frac{d\mathbf{f}(t)}{dt} = \mathbf{A} \mathbf{f}(t) + \mathbf{L} w(t)$
Spatio-temporal $k(\mathbf{x}, t; \mathbf{x}', t')$	Stochastic evolution equation $\frac{\partial}{\partial t} \mathbf{f}(\mathbf{x}, t) = \mathcal{A}_x \mathbf{f}(\mathbf{x}, t) + \mathbf{L} w(\mathbf{x}, t)$

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Why use S(P)DE solvers for GPs?

- The $O(n^3)$ computational complexity is a challenge.
- What do we get:
 - $O(n)$ state-space methods for SDEs/SPDEs.
 - Sparse approximations developed for SPDEs.
 - Reduced rank Fourier/basis function approximations.
 - Path to non-Gaussian processes.
- Downsides:
 - We often need to approximate.
 - Mathematics can become messy.

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Ornstein-Uhlenbeck process

- The mean and covariance functions:

$$m(x) = 0$$

$$k(x, x') = \sigma^2 \exp(-\lambda|x - x'|)$$

- This has a *path representation* as a stochastic differential equation (SDE):

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t).$$

where $w(t)$ is a white noise process with x relabeled as t .

- Ornstein–Uhlenbeck process is a Markov process.
- What does this actually mean \implies white board.

Ornstein-Uhlenbeck process

- The mean and covariance functions:

$$m(x) = 0$$

$$k(x, x') = \sigma^2 \exp(-\lambda |x - x'|)$$

- This has a *path representation* as a stochastic differential equation (SDE):

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t).$$

where $w(t)$ is a white noise process with x relabeled as t .

- Ornstein–Uhlenbeck process is a Markov process.
- What does this actually mean \implies white board.

Ornstein-Uhlenbeck process

- The mean and covariance functions:

$$m(x) = 0$$

$$k(x, x') = \sigma^2 \exp(-\lambda|x - x'|)$$

- This has a *path representation* as a stochastic differential equation (SDE):

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t).$$

where $w(t)$ is a white noise process with x relabeled as t .

- Ornstein-Uhlenbeck process is a Markov process.
- What does this actually mean \implies white board.

Ornstein-Uhlenbeck process

- The mean and covariance functions:

$$m(x) = 0$$

$$k(x, x') = \sigma^2 \exp(-\lambda |x - x'|)$$

- This has a path representation as a stochastic differential equation (SDE):

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t)$$

$$\mathcal{F}(\omega) \hat{f} = -\lambda \hat{f} + \hat{w}$$

$$\hat{f} = \frac{\hat{w}}{\lambda + i\omega}$$

Spectral Density:

$$S(\omega) = \frac{E[\hat{w}^2]}{\omega^2 + \lambda^2} = \frac{2}{\omega^2 + \lambda^2}$$

where $w(t)$ is a white noise process with x relabeled as t .

- Ornstein-Uhlenbeck process is a Markov process.

- What does this actually mean \Rightarrow white board.

$$h(t) = \pi \int_{-\infty}^{\infty} \frac{1}{\omega^2 + \lambda^2} \exp(i\omega \tau) d\omega$$

$$\tau = t - t'$$

$$= \frac{2}{2\lambda} \exp(-\lambda |\tau|)$$

Residual transform.

=

Ornstein-Uhlenbeck process (cont.)

- Consider a Gaussian process regression problem

$$f(x) \sim \text{GP}(0, \sigma^2 \exp(-\lambda|x - x'|))$$

$$y_k = f(x_k) + \varepsilon_k$$

- This is equivalent to the state-space model

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t)$$

$$y_k = f(t_k) + \varepsilon_k$$

that is, with $f_k = f(t_k)$ we have a Gauss-Markov model

$$f_{k+1} \sim p(f_{k+1} | f_k)$$

$$y_k \sim p(y_k | f_k)$$

- Solvable in $O(n)$ time using Kalman filter/smoother.

Ornstein-Uhlenbeck process (cont.)

- Consider a Gaussian process regression problem

$$f(x) \sim \text{GP}(0, \sigma^2 \exp(-\lambda|x - x'|))$$
$$y_k = f(x_k) + \varepsilon_k$$

- This is equivalent to the state-space model

$$\frac{df(t)}{dt} = -\lambda f(t) + w(t)$$
$$y_k = f(t_k) + \varepsilon_k$$

that is, with $f_k = f(t_k)$ we have a Gauss-Markov model

$$f_{k+1} \sim p(f_{k+1} | f_k)$$
$$y_k \sim p(y_k | f_k)$$

- Solvable in $O(n)$ time using Kalman filter/smoother.

Ornstein-Uhlenbeck process (cont.)

- Consider a Gaussian process regression problem

$$f(x) \sim \text{GP}(0, \sigma^2 \exp(-\lambda|x - x'|))$$

$$y_k = f(x_k) + \varepsilon_k$$

- This is equivalent to the state-space model

$$\frac{df(t)}{dt}$$

$$= -\lambda f(t) + w(t)$$

$$y_k = f(t_k) + \varepsilon_k$$

that is, with $f_k = f(t_k)$ we have a Gauss-Markov model

$$f_{k+1} \sim p(f_{k+1} | f_k)$$

$$y_k \sim p(y_k | f_k)$$

- Solvable in $O(n)$ time using Kalman filter/smoothing.

* $e^{\lambda t}$ (integrating factor)

$$e^{\lambda t} \cdot \frac{df(t)}{dt} = \lambda f(t) e^{\lambda t} = w(t) e^{\lambda t}$$

$$\frac{d}{dt} (e^{\lambda t} f(t)) = e^{\lambda t} \cdot w(t)$$

$$e^{\lambda t} f(t) - e^{\lambda t_0} f(t_0) = \int_{t_0}^t w(s) e^{\lambda s} ds$$

$$f(t) = e^{-\lambda(t-t_0)} f(t_0) + \int_{t_0}^t e^{-\lambda(t-s)} w(s) ds$$

k.p.

$$\dot{f}_{k+1} = A \cdot f_k + q$$

$$q \sim N(0, Q)$$

$$p(f_{k+1} | f_k) = N(f_{k+1} | A f_k, Q)$$

$$p(f_{k+1} | f_k) = N(f_{k+1} | A f_k, Q)$$

$$Q = \int_{t_0}^t e^{-\lambda(t-s)} Q ds$$

State Space Form of Linear Time-Invariant SDEs

- Consider a N th order LTI SDE of the form

$$\frac{d^N f}{dt^N} + a_{N-1} \frac{d^{N-1} f}{dt^{N-1}} + \cdots + a_0 f = w(t).$$

- If we define $\mathbf{f} = (f, \dots, d^{N-1} f / dt^{N-1})$, we get a state space model:

$$\frac{d\mathbf{f}}{dt} = \underbrace{\begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ -a_0 & -a_1 & \dots & -a_{N-1} \end{pmatrix}}_{\mathbf{A}} \mathbf{f} + \underbrace{\begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}}_{\mathbf{L}} w(t)$$
$$f(t) = \underbrace{(1 \ 0 \ \dots \ 0)}_{\mathbf{H}} \mathbf{f}.$$

- The vector process $\mathbf{f}(t)$ is Markovian although $f(t)$ isn't.

State Space Form of Linear Time-Invariant SDEs

- Consider a N th order LTI SDE of the form

$$\frac{d^N f}{dt^N} + a_{N-1} \frac{d^{N-1} f}{dt^{N-1}} + \cdots + a_0 f = w(t).$$

- If we define $\mathbf{f} = (f, \dots, d^{N-1} f / dt^{N-1})$, we get a state space model:

$$\frac{d\mathbf{f}}{dt} = \underbrace{\begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ -a_0 & -a_1 & \dots & -a_{N-1} \end{pmatrix}}_{\mathbf{A}} \mathbf{f} + \underbrace{\begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}}_{\mathbf{L}} w(t)$$
$$f(t) = \underbrace{\begin{pmatrix} 1 & 0 & \dots & 0 \end{pmatrix}}_{\mathbf{H}} \mathbf{f}.$$

- The vector process $\mathbf{f}(t)$ is Markovian although $f(t)$ isn't.

State Space Form of Linear Time-Invariant SDEs

- Consider a N th order LTI SDE of the form

$$\frac{d^N f}{dt^N} + a_{N-1} \frac{d^{N-1} f}{dt^{N-1}} + \cdots + a_0 f = w(t).$$

- If we define $\mathbf{f} = (f, \dots, d^{N-1} f / dt^{N-1})$, we get a state space model:

$$\frac{d\mathbf{f}}{dt} = \underbrace{\begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ -a_0 & -a_1 & \dots & -a_{N-1} \end{pmatrix}}_{\mathbf{A}} \mathbf{f} + \underbrace{\begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}}_{\mathbf{L}} w(t)$$
$$f(t) = \underbrace{\begin{pmatrix} 1 & 0 & \dots & 0 \end{pmatrix}}_{\mathbf{H}} \mathbf{f}.$$

- The vector process $\mathbf{f}(t)$ is Markovian although $f(t)$ isn't.

Spectra of Linear Time-Invariant SDEs

- By taking the Fourier transform of the LTI SDE, we can derive the **spectral density** which has the form:

$$S(\omega) = \frac{(\text{constant})}{(\text{polynomial in } \omega^2)}$$

Handwritten: $\frac{1}{\omega^2 + \lambda^2}$

- We can also do this conversion to the **other direction**:

- With certain parameter values, the Matérn has the form:

$$S(\omega) \propto (\lambda^2 + \omega^2)^{-(\rho+1)}.$$

- Many non-rational spectral densities can be approximated:

$$S(\omega) = \sigma^2 \sqrt{\frac{\pi}{\kappa}} \exp\left(-\frac{\omega^2}{4\kappa}\right) \approx \frac{(\text{const})}{N!0!(4\kappa)^N + \dots + \omega^{2N}}$$

- For the conversion of a rational spectral density to a Markovian (state-space) model, we can use the **spectral factorization**.

Spectra of Linear Time-Invariant SDEs

- By taking the Fourier transform of the LTI SDE, we can derive the **spectral density** which has the form:

$$S(\omega) = \frac{(\text{constant})}{(\text{polynomial in } \omega^2)}$$

- We can also do this conversion to the **other direction**:
 - With certain parameter values, the **Matérn** has the form:

$$S(\omega) \propto (\lambda^2 + \omega^2)^{-(p+1)}.$$

- Many **non-rational** spectral densities can be approximated:

$$S(\omega) = \sigma^2 \sqrt{\frac{\pi}{\kappa}} \exp\left(-\frac{\omega^2}{4\kappa}\right) \approx \frac{(\text{const})}{N!/0!(4\kappa)^N + \dots + \omega^{2N}}$$

- For the conversion of a rational spectral density to a Markovian (state-space) model, we can use the **spectral factorization**.

Spectra of Linear Time-Invariant SDEs

- By taking the Fourier transform of the LTI SDE, we can derive the **spectral density** which has the form:

$$S(\omega) = \frac{(\text{constant})}{(\text{polynomial in } \omega^2)}$$

- We can also do this conversion to the **other direction**:
 - With certain parameter values, the **Matérn** has the form:

$$S(\omega) \propto (\lambda^2 + \omega^2)^{-(p+1)}.$$

- Many **non-rational** spectral densities can be approximated:

$$S(\omega) = \sigma^2 \sqrt{\frac{\pi}{\kappa}} \exp\left(-\frac{\omega^2}{4\kappa}\right) \approx \frac{(\text{const})}{N!/0!(4\kappa)^N + \dots + \omega^{2N}}$$

- For the conversion of a rational spectral density to a Markovian (state-space) model, we can use the **spectral factorization**.

Spectra of Linear Time-Invariant SDEs

- By taking the Fourier transform of the LTI SDE, we can derive the **spectral density** which has the form:

$$S(\omega) = \frac{(\text{constant})}{(\text{polynomial in } \omega^2)}$$

- We can also do this conversion to the **other direction**:
 - With certain parameter values, the **Matérn** has the form:

$$S(\omega) \propto (\lambda^2 + \omega^2)^{-(p+1)}.$$

- Many **non-rational** spectral densities can be approximated:

$$S(\omega) = \sigma^2 \sqrt{\frac{\pi}{\kappa}} \exp\left(-\frac{\omega^2}{4\kappa}\right) \approx \frac{(\text{const})}{N!/0!(4\kappa)^N + \dots + \omega^{2N}}$$

- For the conversion of a rational spectral density to a Markovian (state-space) model, we can use the **spectral factorization**.

Spectra of Linear Time-Invariant SDEs

- By taking the Fourier transform of the LTI SDE, we can derive the **spectral density** which has the form:

$$S(\omega) = \frac{(\text{constant})}{(\text{polynomial in } \omega^2)}$$

- We can also do this conversion to the **other direction**:
 - With certain parameter values, the **Matérn** has the form:

$$S(\omega) \propto (\lambda^2 + \omega^2)^{-(p+1)}.$$

- Many **non-rational** spectral densities can be approximated:


$$S(\omega) = \sigma^2 \sqrt{\frac{\pi}{\kappa}} \exp\left(-\frac{\omega^2}{4\kappa}\right) \approx \frac{(\text{const})}{N!/0!(4\kappa)^N + \dots + \omega^{2N}}$$

- For the conversion of a rational spectral density to a Markovian (state-space) model, we can use the **spectral factorization**.

State-space methods for Gaussian processes

- Approximation:

$$S(\omega) \approx \frac{b_0 + b_1 \omega^2 + \dots + b_M \omega^{2M}}{a_0 + a_1 \omega^2 + \dots + a_N \omega^{2N}}$$



- Results in a linear stochastic differential equation (SDE)


$$d\mathbf{f}(t) = \mathbf{A} \mathbf{f}(t) dt + \mathbf{L} d\mathbf{W}$$

- More generally stochastic evolution equations.
- $O(n)$ GP regression with Kalman filters and smoothers.
- Parallel block-sparse precision methods $\rightarrow O(\log n)$.

State-space methods for Gaussian processes

- Approximation:

$$S(\omega) \approx \frac{b_0 + b_1 \omega^2 + \dots + b_M \omega^{2M}}{a_0 + a_1 \omega^2 + \dots + a_N \omega^{2N}}$$



- Results in a linear stochastic differential equation (SDE)


$$d\mathbf{f}(t) = \mathbf{A} \mathbf{f}(t) dt + \mathbf{L} d\mathbf{W}$$

- More generally stochastic evolution equations.
- $O(n)$ GP regression with Kalman filters and smoothers.
- Parallel block-sparse precision methods $\rightarrow O(\log n)$.

State-space methods for Gaussian processes

- Approximation:

$$S(\omega) \approx \frac{b_0 + b_1 \omega^2 + \dots + b_M \omega^{2M}}{a_0 + a_1 \omega^2 + \dots + a_N \omega^{2N}}$$



- Results in a linear stochastic differential equation (SDE)


$$d\mathbf{f}(t) = \mathbf{A} \mathbf{f}(t) dt + \mathbf{L} d\mathbf{W}$$

- More generally stochastic evolution equations.
- $O(n)$ GP regression with Kalman filters and smoothers.
- Parallel block-sparse precision methods $\rightarrow O(\log n)$.

State-space methods for Gaussian processes

- Approximation:

$$S(\omega) \approx \frac{b_0 + b_1 \omega^2 + \dots + b_M \omega^{2M}}{a_0 + a_1 \omega^2 + \dots + a_N \omega^{2N}}$$



- Results in a linear stochastic differential equation (SDE)


$$d\mathbf{f}(t) = \mathbf{A} \mathbf{f}(t) dt + \mathbf{L} d\mathbf{W}$$

- More generally stochastic evolution equations.
- $O(n)$ GP regression with Kalman filters and smoothers.
- Parallel block-sparse precision methods $\rightarrow O(\log n)$.

State-space methods for Gaussian processes

- Approximation:

$$S(\omega) \approx \frac{b_0 + b_1 \omega^2 + \dots + b_M \omega^{2M}}{a_0 + a_1 \omega^2 + \dots + a_N \omega^{2N}}$$



- Results in a linear stochastic differential equation (SDE)

$$d\mathbf{f}(t) = \mathbf{A} \mathbf{f}(t) dt + \mathbf{L} d\mathbf{W}$$

- More generally stochastic evolution equations.
- $O(n)$ GP regression with Kalman filters and smoothers. *NIS*
- Parallel block-sparse precision methods $\rightarrow O(\log n)$.

State-space methods – temporal example

Example (Matérn class 1d)

The Matérn class of covariance functions is

$$k(t, t') = \sigma^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\frac{\sqrt{2\nu}}{\ell} |t - t'| \right)^\nu K_\nu \left(\frac{\sqrt{2\nu}}{\ell} |t - t'| \right).$$

When, e.g., $\nu = 3/2$, we have

$$\begin{aligned} d\mathbf{f}(t) &= \begin{pmatrix} 0 & 1 \\ -\lambda^2 & -2\lambda \end{pmatrix} \mathbf{f}(t) dt + \begin{pmatrix} 0 \\ q^{1/2} \end{pmatrix} dW(t), \\ f(t) &= (1 \quad 0) \mathbf{f}(t). \end{aligned}$$

State-space methods – spatio-temporal example

Example (2D Matérn covariance function)

- Consider a space-time Matérn covariance function

$$k(x, t; x', t') = \sigma^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\sqrt{2\nu} \frac{\rho}{l} \right)^\nu K_\nu \left(\sqrt{2\nu} \frac{\rho}{l} \right).$$

where we have $\rho = \sqrt{(t - t')^2 + (x - x')^2}$, $\nu = 1$ and $d = 2$.

- We get the following representation:

$$d\mathbf{f}(x, t) = \begin{pmatrix} 0 & 1 \\ \frac{\partial^2}{\partial x^2} - \lambda^2 & -2\sqrt{\lambda^2 - \frac{\partial^2}{\partial x^2}} \end{pmatrix} \mathbf{f}(x, t) dt + \begin{pmatrix} 0 \\ 1 \end{pmatrix} dW(x, t).$$

State-space methods – spatio-temporal example

Example (2D Matérn covariance function)

- Consider a space-time Matérn covariance function

$$k(x, t; x', t') = \sigma^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\sqrt{2\nu} \frac{\rho}{l} \right)^\nu K_\nu \left(\sqrt{2\nu} \frac{\rho}{l} \right).$$

where we have $\rho = \sqrt{(t - t')^2 + (x - x')^2}$, $\nu = 1$ and $d = 2$.

- We get the following representation:

$$d\mathbf{f}(x, t) = \begin{pmatrix} 0 & 1 \\ \frac{\partial^2}{\partial x^2} - \lambda^2 & -2\sqrt{\lambda^2 - \frac{\partial^2}{\partial x^2}} \end{pmatrix} \mathbf{f}(x, t) dt + \begin{pmatrix} 0 \\ 1 \end{pmatrix} dW(x, t).$$

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes**
- 4 Conclusion

Basic idea of SPDE inference on GPs [1/2]

- Consider e.g. the stochastic partial differential equation:

$$\frac{\partial^2 f(x, y)}{\partial x^2} + \frac{\partial^2 f(x, y)}{\partial y^2} - \lambda^2 f(x, y) = w(x, y)$$

- Fourier transforming gives the spectral density:

$$S(\omega_x, \omega_y) \propto \left(\lambda^2 + \omega_x^2 + \omega_y^2 \right)^{-2}.$$

- Inverse Fourier transform gives the covariance function:

$$k(x, y; x', y') = \frac{\sqrt{(x - x')^2 + (y - y')^2}}{2\lambda} K_1(\lambda \sqrt{(x - x')^2 + (y - y')^2})$$

- But this is just the Matérn covariance function.

Basic idea of SPDE inference on GPs [1/2]

- Consider e.g. the stochastic partial differential equation:

$$\frac{\partial^2 f(x, y)}{\partial x^2} + \frac{\partial^2 f(x, y)}{\partial y^2} - \lambda^2 f(x, y) = w(x, y)$$

- Fourier transforming gives the spectral density:

$$S(\omega_x, \omega_y) \propto \left(\lambda^2 + \omega_x^2 + \omega_y^2 \right)^{-2}.$$

- Inverse Fourier transform gives the covariance function:

$$k(x, y; x', y') = \frac{\sqrt{(x - x')^2 + (y - y')^2}}{2\lambda} K_1(\lambda \sqrt{(x - x')^2 + (y - y')^2})$$

- But this is just the Matérn covariance function.

Basic idea of SPDE inference on GPs [1/2]

- Consider e.g. the stochastic partial differential equation:

$$\frac{\partial^2 f(x, y)}{\partial x^2} + \frac{\partial^2 f(x, y)}{\partial y^2} - \lambda^2 f(x, y) = w(x, y)$$

- Fourier transforming gives the spectral density:

$$S(\omega_x, \omega_y) \propto \left(\lambda^2 + \omega_x^2 + \omega_y^2 \right)^{-2}.$$

- Inverse Fourier transform gives the covariance function:

$$k(x, y; x', y') = \frac{\sqrt{(x - x')^2 + (y - y')^2}}{2\lambda} K_1(\lambda \sqrt{(x - x')^2 + (y - y')^2})$$

- But this is just the Matérn covariance function.

Basic idea of SPDE inference on GPs [1/2]

- Consider e.g. the stochastic partial differential equation:

$$\frac{\partial^2 f(x, y)}{\partial x^2} + \frac{\partial^2 f(x, y)}{\partial y^2} - \lambda^2 f(x, y) = w(x, y)$$

- Fourier transforming gives the spectral density:

$$S(\omega_x, \omega_y) \propto \left(\lambda^2 + \omega_x^2 + \omega_y^2 \right)^{-2}.$$

- Inverse Fourier transform gives the covariance function:

$$k(x, y; x', y') = \frac{\sqrt{(x - x')^2 + (y - y')^2}}{2\lambda} K_1(\lambda \sqrt{(x - x')^2 + (y - y')^2})$$

- But this is just the Matérn covariance function.

Basic idea of SPDE inference on GPs [2/2]

- More generally, **SPDE** for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with **precision and covariance operators**:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$

$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- **Idea**: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:

- Finite-differences/FEM methods lead to sparse precision approximations.
- Fourier/basis-function methods lead to reduced rank covariance approximations.
- Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Basic idea of SPDE inference on GPs [2/2]

- More generally, SPDE for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with precision and covariance operators:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$

$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- Idea: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:

- Finite-differences/FEM methods lead to sparse precision approximations.
- Fourier/basis-function methods lead to reduced rank covariance approximations.
- Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Basic idea of SPDE inference on GPs [2/2]

- More generally, SPDE for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with precision and covariance operators:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$

$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- Idea: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:

- 1 Finite-differences/FEM methods lead to sparse precision approximations.
- 2 Fourier/basis-function methods lead to reduced rank covariance approximations.
- 3 Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Basic idea of SPDE inference on GPs [2/2]

- More generally, SPDE for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with precision and covariance operators:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$

$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- Idea: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:

- 1 Finite-differences/FEM methods lead to sparse precision approximations.
- 2 Fourier/basis-function methods lead to reduced rank covariance approximations.
- 3 Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Basic idea of SPDE inference on GPs [2/2]

- More generally, SPDE for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with precision and covariance operators:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$

$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- Idea: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:

- 1 Finite-differences/FEM methods lead to sparse precision approximations.
- 2 Fourier/basis-function methods lead to reduced rank covariance approximations.
- 3 Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Basic idea of SPDE inference on GPs [2/2]

- More generally, SPDE for some linear operator \mathcal{L} :

$$\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$$

- Now f is a GP with precision and covariance operators:

$$\mathcal{K}^{-1} = \mathcal{L}^* \mathcal{L}$$


$$\mathcal{K} = (\mathcal{L}^* \mathcal{L})^{-1}$$

- Idea: approximate \mathcal{L} or \mathcal{L}^{-1} using PDE/ODE methods:
 - 1 Finite-differences/FEM methods lead to sparse precision approximations.
 - 2 Fourier/basis-function methods lead to reduced rank covariance approximations.
 - 3 Spectral factorization leads to state-space (Kalman) methods which are time-recursive (or sparse in precision).

Finite-differences/FEM – sparse precision

- Basic idea:

$$\frac{\partial f(x)}{\partial x} \approx \frac{f(x+h) - f(x)}{h}$$
$$\frac{\partial^2 f(x)}{\partial x^2} \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$



- We get an SPDE approximation $\mathcal{L} \approx \mathbf{L}$, where \mathbf{L} is sparse
- The precision operator approximation is then sparse:


$$\mathcal{K}^{-1} \approx \mathbf{L}^T \mathbf{L} = \text{sparse}$$

- \mathcal{L} need to be approximated as integro-differential operator.
- Requires formation of a grid, but parallelizes well.

Finite-differences/FEM – sparse precision

- Basic idea:

$$\frac{\partial f(x)}{\partial x} \approx \frac{f(x+h) - f(x)}{h}$$
$$\frac{\partial^2 f(x)}{\partial x^2} \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$



- We get an SPDE approximation $\mathcal{L} \approx \mathbf{L}$, where \mathbf{L} is **sparse**
- The **precision operator** approximation is then **sparse**:


$$\mathcal{K}^{-1} \approx \mathbf{L}^T \mathbf{L} = \text{sparse}$$

- \mathcal{L} need to be approximated as **integro-differential operator**.
- Requires formation of a **grid**, but parallelizes well.

Finite-differences/FEM – sparse precision

- Basic idea:

$$\frac{\partial f(x)}{\partial x} \approx \frac{f(x+h) - f(x)}{h}$$
$$\frac{\partial^2 f(x)}{\partial x^2} \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$



- We get an SPDE approximation $\mathcal{L} \approx \mathbf{L}$, where \mathbf{L} is **sparse**
- The **precision operator** approximation is then **sparse**:


$$\mathcal{K}^{-1} \approx \mathbf{L}^T \mathbf{L} = \text{sparse}$$

- \mathcal{L} need to be approximated as **integro-differential operator**.
- Requires formation of a **grid**, but parallelizes well.

Finite-differences/FEM – sparse precision

- Basic idea:

$$\frac{\partial f(x)}{\partial x} \approx \frac{f(x+h) - f(x)}{h}$$
$$\frac{\partial^2 f(x)}{\partial x^2} \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$



- We get an SPDE approximation $\mathcal{L} \approx \mathbf{L}$, where \mathbf{L} is **sparse**
- The **precision operator** approximation is then **sparse**:


$$\mathcal{K}^{-1} \approx \mathbf{L}^T \mathbf{L} = \text{sparse}$$

- \mathcal{L} need to be approximated as **integro-differential operator**.
- Requires formation of a **grid**, but parallelizes well.

Finite-differences/FEM – sparse precision

- Basic idea:

$$\frac{\partial f(x)}{\partial x} \approx \frac{f(x+h) - f(x)}{h}$$
$$\frac{\partial^2 f(x)}{\partial x^2} \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$



- We get an SPDE approximation $\mathcal{L} \approx \mathbf{L}$, where \mathbf{L} is **sparse**
- The **precision operator** approximation is then **sparse**:

$$\mathcal{K}^{-1} \approx \mathbf{L}^T \mathbf{L} = \text{sparse}$$


- \mathcal{L} need to be approximated as **integro-differential operator**.
- Requires formation of a **grid**, but parallelizes well.

Classical and random Fourier methods – reduced rank approximations and FFT

- Approximation:

$$f(\mathbf{x}) \approx \sum_{\mathbf{k} \in \mathbb{N}^d} \alpha_{\mathbf{k}} \exp(2\pi i \mathbf{k}^T \mathbf{x})$$

$$\alpha_{\mathbf{k}} \sim \text{Gaussian}$$



- We use less coefficients $\alpha_{\mathbf{k}}$ than the number of data points.
- Leads to reduced-rank covariance approximations

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{|\mathbf{k}| \leq N} \sigma_{\mathbf{k}}^2 \exp(2\pi i \mathbf{k}^T \mathbf{x}) \exp(2\pi i \mathbf{k}^T \mathbf{x}')^*$$


- Truncated series, random frequencies, FFT, ...

Classical and random Fourier methods – reduced rank approximations and FFT

- Approximation:

$$f(\mathbf{x}) \approx \sum_{\mathbf{k} \in \mathbb{N}^d} \alpha_{\mathbf{k}} \exp(2\pi i \mathbf{k}^T \mathbf{x})$$

$$\alpha_{\mathbf{k}} \sim \text{Gaussian}$$



- We use **less coefficients** $\alpha_{\mathbf{k}}$ than the **number of data points**.
- Leads to **reduced-rank covariance approximations**

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{|\mathbf{k}| \leq N} \sigma_{\mathbf{k}}^2 \exp(2\pi i \mathbf{k}^T \mathbf{x}) \exp(2\pi i \mathbf{k}^T \mathbf{x}')^*$$


- Truncated series, random frequencies, FFT, ...

Classical and random Fourier methods – reduced rank approximations and FFT

- Approximation:

$$f(\mathbf{x}) \approx \sum_{\mathbf{k} \in \mathbb{N}^d} \alpha_{\mathbf{k}} \exp(2\pi i \mathbf{k}^T \mathbf{x})$$

$$\alpha_{\mathbf{k}} \sim \text{Gaussian}$$



- We use **less coefficients** $\alpha_{\mathbf{k}}$ than the **number of data points**.
- Leads to **reduced-rank covariance approximations**

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{|\mathbf{k}| \leq N} \sigma_{\mathbf{k}}^2 \exp(2\pi i \mathbf{k}^T \mathbf{x}) \exp(2\pi i \mathbf{k}^T \mathbf{x}')^*$$


- Truncated series, random frequencies, FFT, ...

Classical and random Fourier methods – reduced rank approximations and FFT

- Approximation:

$$f(\mathbf{x}) \approx \sum_{\mathbf{k} \in \mathbb{N}^d} \alpha_{\mathbf{k}} \exp(2\pi i \mathbf{k}^T \mathbf{x})$$

$$\alpha_{\mathbf{k}} \sim \text{Gaussian}$$



- We use **less coefficients** $\alpha_{\mathbf{k}}$ than the **number of data points**.
- Leads to **reduced-rank covariance approximations**

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{|\mathbf{k}| \leq N} \sigma_{\mathbf{k}}^2 \exp(2\pi i \mathbf{k}^T \mathbf{x}) \exp(2\pi i \mathbf{k}^T \mathbf{x}')^*$$


- Truncated series, random frequencies, FFT, ...

Hilbert-space/Galerkin methods – reduced rank approximations

- Approximation:

$$f(\mathbf{x}) \approx \sum_i c_i \phi_i(\mathbf{x})$$

$$\langle \phi_i, \phi_j \rangle_H \approx \delta_{ij}, \text{ e.g. } \nabla^2 \phi_i = -\lambda_i \phi_i$$



- Again, use less coefficients than the number of data points.
- Reduced-rank covariance approximations such as

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{i=1}^N \sigma_i^2 \phi_i(\mathbf{x}) \phi_i(\mathbf{x}').$$


- Wavelets, Galerkin, finite elements, ...

Hilbert-space/Galerkin methods – reduced rank approximations

- Approximation:

$$f(\mathbf{x}) \approx \sum_i c_i \phi_i(\mathbf{x})$$

$$\langle \phi_i, \phi_j \rangle_H \approx \delta_{ij}, \text{ e.g. } \nabla^2 \phi_i = -\lambda_i \phi_i$$



- Again, use **less coefficients** than the **number of data points**.
- Reduced-rank covariance approximations such as

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{i=1}^N \sigma_i^2 \phi_i(\mathbf{x}) \phi_i(\mathbf{x}').$$


- Wavelets, Galerkin, finite elements, ...

Hilbert-space/Galerkin methods – reduced rank approximations

- Approximation:

$$f(\mathbf{x}) \approx \sum_i c_i \phi_i(\mathbf{x})$$

$$\langle \phi_i, \phi_j \rangle_H \approx \delta_{ij}, \text{ e.g. } \nabla^2 \phi_i = -\lambda_i \phi_i$$



- Again, use **less coefficients** than the **number of data points**.
- **Reduced-rank covariance approximations** such as

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{i=1}^N \sigma_i^2 \phi_i(\mathbf{x}) \phi_i(\mathbf{x}').$$


- Wavelets, Galerkin, finite elements, ...

Hilbert-space/Galerkin methods – reduced rank approximations

- Approximation:

$$f(\mathbf{x}) \approx \sum_i c_i \phi_i(\mathbf{x})$$

$$\langle \phi_i, \phi_j \rangle_H \approx \delta_{ij}, \text{ e.g. } \nabla^2 \phi_i = -\lambda_i \phi_i$$



- Again, use **less coefficients** than the **number of data points**.
- **Reduced-rank covariance approximations** such as

$$k(\mathbf{x}, \mathbf{x}') \approx \sum_{i=1}^N \sigma_i^2 \phi_i(\mathbf{x}) \phi_i(\mathbf{x}').$$

- Wavelets, Galerkin, finite elements, ...

Contents

- 1 Basic ideas
- 2 Stochastic differential equations and Gaussian processes
- 3 Stochastic partial differential equations and Gaussian processes
- 4 Conclusion

Back to SPDE representations of GPs

GP model $\mathbf{x} \in \mathbb{R}^d, t \in \mathbb{R}$	Equivalent S(P)DE model
Spatial $k(\mathbf{x}, \mathbf{x}')$	SPDE model (\mathcal{L} is an operator) $\mathcal{L} f(\mathbf{x}) = w(\mathbf{x})$
Temporal $k(t, t')$	State-space/SDE model $\frac{d\mathbf{f}(t)}{dt} = \mathbf{A} \mathbf{f}(t) + \mathbf{L} w(t)$
Spatio-temporal $k(\mathbf{x}, t; \mathbf{x}', t')$	Stochastic evolution equation $\frac{\partial}{\partial t} \mathbf{f}(\mathbf{x}, t) = \mathcal{A}_x \mathbf{f}(\mathbf{x}, t) + \mathbf{L} w(\mathbf{x}, t)$

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFM.
- Inverse problems – operators in the measurement model.

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFMs.
- Inverse problems – operators in the measurement model.

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFM.
- Inverse problems – operators in the measurement model.

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFM.
- Inverse problems – operators in the measurement model.

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFMs.
- Inverse problems – operators in the measurement model.

What then?

- Exchange and map approximations between the fields:
 - Inducing points \leftrightarrow point-collocation; spectral methods \leftrightarrow Galerkin methods; finite-differences \leftrightarrow GMRFs;
- Non-Gaussian processes: Student's-t processes, non-linear Itô processes, jump processes, hybrid point/Gaussian processes.
- Hierarchical (deep) SPDE models: we stack SPDEs on top of each other – the SPDE just becomes non-linear.
- Combined first-principles and nonparametric models – latent force models (LFM), also non-linear and non-Gaussian LFMs.
- Inverse problems – operators in the measurement model.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Summary

- Gaussian processes (GPs) are nice, but the computational scaling is bad.
- GPs have representations as solutions to SPDEs.
- In temporal models we can use Kalman/Bayesian filters and smoothers.
- SPDE methods can be used to speed up GP inference.
- New paths towards non-linear GP models.
- Work nicely in latent force models.

Useful references

- **N. Wiener (1950).** *Extrapolation, Interpolation and Smoothing of Stationary Time Series with Engineering Applications.* John Wiley & Sons, Inc.
- **R. L. Stratonovich (1963).** *Topics in the Theory of Random Noise.* Gordon and Breach.
- **J. Hartikainen and S. Särkkä (2010).** *Kalman Filtering and Smoothing Solutions to Temporal Gaussian Process Regression Models.* Proc. MLSP.
- **S. Särkkä, A. Solin, and J. Hartikainen (2013).** *Spatio-Temporal Learning via Infinite-Dimensional Bayesian Filtering and Smoothing.* IEEE Sig.Proc.Mag., 30(5):51–61.
- **S. Särkkä (2013).** *Bayesian Filtering and Smoothing.* Cambridge University Press.
- **S. Särkkä and A. Solin (2017, to appear)** *Applied Stochastic Differential Equations.* Cambridge University Press.