


S-PLUS: An atlas of integrated $H\alpha$ fluxes for planetary nebulae in the Magellanic Clouds

L. A. Gutiérrez-Soto, A. R. Lopes, A. V. Smith Castelli & F. R. Faifer
Instituto de Astrofísica de La Plata, CONICET-UNLP, Argentina

ABSTRACT

We present an atlas of $H\alpha$ fluxes for planetary nebulae of the Magellanic Clouds (MC PNe) with measurements from the Southern Photometric Local Universe Survey (S-PLUS), a 12 band (7 narrow and 5 broad) imaging survey that allows us to perform an spatial analysis of the $H\alpha$ emission. Aperture photometry on the continuum-subtracted images was performed to extract $H\alpha + [N II]$ fluxes of the MC PNe observed by S-PLUS. The dust attenuation and $[N II]$ contribution was corrected with empirical relations. Amongst its many applications, it can provide baseline data for photoionization and hydrodynamical modelling, and allow better estimates of Zanstra temperatures for PN central stars with accurate optical photometry.

Ring Nebula (Messier 57)




Credits: NASA, ESA and the Hubble Heritage (STScI/AURA)-ESA/Hubble Collaboration

What are planetary nebulae?


Planetary nebulae (PNe) are emission line nebulae that represent a short phase in the late evolution of low- and intermediate-mass stars.

About S-PLUS:



As part of its effort to map 9,000 square degrees of the Southern Hemisphere, the S-PLUS (Mendes de Oliveira et al. 2019) project has a crucial feature: images of the entire field captured using the $H\alpha$ narrow-band J0660 filter.


PNe in the Magellanic Clouds



The distribution of the planetary nebulae (green circles) in the Magellanic Clouds can be seen in the figure. The large gray squares indicate the S-PLUS fields that cover both the Large and Small Magellanic Clouds. Almost all the PNe are within the S-PLUS fields.

Extracting $H\alpha$ flux from photometric data

The $H\alpha$ flux for the Magellanic Cloud planetary nebulae observed by S-PLUS was extracted using two broad-band filters (r and i) and one narrow-band filter (J0660), according to Equation 3 of Vilella-Rojo et al. (2015). Below are the $H\alpha$ maps for some planetary nebulae, along with their respective S-PLUS spectra and false color images based on the r, J0660, and i filters:



References

Mendes de Oliveira C. et al., 2019, MNRAS, 489, 241
Vilella-Rojo, G. et al. 2015, A&A, 580, A47

✉ gsotoangel@fcaglp.unlp.edu.ar