


L. A. Gutiérrez-Soto, A. R. Lopes, A. V. Smith Castelli, F. R. Faifer & R. Haack Instituto de Astrofísica de La Plata, CONICET-UNLP, Argentina

18th S-PLUS Collaboration Meeting August 21-23, 2023, Rio de Janeiro, Brasil


## **Context**



#### What are planetary nebulae?

Planetary nebulae (PNe) are emission line nebulae that represent a short phase in the late evolution of low- and intermediatemass stars.


#### PNe in the Magellanic Clouds



The distribution of the planetary nebulae from the literature (green circles) in the Magellanic Clouds.

#### About S-PLUS:

Typical spectrum of a PN: the  $H\alpha$  and [N II] emission lines are detected by J0660 filter for sources with a redshift up to 0.015




As part of its effort to map 9,000 square degrees of the Southern Hemisphere, the S-PLUS project (Mendes de Oliveira et al. 2019) has a crucial feature: images of the entire field captured using the Hα narrow-band **J0660** filter.

# Hα flux PNe with S-PLUS data

Wavelength (A)

### Extracting $H\alpha$ flux from photometric data



Using S-PLUS data, we analyze Hα flux in planetary nebulae. This helps us understand their emission properties, ionization processes, and chemical composition, shedding light on these astronomical objects.