




Mapping H α -Excess Candidate Point Sources in the Southern Hemisphere Using S-PLUS Data

Luis A. Gutiérrez-Soto & S-PLUS collaboration

**19th S-PLUS Collaboration Meeting
August 19-21, 2024**

Context





As part of mapping 9000 deg² of the Southern Hemisphere, the S-PLUS project surveys the H α transition, capturing a wide range of astrophysical processes. In this study, we utilize DR4 data, covering 171 low-galactic-latitude fields, 341 fields from the Main Survey, and 150 fields in the Magellanic Clouds region, to highlight H α -excess point sources using the $(r - \text{J0660})$ vs. $(r - i)$ color-color diagram.




Selecting the H α Excess Sources

The identification of H α emitters was based on the method applied by Witham et al. (2008)

Iteratively fit the main stellar locus and select all H α -excess sources which lie more than 4σ away from the fitted locus



Selecting the H α Excess Sources



Galactic Disk Challenges

Stellar Loci Duplication:

Disk fields show two stellar loci due to differential reddening and mixed populations (**main-sequence stars** and **giants**).


Fitting Issues:

Initial fit often aligns with the reddened locus, complicating H-alpha excess detection.

Solution:

Adjusted fit upward to align with the unreddened sequence.

Reverted to the original fit if adjustments worsened the accuracy.



Selecting the H α Excess Sources

Identification:

Sources significantly above the locus of main-sequence and giant stars exhibit excess in the J0660 filter, attributed to the H-alpha line.


Current Data:

Main Survey: 3,637 outliers identified.


Galactic Disk: 3,734 outliers identified.

Types of Sources:





In agreement with SIMBAD, these include emission-line stars, PNe, CVs, SySt, YSOs, Be stars, QSOs, and galaxies.



Example of two H α Emitters





Extragalactic Compact Sources



Our selection criteria effectively isolated compact sources, minimizing the inclusion of extended sources. In agreement with SIMBAD, only 23 AGN (3.1%) and 9 galaxies (1.2%) were identified in the main survey, while 143 QSOs (19.6%) were selected. The spectra of these QSOs and Seyfert galaxy show other emission lines that fall into the J0660 filter, resulting in a significant H α excess.


Machine Learning Approaches

We applied **UMAP** for dimensionality reduction on S-PLUS data and used **HDBSCAN** for clustering H α excess sources. Two experiments were conducted: one using only the **66 S-PLUS** colors and another incorporating WISE bands. The results revealed the formation of five distinct groups, effectively separating galactic and extragalactic sources.



Machine Learning Approaches


We applied **UMAP** for dimensionality reduction on S-PLUS data and used **HDBSCAN** for clustering H α excess sources. Two experiments were conducted: one using only the **66 S-PLUS** colors and another incorporating WISE bands. The results revealed the formation of five distinct groups, effectively separating galactic and extragalactic sources.



Extracting Main Features: Color Analysis


Analysis: Key colors from S-PLUS and WISE filters identified by **Random Forest** as most influential in classifying H-alpha excess objects.

Results: **Top 20 features** contribute significantly to distinguishing between different object classes identified by UMAP + HDBSCAN.



S-PLUS: H α emitters

Extracting Main Features: Color Analysis





Conclusions

- **Identification of H-alpha Excess Candidates:**

- ✓ 7,371 candidates identified using the S-PLUS J0660 filter.
- ✓ Distribution: 3,637 in the high-latitude main survey and 3,734 in the Galactic disk.

- **Classification of Objects:**

- ✓ Cross-referencing with SIMBAD: EM stars, YSOs, Be stars, CVs, PNe, QSOs, non-local galaxies, and RR Lyrae stars.

- **Machine Learning Techniques:**

- ✓ UMAP and HDBSCAN: Effective differentiation between Galactic and extragalactic objects
- ✓ Challenges in distinguishing cataclysmic variables from QSOs or AGN with $z \approx 1.35$.

- **Integration of WISE Data:**

- ✓ Improved separation of extragalactic and Galactic sources.

- **Random Forest Model:**

- ✓ WISE data crucial for identifying significant features.
- ✓ Improved color-color diagrams and understanding of H-alpha related phenomena.

THANK!!