

Computational Astrophysics

E. Larrañaga

Observatorio Astronómico Nacional
Universidad Nacional de Colombia

April 20, 2019

Outline

1 Integration]

2 Integration based on Piecewise Polynomial Interpolation

- MidPoint Rule
- Trapezoidal Rule
- Simpson's Rule
- Simpson's Rule (non-equidistant intervals)

3 Gaussian Quadrature

- Change of Interval

Integration

We want to evaluate numerically the integral

$$Q = \int_a^b f(x) dx , \quad (1)$$

where $f(x)$ may be a well behaved analytical function or a function given as discrete data $f(x_i)$.

Integration based on Piecewise Polynomial Interpolation

Suppose that we know the values of $f(x)$ at a finite set of points (nodes) $\{x_j\}$, ($j = 0, \dots, n$), in the interval $[a, b]$.

Integration based on Piecewise Polynomial Interpolation

Suppose that we know the values of $f(x)$ at a finite set of points (nodes) $\{x_j\}$, ($j = 0, \dots, n$), in the interval $[a, b]$.

It is possible to replace $f(x)$ with an interpolated polynomial $p(x)$ whose analytic integral is known.

Integration based on Piecewise Polynomial Interpolation

Suppose that we know the values of $f(x)$ at a finite set of points (nodes) $\{x_j\}$, ($j = 0, \dots, n$), in the interval $[a, b]$.

It is possible to replace $f(x)$ with an interpolated polynomial $p(x)$ whose analytic integral is known.

This integration, based on interpolation polynomials, is generally called *Newton-Cotes quadrature formulae*.

Integration based on Piecewise Polynomial Interpolation

Now, consider that the full interval over which we intend to integrate can be broken down into N sub-intervals $[a_i, b_i]$ that encompass $N + 1$ nodes x_i ($i = 0, \dots, N$) at which we know the integrand $f(x)$.

Integration based on Piecewise Polynomial Interpolation

Now, consider that the full interval over which we intend to integrate can be broken down into N sub-intervals $[a_i, b_i]$ that encompass $N + 1$ nodes x_i ($i = 0, \dots, N$) at which we know the integrand $f(x)$.

The full integral Q can be written as the sum of the sub-integrals Q_i :

$$Q = \sum_{i=0}^{N-1} Q_i = \sum_{i=0}^{N-1} \int_{a_i}^{b_i} f(x) dx . \quad (2)$$

Integration based on Piecewise Polynomial Interpolation

Now, consider that the full interval over which we intend to integrate can be broken down into N sub-intervals $[a_i, b_i]$ that encompass $N + 1$ nodes x_i ($i = 0, \dots, N$) at which we know the integrand $f(x)$.

The full integral Q can be written as the sum of the sub-integrals Q_i :

$$Q = \sum_{i=0}^{N-1} Q_i = \sum_{i=0}^{N-1} \int_{a_i}^{b_i} f(x) dx . \quad (2)$$

There are many ways to evaluate these sub-integrals.

MidPoint Rule

The simplest approximation is to assume that the function $f(x)$ is constant on the interval $[a_i, b_i]$.

MidPoint Rule

The simplest approximation is to assume that the function $f(x)$ is constant on the interval $[a_i, b_i]$.

Hence, using its central value we write

$$Q_i = \int_{a_i}^{b_i} f(x) dx = (b_i - a_i) f\left(\frac{a_i + b_i}{2}\right) . \quad (3)$$

MidPoint Rule

The simplest approximation is to assume that the function $f(x)$ is constant on the interval $[a_i, b_i]$.

Hence, using its central value we write

$$Q_i = \int_{a_i}^{b_i} f(x) dx = (b_i - a_i) f\left(\frac{a_i + b_i}{2}\right) . \quad (3)$$

This is also called the “rectangle rule” and is an open Newton-Cotes formula.

MidPoint Rule

The simplest approximation is to assume that the function $f(x)$ is constant on the interval $[a_i, b_i]$.

Hence, using its central value we write

$$Q_i = \int_{a_i}^{b_i} f(x) dx = (b_i - a_i) f\left(\frac{a_i + b_i}{2}\right) . \quad (3)$$

This is also called the “rectangle rule” and is an open Newton-Cotes formula.

Note that we need to be able to evaluate $f(x)$ at the midpoint (knowing the value or using an interpolating polynomial).

MidPoint Rule

The error in the midpoint quadrature can be estimated using a Taylor expansion about the midpoint $m_i = (a_i + b_i)/2$,

$$f(x) = f(m_i) + f'(m_i)(x - m_i) + \frac{f''(m_i)}{2}(x - m_i)^2 + \frac{f'''(m_i)}{6}(x - m_i)^3 + \dots \quad (4)$$

Integrating this expression from a_i to b_i , the odd-order terms drop out, and what is left is

$$Q_i = \int_{a_i}^{b_i} f(x) dx = f(m_i)(b_i - a_i) + \frac{f''(m_i)}{24}(b_i - a_i)^3 + \dots \quad (5)$$

$$Q_i = \int_{a_i}^{b_i} f(x) dx = f(m_i)(b_i - a_i) + \mathcal{O}(h^3) + \dots \quad (6)$$

where $h = (b_i - a_i) \approx \frac{(b-a)}{N}$.

Trapezoidal Rule

This method approximates $f(x)$ with a linear polynomial on $[a_i, b_i]$,

$$Q_i = \int_{a_i}^{b_i} f(x) dx = \frac{1}{2}(b_i - a_i) [f(b_i) + f(a_i)] . \quad (7)$$

Trapezoidal Rule

This method approximates $f(x)$ with a linear polynomial on $[a_i, b_i]$,

$$Q_i = \int_{a_i}^{b_i} f(x) dx = \frac{1}{2}(b_i - a_i) [f(b_i) + f(a_i)] . \quad (7)$$

Using a Taylor expansion it is shown that the trapezoidal rule has a local error that, like the midpoint rule, goes with h^3 .

Simpson's Rule

Simpson's Rule approximates $f(x)$ on $[a_i, b_i]$ by a polynomial of degree 2.

Simpson's Rule

Simpson's Rule approximates $f(x)$ on $[a_i, b_i]$ by a polynomial of degree 2.

Writing out the interpolating polynomial and integrating we get

$$Q_i = \frac{b_i - a_i}{6} \left[f(a_i) + 4f\left(\frac{a_i + b_i}{2}\right) + f(b_i) \right] + \mathcal{O}([b_i - a_i]^5) . \quad (8)$$

Simpson's Rule

Simpson's Rule approximates $f(x)$ on $[a_i, b_i]$ by a polynomial of degree 2.

Writing out the interpolating polynomial and integrating we get

$$Q_i = \frac{b_i - a_i}{6} \left[f(a_i) + 4f\left(\frac{a_i + b_i}{2}\right) + f(b_i) \right] + \mathcal{O}([b_i - a_i]^5) . \quad (8)$$

Note that it is needed the value $f((a_i + b_i)/2)$ and that a special care must be taken at the boundaries.

Simpson's Rule

Simpson's Rule approximates $f(x)$ on $[a_i, b_i]$ by a polynomial of degree 2.

Writing out the interpolating polynomial and integrating we get

$$Q_i = \frac{b_i - a_i}{6} \left[f(a_i) + 4f\left(\frac{a_i + b_i}{2}\right) + f(b_i) \right] + \mathcal{O}([b_i - a_i]^5) . \quad (8)$$

Note that it is needed the value $f((a_i + b_i)/2)$ and that a special care must be taken at the boundaries.

This method has an error going with h^5

Simpson's Rule (non-equidistant intervals)

If $f(x)$ is non-equidistantly sampled, i.e. $h = [a_i, b_i]$ is not constant, Simpson's rule modifies.

Simpson's Rule (non-equidistant intervals)

If $f(x)$ is non-equidistantly sampled, i.e. $h = [a_i, b_i]$ is not constant, Simpson's rule modifies.

Write

$$f(x) = ax^2 + bx + c \quad \text{for } x \in [x_{i-1}, x_{i+1}] , \quad (9)$$

where,

$$\begin{aligned} a &= \frac{h_{i-1}f(x_{i+1}) - (h_{i-1} + h_i)f(x_i) + h_if(x_{i-1})}{h_{i-1}h_i(h_{i-1} + h_i)} , \\ b &= \frac{h_{i-1}^2f(x_{i+1}) + (h_i^2 - h_{i-1}^2)f(x_i) - h_i^2f(x_{i-1})}{h_{i-1}h_i(h_{i-1} + h_i)} , \\ c &= f(x_i) , \end{aligned} \quad (10)$$

with $h_i = x_{i+1} - x_i$ and $h_{i-1} = x_i - x_{i-1}$.

Simpson's Rule (non-equidistant intervals)

Since the integral

$$\int_{x_{i-1}}^{x_{i+1}} f(x) dx$$

is independent of the origin of the coordinates, we can choose $x_i = 0$, and hence $-h_{i-1} = x_{i-1}$, $h_i = x_{i+1}$ to obtain

$$\int_{x_{i-1}}^{x_{i+1}} f(x) dx = \int_{-h_{i-1}}^{h_i} f(x) dx = \alpha f(x_{i+1}) + \beta f(x_i) + \gamma f(x_{i-1}) , \quad (11)$$

with

$$\begin{aligned} \alpha &= \frac{2h_i^2 + h_i h_{i-1} - h_{i-1}^2}{6h_i} , & \beta &= \frac{(h_i + h_{i-1})^3}{6h_i h_{i-1}} , \\ \gamma &= \frac{-h_i^2 + h_i h_{i-1} + 2h_{i-1}^2}{6h_{i-1}} , \end{aligned} \quad (12)$$

Gaussian Quadrature

Gaussian quadrature is an integration method in which there are chosen two quantities, nodes and weights, to maximize the degree of the resulting integration rule.

Gaussian Quadrature

Gaussian quadrature is an integration method in which there are chosen two quantities, nodes and weights, to maximize the degree of the resulting integration rule.

Using $2n$ degrees of freedom, it is possible to integrate polynomials of degree $2n - 1$. The condition is that, for a set of n nodes x_i and n weights w_i at which a function f is known,

$$\int_a^b f(x) dx \approx \sum_{i=1}^n w_i f(x_i) \quad (13)$$

2-point Gaussian Quadrature

In the 2-point Gaussian quadrature we consider 2 nodes x_i and 2 weights w_i on the interval $[a, b]$ to integrate exactly a polynomial of degree 3. Hence

$$\int_a^b f(x)dx = w_1 f(x_1) + w_2 f(x_2). \quad (14)$$

We can find the unknowns w_1 , w_2 , x_1 , and x_2 by demanding that the formula give exact results for integrating a general polynomial of degree 3,

$$\begin{aligned} \int_a^b f(x)dx &= \int_a^b (c_0 + c_1x + c_2x^2 + c_3x^3)dx, \\ &= c_0(b-a) + c_1 \left(\frac{b^2 - a^2}{2} \right) + c_2 \left(\frac{b^3 - a^3}{3} \right) + c_3 \left(\frac{b^4 - a^4}{4} \right). \end{aligned} \quad (15)$$

2-point Gaussian Quadrature

However, we have

$$\begin{aligned}\int_a^b f(x) dx &= w_1 f(x_1) + w_2 f(x_2) \\ &= w_1(c_0 + c_1 x_1 + c_2 x_1^2 + c_3 x_1^3) + w_2(c_0 + c_1 x_2 + c_2 x_2^2 + c_3 x_2^3).\end{aligned}$$

Therefore, comparing these results gives

$$w_1 = \frac{b-a}{2}, \quad w_2 = \frac{b-a}{2}, \quad (16)$$

$$x_1 = \left(\frac{b-a}{2}\right) \left(-\frac{1}{\sqrt{3}}\right) + \left(\frac{b+a}{2}\right), \quad (17)$$

$$x_2 = \left(\frac{b-a}{2}\right) \left(\frac{1}{\sqrt{3}}\right) + \left(\frac{b+a}{2}\right). \quad (18)$$

2-point Gaussian Quadrature

In the special case of the interval $[-1, 1]$:

$$w_1 = w_2 = 1, \quad x_1 = -\frac{1}{\sqrt{3}}, \quad x_2 = \frac{1}{\sqrt{3}}, \quad (19)$$

and

$$\int_{-1}^1 f(x) dx = f\left(-\frac{1}{\sqrt{3}}\right) + f\left(\frac{1}{\sqrt{3}}\right). \quad (20)$$

Change of Interval

Using an affine transformation that maps $[a, b] \rightarrow [-1, 1]$ as

$$t = \frac{b-a}{2}x + \frac{a+b}{2}, \quad (21)$$

and a change of variables

$$dt = \frac{b-a}{2}dx, \quad (22)$$

it is possible to change the interval of integration,

$$\int_a^b f(x)dx = \frac{b-a}{2} \int_{-1}^1 f\left(\frac{b-a}{2}x + \frac{a+b}{2}\right) dx. \quad (23)$$

Cubic Natural Spline Interpolation

Integration of this interpolated second derivative two times and using $f_i = p_i(x_i) = f(x_i)$ gives

$$p_i(x) = \alpha_i(x - x_i)^3 + \beta_i(x - x_{i+1})^3 + \gamma_i(x - x_i) + \eta_i(x - x_{i+1}), \quad (24)$$

where

$$\alpha_i = \frac{p''_{i+1}}{6h_i} \qquad \beta_i = \frac{-p''_i}{6h_i}, \quad (25)$$

$$\gamma_i = \frac{f_{i+1}}{h_i} - \frac{h_i p''_{i+1}}{6} \qquad \eta_i = \frac{h_i p''_i}{6} - \frac{f_i}{h_i}, \quad (26)$$

with $h_i = x_{i+1} - x_i$.

Cubic Natural Spline Interpolation

Integration of this interpolated second derivative two times and using $f_i = p_i(x_i) = f(x_i)$ gives

$$p_i(x) = \alpha_i(x - x_i)^3 + \beta_i(x - x_{i+1})^3 + \gamma_i(x - x_i) + \eta_i(x - x_{i+1}), \quad (24)$$

where

$$\alpha_i = \frac{p''_{i+1}}{6h_i} \qquad \beta_i = \frac{-p''_i}{6h_i}, \quad (25)$$

$$\gamma_i = \frac{f_{i+1}}{h_i} - \frac{h_i p''_{i+1}}{6} \qquad \eta_i = \frac{h_i p''_i}{6} - \frac{f_i}{h_i}, \quad (26)$$

with $h_i = x_{i+1} - x_i$.

Hence, all we need to find the spline is to know all of its second derivatives $p''_i = p''_i(x_i)$.

Cubic Natural Spline Interpolation

Now apply the condition $p'_{i-1}(x_i) = p'_i(x_i)$ to the interpolated polynomial to obtain the equation

$$h_{i-1}p''_{i-1} + 2(h_{i-1} + h_i)p''_i + h_i p''_{i+1} = 6 \left(\frac{g_i}{h_i} - \frac{g_{i-1}}{h_{i-1}} \right), \quad (27)$$

where $g_i = f_{i+1} - f_i$.

Cubic Natural Spline Interpolation

Now apply the condition $p'_{i-1}(x_i) = p'_i(x_i)$ to the interpolated polynomial to obtain the equation

$$h_{i-1}p''_{i-1} + 2(h_{i-1} + h_i)p''_i + h_i p''_{i+1} = 6 \left(\frac{g_i}{h_i} - \frac{g_{i-1}}{h_{i-1}} \right), \quad (27)$$

where $g_i = f_{i+1} - f_i$.

This is a linear system with $n - 1$ unknowns: p''_i for $i = 1, \dots, n - 1$ (remember that $p''_0 = p''_n = 0$, as set by the natural spline condition).

Cubic Natural Spline Interpolation

Defining $d_i = 2(h_{i-1} + h_i)$ and $b_i = 6 \left(\frac{g_i}{h_i} - \frac{g_{i-1}}{h_{i-1}} \right)$, we can write

$$Ap'' = b \quad \text{with } A_{ij} = \begin{cases} d_i & \text{if } i = j, \\ h_i & \text{if } i = j - 1, \\ h_{i-1} & \text{if } i = j + 1, \\ 0 & \text{otherwise.} \end{cases} \quad (28)$$

Cubic Natural Spline Interpolation

Defining $d_i = 2(h_{i-1} + h_i)$ and $b_i = 6 \left(\frac{g_i}{h_i} - \frac{g_{i-1}}{h_{i-1}} \right)$, we can write

$$Ap'' = b \quad \text{with } A_{ij} = \begin{cases} d_i & \text{if } i = j, \\ h_i & \text{if } i = j - 1, \\ h_{i-1} & \text{if } i = j + 1, \\ 0 & \text{otherwise.} \end{cases} \quad (28)$$

The coefficient matrix A_{ij} is real, symmetric, and tri-diagonal.

The solution of such a linear system of equations will be presented later in the course.