

3.3.4

### TVD methods for conservation laws:

def: Total Variation of a grid function

consider grid function  $\vec{v}^n = \begin{bmatrix} v_1^n \\ \vdots \\ v_m^n \end{bmatrix}$  on a

grid with  $m$  spatial grid points at time  $t_n$ ,

and assume periodic boundary conditions ( $v_m^n = v_0^n$ )

then the total variation of  $\vec{v}^n$  is defined by

$$TV(\vec{v}^n) = \sum_{i=1}^m |v_i^n - v_{i-1}^n|$$

idea: if we want to preclude spurious numerical oscillations, we can require a TVD property for the numerical method:

$$TV(\vec{v}^{n+1}) \leq TV(\vec{v}^n)$$

(since the exact solution also satisfies such a property)

when is a two-level scheme TVD?

consider the general form:

$$v_i^{n+1} = v_i^n - C_{i-\frac{1}{2}} (v_i^n - v_{i-1}^n) + D_{i+\frac{1}{2}} (v_{i+1}^n - v_i^n)$$

use notation:  $\Delta^- v_i^n = v_i^n - v_{i-1}^n$

$$\Delta^+ v_i^n = v_{i+1}^n - v_i^n$$

then we can derive conditions on the coefficients  $C_{i-\frac{1}{2}}$  and  $D_{i+\frac{1}{2}}$  for TVD:

### Theorem 3.22: TVD Conditions

Consider numerical method

$$v_i^{n+1} = v_i^n - C_{i-\frac{1}{2}} \Delta^- v_i^n + D_{i+\frac{1}{2}} \Delta^+ v_i^n, \quad (3.66)$$

with periodic boundary conditions. If, for all  $i$ ,

$$\begin{cases} C_{i+\frac{1}{2}} \geq 0, \\ D_{i+\frac{1}{2}} \geq 0, \\ C_{i+\frac{1}{2}} + D_{i+\frac{1}{2}} \leq 1 \end{cases} \quad (3.67)$$

then the numerical method is TVD.

**Theorem 3.22: TVD Conditions**

Consider numerical method

$$v_i^{n+1} = v_i^n - C_{i-\frac{1}{2}} \Delta^- v_i^n + D_{i+\frac{1}{2}} \Delta^+ v_i^n, \quad (3.66)$$

with periodic boundary conditions. If, for all  $i$ ,

$$\begin{cases} C_{i+\frac{1}{2}} \geq 0, \\ D_{i+\frac{1}{2}} \geq 0, \\ C_{i+\frac{1}{2}} + D_{i+\frac{1}{2}} \leq 1 \end{cases} \quad (3.67)$$

then the numerical method is TVD.

$$\Delta^- v_i^n = v_i^n - v_{i-1}^n$$

$$\Delta^+ v_i^n = v_{i+1}^n - v_i^n$$

proof:  $TV(\bar{v}^{n+1}) = \sum_i |v_{i+1}^{n+1} - v_i^{n+1}|$

$$= \sum_i \left| \Delta^+ v_i^n - C_{i+\frac{1}{2}} \underbrace{\Delta^- v_{i+1}^n}_{\Delta^+ v_i^n} + D_{i+\frac{3}{2}} \Delta^+ v_{i+1}^n + C_{i-\frac{1}{2}} \underbrace{\Delta^- v_i^n}_{\Delta^+ v_{i-1}^n} - D_{i+\frac{1}{2}} \Delta^+ v_i^n \right|$$

$$\leq \sum_i \left| (1 - C_{i+\frac{1}{2}} - D_{i+\frac{1}{2}}) \Delta^+ v_i^n + |D_{i+\frac{3}{2}} \Delta^+ v_{i+1}^n| + |C_{i-\frac{1}{2}} \Delta^+ v_{i-1}^n| \right|$$

$$= \sum_i \left| (1 - C_{i+\frac{1}{2}} - D_{i+\frac{1}{2}}) \Delta^+ v_i^n + |D_{i+\frac{1}{2}} \Delta^+ v_i^n| + |C_{i+\frac{1}{2}} \Delta^+ v_i^n| \right| \quad (\text{by periodicity})$$

$$= \sum_i (1 - C_{i+\frac{1}{2}} - D_{i+\frac{1}{2}} + D_{i+\frac{1}{2}} + C_{i+\frac{1}{2}}) |\Delta^+ v_i^n|$$

(by the assumptions on the coefficients)

$$= \sum_i |v_{i+1}^n - v_i^n| = TV(\bar{v}^n)$$



example: first-order upwind FV method for linear advection is TVD

$$u_t + (au)_x = 0 \quad a > 0$$

$$\text{FOU: } \frac{v_j^{n+1} - v_j^n}{\Delta t} + \frac{a v_j^n - a v_{j-1}^n}{\Delta x} = 0$$

is also FV method with LF flux function  
so  $v_j^{n+1} = v_j^n - a \frac{\Delta t}{\Delta x} (v_j^n - v_{j-1}^n)$

$$C_{j-\frac{1}{2}} = a \frac{\Delta t}{\Delta x}$$

$$D_{j+\frac{1}{2}} = 0$$

TVD conditions:

$$C \geq 0 : \text{OK}$$

$$D \geq 0 : \text{OK}$$

$$C + D \leq 1 : \quad \text{or} \quad C \leq 1$$

$$a \frac{\Delta t}{\Delta x}$$

$\Rightarrow$  this explains why we see no spurious oscillations at shocks for this method

$$\boxed{\Delta t \leq \frac{\Delta x}{a}} \quad \text{OK!} \quad (\text{CFL condition!})$$

Can we have linear FV methods that are TVD?

def: linear FV scheme

A FV method is called linear if, when the scheme is applied to a linear PDE, all the coefficients  $C_i$  in

$$v_j^{n+1} = \sum_i C_i v_{j-i}^n$$

are constant (i.e., do not depend on  $v^n$ )

the following theorem can be proved:

Thm: Godunov's theorem

Linear TVD schemes are at most first-order accurate.

$\Rightarrow$  so if we want second-order accurate schemes, we need schemes with non-constant coefficients  
(see section 3.3.5)