

COPPER-MEDIATED CROSS-COUPLING REACTIONS

COPPER-MEDIATED CROSS-COUPLING REACTIONS

Edited by

GWILHERM EVANO

Laboratoire de Chimie Organique
Service de Chimie et PhysicoChimie Organiques
Université Libre de Bruxelles
Brussels, Belgium

NICOLAS BLANCHARD

Université de Strasbourg
Ecole Européenne de Chimie
Polymères et Matériaux
Laboratoire de Chimie Moléculaire associé au CNRS
Strasbourg, France

WILEY

Copyright © 2014 by John Wiley & Sons, Inc. All rights reserved

Published by John Wiley & Sons, Inc., Hoboken, New Jersey
Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permissions>.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Copper-mediated cross-coupling reactions / edited by Gwilherm Evano, Laboratoire de chimie organique, Service de chimie et physicochimie organiques, Université libre de Bruxelles, Brussels, Belgium; Nicolas Blanchard, Université de Strasbourg, Ecole européenne de chimie, polymères et Matériaux, Laboratoire de chimie moléculaire associé au CNRS, Strasbourg, France.

pages cm

Includes bibliographical references and index.

ISBN 978-1-118-06045-2 (hardback)

1. Copper catalysts. 2. Copper—Reactivity. 3. Chemical bonds. 4. Organic compounds—Synthesis. I. Evano, Gwilherm, editor of compilation. II. Blanchard, Nicolas, editor of compilation.

QD505.C685 2013

546'.652595—dc23

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

CONTENTS

FOREWORD	xvii
<i>Stephen L. Buchwald</i>	
PREFACE: COPPER CATALYSIS FROM A HISTORICAL PERSPECTIVE: A LEGACY FROM THE PAST	xix
<i>Gwilherm Evano and Nicolas Blanchard</i>	
CONTRIBUTORS	xxxiii
PART I FORMATION OF C–HETEROATOM BONDS	1
1 Modern Ullmann–Goldberg Chemistry: Arylation of N-Nucleophiles with Aryl Halides	3
<i>Yongwen Jiang and Dawei Ma</i>	
1.1 Introduction,	3
1.2 Arylation of Amines,	4
1.3 Arylation of Amides, Imides, and Carbamates,	15
1.4 Arylation of Conjugated N-Heterocycles,	24
1.5 Synthesis of Anilines by Coupling with Ammonia or Synthetic Equivalents,	32
1.6 Conclusion and Future Prospects,	37
References,	37

2 Ullmann Condensation Today: Arylation of Alcohols and Thiols with Aryl Halides **41**

Anis Tlili and Marc Taillefer

- 2.1 Introduction, 41
- 2.2 Formation of C–O Bonds via Copper-Catalyzed Cross-Coupling Reactions with Aryl Halides, 42
 - 2.2.1 Coupling of Aryl Halides with Hydroxide Sources: Synthesis of Phenols, 42
 - 2.2.2 Coupling of Aryl Halides with Aliphatic Alcohols: Synthesis of Alkyl Aryl Ethers, 49
 - 2.2.3 Coupling Reactions of Aryl Halides with Phenols: Synthesis of Biaryl Ethers, 54
- 2.3 Formation of C–S Bonds via Copper-Catalyzed Cross-Coupling Reactions with Aryl Halides, 67
 - 2.3.1 Coupling of Aryl Halides with Sulfur Sources: Synthesis of Thiophenols, 67
 - 2.3.2 Coupling of Aryl Halides with Aliphatic Thiols: Synthesis of Alkyl Aryl Thioethers (or Alkyl Aryl Sulfides), 70
 - 2.3.3 Coupling of Aryl Halides with Aryl Thiols: Synthesis of Diaryl Thioethers (or Diaryl Sulfides), 74
- 2.4 Conclusion, 84
References, 86

3 Copper-Catalyzed Formation of C–P Bonds with Aryl Halides **93**

Carole Alayrac and Annie-Claude Gaumont

- 3.1 Introduction, 93
- 3.2 Arylation of Phosphines, 94
 - 3.2.1 Arylation of Diarylphosphines, 94
 - 3.2.2 Arylation of Dialkylphosphines, 95
 - 3.2.3 Application to the Synthesis of Bulky Tertiary Phosphine Ligands, 97
- 3.3 Arylation of Phosphine Oxides and Phosphites, 98
 - 3.3.1 Arylation of Phosphine Oxides, 98
 - 3.3.2 Synthesis of Aryl Phosphonates by Arylation of Trialkylphosphites and H-Phosphonates, 101
 - 3.3.3 Arylation of Other Phosphorus Derivatives, 108
- 3.4 Conclusion, 110
References, 110

4 Alternative and Emerging Reagents for the Arylation of Heteronucleophiles **113**

Luc Neuville

- 4.1 Introduction, 113
- 4.2 Chan–Lam–Evans Coupling: Copper(II)-Promoted Oxidative Aryl Transfer from Arylboron Derivatives, 115
 - 4.2.1 Initial Discovery of the Chan–Lam–Evans Coupling, 115
 - 4.2.2 Developments in the Oxidative Copper-Mediated Arylation of Heteronucleophiles with Boronic Acids, 117
 - 4.2.3 Mechanism of the Coupling, 129
 - 4.2.4 Scope, 130
- 4.3 Copper-Promoted Aryl Transfer from Metallated Aryl Derivatives (Nonboron), 141
 - 4.3.1 Bismuth and Lead, 142
 - 4.3.2 Arylstannanes and Aryl-Alkoxysilanes, 146
 - 4.3.3 Aryliodonium Salts, 148
- 4.4 Copper-Catalyzed Arylation Reactions Involving Masked S- and N-Nucleophiles, 151
 - 4.4.1 C–S Bond Formation, 152
 - 4.4.2 C–N Bond Formation, 152
- 4.5 Copper-Catalyzed Direct Heterofunctionalization of Aromatic C–H Bonds, 160
 - 4.5.1 Functionalization of Acidic Aromatic C–H Bonds, 161
 - 4.5.2 Directed Aromatic C–H Bonds Functionalization, 166
 - 4.5.3 Intramolecular Copper-Catalyzed Heterofunctionalization of Aromatic C–H Bonds, 172
- 4.6 Conclusion and Future Prospects, 178
 - References, 178

5 Beyond Ullmann–Goldberg Chemistry: Vinylation, Alkynylation, and Allenylation of Heteronucleophiles **187**

Kévin Jouvin and Gwilherm Evano

- 5.1 Introduction, 187
- 5.2 Copper-Mediated Alkenylation of Heteronucleophiles: Among the Best Routes to Heteroatom-Substituted Alkenes, 189
 - 5.2.1 Synthesis of Enamides, Enamines, and N-Alkenyl-Heterocycles, 189
 - 5.2.2 Synthesis of Enol Ethers, 201
 - 5.2.3 Synthesis of Vinyl Sulfides and Other Sulfur-Substituted Alkenes, 210

5.2.4	Synthesis of Vinylphosphonates and Phosphine Oxides, 215	
5.3	Alkynylation of Heteronucleophiles: The Emergence of General Methods for the Synthesis of Heteroatom-Substituted Alkynes, 219	
5.3.1	Bringing Ynamides into the New Millennium: Copper-Mediated Alkynylation of Nitrogen Nucleophiles, 219	
5.3.2	Copper-Mediated Synthesis of Ynol Ethers: The First Development, 228	
5.3.3	Copper-Mediated Preparation of Alkynyl Chalcogenides, 229	
5.3.4	Synthesis of Phosphorus-Substituted Alkynes: The Development of Mild Protocols Based on Copper Catalysis, 230	
5.4	Allenylation of Heteronucleophiles: New Tools for the Synthesis of Allenamides, 232	
5.5	Conclusion and Future Prospects, 233 References, 234	
6	Aromatic/Vinyllic Finkelstein Reaction	239
	<i>Alicia Casitas and Xavi Ribas</i>	
6.1	Introduction, 239	
6.2	Copper-Mediated Halogen Exchange Reactions in Aryl Halides, 241	
6.2.1	From Early Discoveries to Most Recent Reports, 241	
6.2.2	Copper-Mediated Fluorination of Haloarenes, 245	
6.2.3	Mechanistic Investigations in Aryl–Cu(III) Model Systems, 246	
6.3	Most Recent Developments and Overview, 247 References, 249	
7	Insights into the Mechanism of Modern Ullmann–Goldberg Coupling Reactions	253
	<i>Alicia Casitas and Xavi Ribas</i>	
7.1	General View and Key Mechanistic Aspects, 253	
7.2	Oxidation State of Copper Catalysts, 254	
7.3	Identity of the Active Copper(I) Complex, 255	
7.4	Activation Mode of Aryl Halides by Copper Complexes, 261	
7.4.1	Oxidative Addition/Reductive Elimination Pathway, 262	
7.4.2	SET Mechanism, 270	
7.4.3	AT Mechanism, 272	

7.4.4	Mechanism Involving π -Complexation of Copper(I) to Aryl Halides, 273	
7.4.5	Mechanism Involving σ -Bond Metathesis, 275	
7.5	Overview, Conclusions, and Future Prospects, 275 References, 277	
PART II FORMATION OF C–C BONDS		281
8	Modern Copper-Catalyzed Hurtley Reaction: Efficient C-Arylation of CH-Acid Derivatives	283
	<i>Irina P. Beletskaya and Alexey Yu. Fedorov</i>	
8.1	Introduction, 283	
8.2	Classical Hurtley Reaction, 285	
8.3	Ligation Effect in Copper-Catalyzed Reactions of Aryl Halides with Carbanions, 286	
8.4	Cascade Reactions Proceeding via a Hurtley Arylation Reaction, 293	
8.5	Mechanism of the Copper-Catalyzed C-Arylation Reactions, 303	
8.6	Concluding Remarks, 308 References, 308	
9	Copper-Catalyzed Cyanations of Aryl Halides and Related Compounds	313
	<i>Thomas Schareina and Matthias Beller</i>	
9.1	Introduction, 313	
9.2	Modifications and Updates of Classical Cyanation Reactions (Rosenmund–von Braun, Sandmeyer), 315	
9.3	Copper-Catalyzed Cyanations of Aryl Halides, 316	
9.4	Copper-Mediated Oxidative Cyanations, 324	
9.5	Conclusion, 331 References, 331	
10	Copper-Mediated Aryl–Aryl Bond Formation Leading to Biaryls: A Century after the Ullmann Breakthrough	335
	<i>Yoshihiko Yamamoto</i>	
10.1	Introduction, 335	
10.2	Biaryl Synthesis by Coupling of Aryl Halides and Diazonium Salts, 336	
10.2.1	Ullmann Couplings of Aryl Halides, 336	
10.2.2	Coupling of Organocopper Reagents Generated from Aryl Halides, 342	
10.2.3	Coupling of Aryl Diazonium Compounds, 346	

- 10.3 Biaryl Synthesis by Coupling of Aryltin, Boron, and Silanes, 347
 - 10.3.1 Coupling of Organostannanes, 347
 - 10.3.2 Coupling of Organoboranes, 351
 - 10.3.3 Coupling of Organosilanes, 356
- 10.4 Biaryl Synthesis by Arylation Involving Arene C–H or C–C Bond Fission, 357
 - 10.4.1 Direct C–H Arylation of Arenes with Prefunctionalized Arylating Reagents, 357
 - 10.4.2 Direct C–H Arylation with Nonfunctionalized Arylating Agents, 368
 - 10.4.3 Decarboxylative Coupling of Benzoic Acid Derivatives, 373
- 10.5 Biaryl Synthesis by Oxidative Coupling of 2-Naphthols, 376
- 10.6 Conclusions and Outlook, 387
 - References, 388

11 Copper-Catalyzed Alkynylation, Alkenylation, and Allylation Reactions of Aryl Derivatives

401*Ren-Jie Song and Jin-Heng Li*

- 11.1 Introduction, 401
- 11.2 Copper-Catalyzed Alkynylation of Aryl Derivatives, 402
 - 11.2.1 Synthesis of Aryl–Ynes by Stille-Type Cross-Coupling Reaction, 402
 - 11.2.2 Synthesis of Aryl–Ynes by Sonogashira-Type Cross-Coupling Reaction, 405
 - 11.2.3 Synthesis of Aryl–Ynes by Oxidative Cross Coupling of Alkynes and Arylboronic Acids, 422
 - 11.2.4 Synthesis of Aryl–Ynes by Decarboxylative Cross Coupling from Propiolic Acids, 424
 - 11.2.5 Synthesis of Aryl–Ynes by Direct C–H Functionalization of Arenes, 427
- 11.3 Copper-Catalyzed Alkenylation of Aryl Derivatives, 432
 - 11.3.1 Synthesis of Aryl–Enes by Stille-Type Cross Coupling, 432
 - 11.3.2 Synthesis of Aryl–Enes by Suzuki–Miyaura-Type Cross Coupling, 435
 - 11.3.3 Synthesis of Aryl–Enes by Heck-Type Cross Coupling, 437
 - 11.3.4 Synthesis of Aryl–Enes by Direct C–H Functionalization of Arenes, 441
 - 11.3.5 Synthesis of Aryl–Enes by Decarboxylative Cross Coupling, 444

11.4	Copper-Catalyzed Strategies for the Formation of Allyl–Aryl Bonds, 445	
11.5	Conclusion and Outlook, 450 References, 450	
12	Copper-Catalyzed Alkynylation and Alkenylation Reactions of Alkynyl Derivatives: New Access to Diynes and Enynes	455
	<i>Ruimao Hua</i>	
12.1	Introduction, 455	
12.2	Copper-Catalyzed Synthesis of Symmetrical and Unsymmetrical 1,3-Diynes, 456	
12.3	Copper-Catalyzed Synthesis of 1,4-Diynes, 468	
12.4	Synthesis of 1,3-Enynes by Direct Reaction of Vinyl Halides with Alkynes, 468	
12.5	Synthesis of 1,3-Enynes by Stille-Type Cross-Coupling Reaction, 474	
12.6	Synthesis of 1,3-Enynes by the Suzuki–Miyaura-Type Cross-Coupling Reaction, 476	
12.7	Synthesis of 1,4-Enynes by Allylation Reaction of Terminal Alkynes, 478	
12.8	Conclusion, 480 References, 480	
13	Copper-Mediated Alkenylation Reaction of Alkenyl Derivatives: A Straightforward Elaboration of 1,3-Dienes	485
	<i>Hao Li, Songbai Liu, and Lanny S. Liebeskind</i>	
13.1	Introduction, 485	
13.2	Symmetrical 1,3-Dienes by Homocoupling Reaction of Vinyl Derivatives, 486	
13.2.1	Coupling of Vinylboranes, 486	
13.2.2	Coupling of Vinylsilanes, 487	
13.2.3	Coupling of Vinylstannanes, 489	
13.2.4	Coupling of Vinyl Halides, 493	
13.2.5	Miscellaneous, 496	
13.3	Unsymmetrical 1,3-Dienes by Cross-Coupling Reactions, 496	
13.3.1	Synthesis of 1,3-Dienes by Stille-Type Coupling, 497	
13.3.2	Synthesis of 1,3-Dienes by Suzuki-Type Coupling, 505	
13.3.3	Synthesis of 1,3-Dienes by Heck-Type Coupling, 508	
13.4	Conclusions, 510 References, 511	

14 Emerging Areas in Copper-Mediated Trifluoromethylations of Aryl Derivatives: Catalytic and Oxidative Cross-Coupling Processes **515**

Kévin Jouvin, Céline Guissart, Cédric Theunissen, and Gwilherm Evano

- 14.1 Introduction, 515
- 14.2 Copper-Catalyzed Trifluoromethylation of Aryl Halides: A Long-Lasting Quest Finally Reached, 517
- 14.3 Copper-Mediated Oxidative Trifluoromethylation Reactions, 523
- 14.4 Conclusion and Future Prospects, 528
References, 528

PART III APPLICATIONS OF COPPER-CATALYZED CROSS-COUPPLING REACTIONS: HETEROCYCLES, NATURAL PRODUCTS, PROCESS, AND SUSTAINABLE CHEMISTRY **531**

15 Copper-Mediated Cyclization Reactions: New Entries to Heterocycles **533**

Daoshan Yang and Hua Fu

- 15.1 Introduction, 533
- 15.2 Cyclization by C–N Bond Formation, 534
 - 15.2.1 Copper-Catalyzed Synthesis of Pyrroloindole Derivatives, 536
 - 15.2.2 Copper-Catalyzed Synthesis of Benzimidazole Derivatives, 536
 - 15.2.3 Copper-Catalyzed Synthesis of 1,3-Dihydrobenzimidazol-2-One Derivatives, 544
 - 15.2.4 Copper-Catalyzed Synthesis of Quinazolines, 547
 - 15.2.5 Copper-Catalyzed Synthesis of Quinazolinone Derivatives, 551
 - 15.2.6 Copper-Catalyzed Synthesis of 1,2,4-Benzothiadiazine 1,1-Dioxides, 552
 - 15.2.7 Copper-Catalyzed Synthesis of 1-Aryl-2-Thioxo-2,3-Dihydroquinazolin-4(1H)-Ones, 553
 - 15.2.8 Copper-Catalyzed Synthesis of Indole Derivatives, 554
 - 15.2.9 Copper-Catalyzed Formation of Cyclic Enamides: Efficient Routes to Unsaturated Lactams, Alkylideneazetidines, and Dihydropyrroles, 556
 - 15.2.10 Copper-Catalyzed Synthesis of 1,4-Benzoxazines, 558
- 15.3 Cyclization by C–O Bond Formation, 560
 - 15.3.1 Copper-Catalyzed Synthesis of Benzoxazole Derivatives, 560

- 15.3.2 Copper-Catalyzed Synthesis of Cyclic Enol and Aryl Ethers, 562
- 15.3.3 Copper-Catalyzed Synthesis of Enol Lactones and Benzopyranones, 565
- 15.4 Cyclization by C–C Bond Formation, 567
 - 15.4.1 Copper-Catalyzed Synthesis of Indole Derivatives, 567
 - 15.4.2 Copper-Catalyzed Synthesis of 3-Acyloxindoles, 569
 - 15.4.3 Copper-Catalyzed Synthesis of Isoquinoline and Isoquinolinone Derivatives, 571
 - 15.4.4 Heterocyclic Synthesis Based on a Copper-Catalyzed Sonogashira-Type Cross Coupling, 574
- 15.5 Copper-Catalyzed Double Cross-Coupling Reactions for the Assembly of Heterocycles, 576
 - 15.5.1 Synthesis of Benzoxazoles and Oxazoles by Double Copper-Catalyzed Cross Coupling, 576
 - 15.5.2 Synthesis of Benzimidazoles by Double Copper-Catalyzed Cross Coupling, 578
 - 15.5.3 Synthesis of Benzofurans by Double Copper-Catalyzed Cross Coupling, 579
 - 15.5.4 Synthesis of Benzimidazo[1,2-*b*]isoquinolin-11-one Derivatives by Double Copper-Catalyzed Cross Coupling, 580
 - 15.5.5 Synthesis of Pyrrole and Thiophene Derivatives by Double Copper-Catalyzed Cross Coupling, 581
- 15.6 Conclusion and Future Prospects, 583
- References, 584

16 Application of Copper-Mediated C–N Bond Formation in Complex Molecules Synthesis

589*Jihoon Lee and James S. Panek*

- 16.1 Introduction, 589
- 16.2 Aryl Amination in Complex Molecule Synthesis, 590
- 16.3 Aryl Amidation in Complex Molecule Synthesis, 595
- 16.4 Arylation of N-Heterocycles in Complex Molecule Synthesis, 601
- 16.5 Vinyl Amidation in Complex Molecule Synthesis, 606
- 16.6 Alkyne Amidation in Complex Molecule Synthesis, 620
- 16.7 Intramolecular C–N Bond Formation in Natural Product Synthesis, 622
 - 16.7.1 Copper-Mediated Synthesis of Small- to Medium-Sized N-Heterocycles in Total Synthesis, 622
 - 16.7.2 Copper-Mediated Macrocyclization by C–N Bond Formation in Natural Product Synthesis, 632
- 16.8 Summary and Outlook, 637
- References, 638

17	Natural Products and C–O/C–S Bond-Forming Reactions: Copper Showed the Way	643
	<i>Doron Pappo</i>	
17.1	Introduction, 643	
17.2	Total Synthesis of Naturally Occurring Diaryl Ethers by Arylation of Phenols, 644	
17.3	Intramolecular Diaryl Ether Bond-Forming Reactions, 659	
17.3.1	Macrocyclization, 659	
17.3.2	Small- to Medium-Sized Oxygenated Heterocycles, 663	
17.4	Arylation of Alcohols, 666	
17.5	Vinylation of Alcohols, 673	
17.6	Copper-Mediated C–S Bond Formation in Natural Product Synthesis, 675	
17.7	Conclusion and Future Prospects, 677	
	References, 678	
18	Copper-Catalyzed C–C Bond Formation in Natural Product Synthesis: Elegant and Efficient Solutions to a Key Bond Disconnection	683
	<i>Morgan Donnard and Nicolas Blanchard</i>	
18.1	Introduction, 683	
18.2	Natural Biaryls by Copper-Catalyzed Cross Coupling, 684	
18.2.1	Classical Ullmann Cross-Coupling Reaction, 685	
18.2.2	Oxidative Biaryl Coupling of Phenols and Naphthols, 687	
18.3	Copper-Catalyzed 1,3-Enyne Formation, 691	
18.4	Copper-Mediated Synthesis of Dienes, Trienes, and Extended Polyenes, 694	
18.4.1	Copper-Mediated Synthesis of 1,3-Dienes, 694	
18.4.2	Copper-Mediated Synthesis of 1,3,5-Trienes, 701	
18.4.3	Copper-Mediated Synthesis of Extended Polyenes, 704	
18.4.4	Copper Salts as Essential Cocatalysts of Palladium-Catalyzed Stille Cross-Coupling Reactions, 704	
18.5	Copper-Catalyzed Synthesis of 1,N-Polyyne Natural Products, 711	
18.5.1	Cadiot–Chodkiewicz Strategies, 712	
18.5.2	Eglinton Strategies, 714	
18.5.3	Hay Strategies, 716	
18.5.4	Copper-Catalyzed Synthesis of Skipped Dienes, 718	
18.6	Conclusions and Future Prospects, 718	
	References, 719	

19 Process Chemistry and Copper Catalysis 725*Klaus Kunz and Norbert Lui*

- 19.1 Introduction and Scope, 725
- 19.2 Copper versus Palladium, 727
- 19.3 Applications, 727
 - 19.3.1 Materials Sciences, 727
 - 19.3.2 Pharmaceutical Chemistry, 731
 - 19.3.3 Crop Science Chemistry, 737
- 19.4 Conclusion, 739
- References, 740

20 Reusable Catalysts for Copper-Mediated Cross-Coupling Reactions under Heterogeneous Conditions 745*Zhiyong Wang, Changfeng Wan, and Ye Wang*

- 20.1 Introduction, 745
- 20.2 Copper Nanoparticle-Catalyzed Cross-Coupling Reactions, 746
 - 20.2.1 C–C Bond Formation, 746
 - 20.2.2 C–N Bond Formation, 750
 - 20.2.3 C–S Bond Formation, 754
 - 20.2.4 C–O Bond Formation, 757
 - 20.2.5 C–Se Bond Formation, 762
- 20.3 Supported Copper-Catalyzed Cross-Coupling Reaction, 766
 - 20.3.1 C–C Bond Formation, 766
 - 20.3.2 C–X Bond Formation, 769
- 20.4 Conclusion, 780
- References, 780

INDEX 785

FOREWORD

The development of efficient methods for use in organic synthesis, particularly applicable to polyfunctionalized molecules and heterocycles, remains one of the top challenges in organic chemistry. Oftentimes older reactions are rediscovered and improved, sometimes substantially, in order to solve modern problems. A prime example of this is copper-catalyzed methodology. This topic, studied at the beginning of the twentieth century by Ullmann, Goldberg, and others has experienced a renaissance in the past two decades. A significant reason for this is the development of new and improved ligands to facilitate the transformations of interest.

While the most obvious difference between catalysts based on palladium and copper is the lower cost of the latter, this is only a small part of what makes copper catalysts so attractive. For most chemists in academia or discovery chemistry, the cost is not a major issue. In addition, one must take in to account many factors, such as volumetric productivity (reactor time, labor cost), substrate cost, yield, and substrate scope, which affect the total cost when comparing any two processes. Among the major advantages of using copper catalysts (in addition to cost) are that they tend to show excellent functional group compatibility and, in fact, are often at their best in promoting reactions of substrates that contain coordinating functional groups. A prime example, as delineated in Evano and Blanchard's preface to this book, is seen in Ullmann's seminal work: he was able to accomplish the coupling of aniline with 2-chlorobenzoic acid well before the same transformation could be carried out with palladium catalysts. In fact, the chemistry based on these two metals is really quite complementary. In support of this notion, today these are used interchangeably for many synthetic operations depending on the nature of the substrate combination to be examined.

The chapters in the book cover a range of synthetically and mechanistically important topics. Many aspects of carbon-heteroatom and carbon-carbon,

bond-forming reactions are described. These have become an everyday part of the practicing synthetic organic chemist's arsenal. In addition, topics such as the formation of trifluoromethylated organic molecules and the application of copper-catalyzed methods to the total synthesis of natural products and pharmaceuticals are also covered. Taken together, the chapters in this book provide an up-to-date overview of the state of the field, providing the reader with the knowledge of both what has been accomplished and what new areas remain for further investigation.

Massachusetts Institute of Technology

STEPHEN L. BUCHWALD

PREFACE

Copper Catalysis from a Historical Perspective: A Legacy from the Past

GWILHERM EVANO AND NICOLAS BLANCHARD

THE HISTORICAL AND REMARKABLE FIRST DISCOVERIES BY FRITZ ULLMANN AND IRMA GOLDBERG

It all started in 1901 in Geneva when Fritz Ullmann, born in 1875 in Fürth, where his father owned a fabric... of metallic powders,^[1] reported, in a remarkable publication that we highly recommend, that “Erhitzt man *o*-Bromnitrobenzol mit fein vertheiltem Kupferpulver, so bemerkt man, dass letzteres seinen Glanz verliert und in eine matte graue Masse verwandelt wird. Bei Aufarbeitung des Reactionsproductes zeigte sich nun, dass das Kupfer zum grössten Theil in Cuprobromid und das Bromnitrobenzol in eine bromfreie Substanz verwandelt worden ist, welche sich bei näherer Untersuchung identisch, mit der von Tauber auf andere Weise dargestellten 2,2'-Dinitrobiphenyl, erwies.” (If one heats *o*-bromonitrobenzene with finely divided copper powder, so one recognizes that the last one is losing its shine and turns into a matt grey mass. After purification of the reaction products, it appears that copper has turned into copper bromide and that the bromonitrobenzene has turned into a bromine-free substance, which, on a closer look, turns out to be identical with the 2,2-dinitrobiphenyl synthesized in a different way by Tauber) (Fig. P1).^{*,[2]} The reaction can also be performed from aryl iodides and, which is just astonishing, even from aryl chlorides (Scheme P.1). Two molecules of aryl

* The editors are truly indebted to Dr. Théophile Tschamber from the University of Mulhouse (France) for his expertise in translating the numerous original German research papers and patents that are cited in the Preface.





Figure P.1 Fritz Ullmann (1875–1939) and an extract from his original publication.


halides can be coupled together in the presence of stoichiometric amounts of metallic copper: The Ullmann reaction had just been discovered.

Obviously fascinated with the power of the metallic copper, Ullmann next reported in 1903 an equally fascinating reaction in a publication entitled “Über eine neue Bildungsweise von Diphenylaminderivaten” (*On a New Method for the Formation of Diphenylamine Derivatives*). By mixing aniline (or other aniline derivatives) and *ortho*-chlorobenzoic acid in the presence of metallic copper at reflux, the aniline was smoothly arylated to the corresponding diarylamine with a remarkable efficiency (Scheme P.2).^[3] Two years after his initial discovery of the biaryl synthesis, Ullmann continued to set the foundations of copper-mediated cross-coupling reactions and the arylation of amines that would later on be named after him was invented. Also remarkable is that the aryl halide substrate chosen for this study, *ortho*-chlorobenzoic acid, already possessed a non-innocent chelating group: the *ortho*-effect, which still has a deep impact in copper catalysis today and which was crucial to the success of many copper-mediated transformations including in Nicolaou’s synthesis of vancomycin (see further), was already touched upon in 1903.


Following these two reports, Ullmann’s assistant in Geneva, Irma Goldberg, reported 3 years later that a related reaction could be performed using only catalytic amounts of copper. By reacting anthranilic acid and bromobenzene in the presence of potassium carbonate and catalytic amounts of “Naturkupfer C” in refluxing nitrobenzene for 3 hours, the reaction proceeded with a remarkable efficiency, the corresponding condensation product being isolated in 99% yield (Scheme P.3).^[4] The arylation could also be efficiently performed with *para*-nitro-bromobenzene and, even better, the arylation of amides (benzamide in this case), known nowadays as the Goldberg condensation reaction, could also be catalyzed by traces of metallic copper. These three remarkable




Scheme P.1 1901: the discovery of the Ullmann reaction.



Scheme P.2 1903: the discovery of the Ullmann condensation reaction for the synthesis of diarylamines.



Scheme P.3 1906: the discovery of the copper-catalyzed arylation of anilines and amides by Irma Goldberg.



Scheme P.4 1905: the discovery of the copper-catalyzed arylation of phenols by Fritz Ullmann.

reactions published between 1901 and 1906 mark the beginning of cross-coupling reactions and clearly paved the way for all recent developments.

Last but not least, Ullmann also studied the cross coupling involving phenols and reported in 1905 that “if you try to react potassium phenoxide with bromobenzene, the yield of the biphenyl ether is 0.9%. If you however add small quantities of copper to the reaction mixture, the yield goes to 90%” (Scheme P.4).^[5] It is therefore possible to catalyze the cross coupling between phenols and aryl bromides with small amounts of copper, which obviously have a dramatic effect on the reaction rate and yields.

FURTHER HISTORICAL DEVELOPMENTS: THE CYANATION OF ARYL HALIDES BY ROSENMUND AND VON BRAUN AND THE ARYLATION OF DIKETONES AND MALONATES BY HURTLEY

In a series of just four publications, Fritz Ullmann and Irma Goldberg, who eventually got married in 1910, clearly set the foundations of modern copper catalysis. This pioneering work was followed by other key reports in the next 30 years or so, which further demonstrated the potential of copper-mediated cross-coupling reactions.

The next major development in this field was stimulated by the interest in the copper-mediated cyanation of aryl halides, which appeared closely after the seminal publications of Ullmann and Goldberg. Although this transformation is known as the “Rosenmund–von Braun” reaction, many academic and industrial research groups contributed to the development of this very useful reaction.^[6–8] Indeed, the first report of such a reaction appeared in the patent literature in 1913 by the Meister, Lucius & Brünig Company who found that chloro- and bromoanthraquinones reacted with copper(I) cyanide in pyridine at 150°C to give the corresponding nitriles (Fig. P.2 and Scheme P.5).^[8]

Karl W. Rosenmund and his PhD student Erich Struck described next in 1919 that potassium cyanide in combination with copper(I) cyanide converted aryl halides to the corresponding benzoic acids in water at 200°C under pressure. The intermediate nitriles were not isolated, but the central role of the copper(I) species was put forward (Scheme P.6, Eq. 1).^[9a] Rosenmund disclosed a year later^[9b] that stoichiometric copper(I) thiocyanate in pyridine

KAISERLICHES



PATENTAMT.

PATENTSCHRIFT

— № 271790 —

KLASSE 12 o. GRUPPE 10.

AUSGEBEN DEN 18. MÄRZ 1913.

FARBWERKE FORM. MEISTER LUCIUS & BRÜNING IN HÖCHST A.M.

Verfahren zur Darstellung von Anthrachinon- α -nitrilen.

Patentiert im Deutschen Reich vom 18. Februar 1913 ab.

Es wurde gefunden, daß Anthrachinon- α -nitrile erhalten werden, wenn man α -Halogenanthrachinone mit Kupfercyanür in Gegenwart von indifferenten Mitteln erhitzt, während eine ähnliche Umsetzung mittels Cyankaliums nicht gelingt.

Dies war nicht vorauszusehen, da in verschiedenen organischen Halogenverbindungen mit leicht beweglichem Halogen sich dieses überhaupt nicht durch eine Cyangruppe ersetzen läßt (vgl. Windisch, Dissertation, Berlin 1887; V. v. Richter, Ber. IV, 1871, Seite 462).

H. Dienel hat bereits das α -Cyananthrachinon sowohl durch Oxydation von α -Cyananthracen wie durch trockne Destillation von anthrachinonsulfosaurem Kalium mit Cyankalium erhalten (Ber. XXXIX, 1906, Seite 932). Ullmann und van der Schalk haben dann nachgewiesen, daß man beim Nacharbeiten des letzteren Verfahrens an α -Cyananthrachinon im günstigsten Falle 1 Prozent der Theorie erhält (Ann. 388, 205), und gezeigt, daß α -Cyananthrachinon im allgemeinen aus den α -Aminoanthrachinonen über die Diazoniumverbindungen mittels der Sandmeyerischen Reaktion gewonnen werden (Ann. 388, 203); Patent 243788.

Den bisher bekannten Verfahren gegenüber bedeutet das vorliegende schon deshalb einen großen technischen Fortschritt, weil es verschiedene neue, auf anderem Wege nicht erhältliche Derivate des Anthrachinon- α -nitrils darzustellen gestattet.



Beispiele.

1. Anthrachinon- α -nitril: 24 Teile 1-Chloranthrachinon werden mit 9 Teilen Kupfercyanür und 100 Teilen Pyridin während einiger Stunden auf 150° erhitzt. Das Reaktionsprodukt wird mit Ammoniak verdünnt und das abgeschiedene rohe Nitril abgesaugt. Zur Reinigung wird es z. B. aus Chlorbenzol umkristallisiert und so in Form brauner Kristalle erhalten, die bei 247° schmelzen und eine reine blaue Küpe geben.





2. 2-Amino-3-bromanthrachinon- α -nitril: 20 Teile 1-3-Dibrom-2-aminoanthrachinon werden mit 5 Teilen Kupfercyanür und 100 Teilen Pyridin etwa 6 Stunden am Rückflußkühler erhitzt. Nach dem Erkalten wird das grünlichgelbe, schön kristallisierte Nitril abgesaugt und gewaschen. Aus Chlorbenzol kristallisiert es in gelblichen Nadeln, die bei 297 bis 300° schmelzen. Die Lösung in konzentrierter Schwefelsäure ist rotorange und wird beim Erwärmen rot. Gießt man die schwefelsaure Lösung des Nitrils in Wasser, so fällt die 2-Amino-3-bromanthrachinon- α -carbonsäure in gelben Flocken aus.

3. 2-Oxy-3-bromanthrachinon- α -nitril: 20 Teile 1-3-Dibrom-2-oxyanthrachinon werden mit 5 Teilen Kupfercyanür und 200 Teilen Pyridin 6 Stunden am Rückflußkühler erhitzt. Nach dem Erkalten wird das rote, in Pyridin schwer lösliche Pyridinsalz des 2-Oxy-3-bromanthrachinon- α -nitrils abgesaugt. Aus der Lö-


Figure P.2 The 1913 Meister, Lucius & Brüning Company patent.



Scheme P.5 1913: The first copper(I)-mediated cyanations of haloanthraquinones by the Meister, Lucius & Brünig Company.



Scheme P.6 1919–1923: Karl W. Rosenmund and Henri de Diesbach early reports on the cyanation of aryl halides.




Scheme P.7 1927–1931: Alfred Pongratz and Julius von Braun cyanation of aryl halides.

converted aryl bromides to intermediate aryl thiocyanates, which spontaneously gave the corresponding nitriles at 180°C in the presence of the copper species,^[6] the nitriles being characterized as carboxylic acids after aqueous acidic treatment (Scheme P6, Eq. 2). Shortly after, Henri de Diesbach reported in two back-to-back articles in 1923 that dibromoxylene derivatives could be converted to the corresponding dinitriles in excellent yield using an excess of copper(I) cyanide in pyridine at 200°C (Scheme P6, Eq. 3),^[10] and even aryl chlorides could be transformed to the corresponding nitriles, as seen in Scheme P6, Eq. 4.^[11]

In 1927, Alfred Pongratz described that copper(I) cyanide in refluxing quinoline was efficient for the cyanation of perylene-3,9-dibromide (Scheme P.7, Eq. 1) and that the corresponding chloride could be converted in the absence of solvent at 300°C, although no yield was given in the latter case.^[12] Finally, during investigation on the chemistry of fluoranthene derivatives, Julius von Braun, who gave his name to this cyanation reaction, demonstrated that 4-bromo-fluoranthene could be transformed to the corresponding nitrile in 80% yield when reacted with copper(I) cyanide at 260°C (Scheme P.7, Eq. 2).^[13]

The pioneering investigations of Ullmann, Goldberg and Rosenmund^[8b] were a source of inspiration for William R. H. Hurtley, who recognized their contributions in his seminal 1929 report, stating that “under the catalytic influence of copper, the halogen in *o*-bromo-benzoic acid is much more reactive than is commonly realized.”^[14] Hurtley showed that copper bronze or copper acetate can promote the C-arylation of some families of CH-acids in reaction with *o*-bromobenzoic acid in the presence of sodium ethoxide as a base (Scheme P.8). Since then, this transformation, commonly known as the “Hurtley reaction”, has triggered mechanistic investigations as well as very elegant applications in organic synthesis. Mild conditions are now available thanks to




Scheme P.8 1929: the discovery of the copper-mediated arylation of CH-acid derivatives by William R. H. Hurtley.

new classes of ligands that even allowed the development of an asymmetric version of the Hurtley reaction.

FROM THE HISTORICAL DISCOVERIES TO THE DEVELOPMENT OF MODERN COPPER-MEDIATED CROSS-COUPLING REACTIONS

While remarkable, these reactions, however, have found only a limited number of applications, especially when compared to their impressive potential. Indeed, drawbacks associated with the use of harsh reaction conditions (strong bases, stoichiometric amounts of copper in some cases, and high reaction temperatures) have clearly hampered the development of these copper-mediated cross-coupling reactions. From the first discoveries in the early 1900s up to the beginning of the twenty-first century, no real breakthrough was made to significantly improve the original systems and the strategy was to tailor the substrate, not the catalyst, to obtain systems that would perform under milder conditions. Nicolaou's remarkable total synthesis of vancomycin is one of the most representative example of such a strategy. Capitalizing on the well-known *ortho*-effect in copper-mediated transformations, a mild copper-mediated arylation reaction was designed based on the ingenious incorporation of a triazene unit in the starting material strategically placed *ortho* to the bromine: this internal helper auxiliary served both as a potential "electron sink" and to coordinate the intermediate copper species. The triazene indeed considerably helped the reaction, which enabled the formation of the C–O–D and D–O–E ring systems under remarkably mild conditions since the reaction could be performed in refluxing acetonitrile (Scheme P.9).^[15]

Besides these strategies based on substrate modifications, the inherent shortcomings of the original copper-mediated cross-coupling reactions were still not overcome at the end of the twentieth century and palladium was in most cases preferred to copper. Despite remarkable work done with palladium catalysis for the construction of an impressive and ever-increasing number of C–C and C–heteroatom bonds, limitations still exist and the high cost of palladium and ligands has forced chemists to consider alternative metals. This evolution eventually led to the renaissance of copper catalysis, which has been extremely revisited over the past decade. Following extensive and remarkable studies of the Ullmann condensation by Payne,^[16] who showed that the active



Scheme P.9 The Ullmann condensation reaction under milder conditions: redesigning the substrate to improve the efficiency (Nicolaou's total synthesis of vancomycin).

catalytic species are soluble cuprous ions and pioneering work by Bryant,^[17] Capdevielle,^[18] and Goodbrand,^[19] who investigated the idea of a “ligated copper catalysis”, the key breakthrough was the introduction of chelating ligands for copper.

Remarkable and pioneering studies in this area from, among others, the Buchwald, Ma, and Taillefer groups had a strong impact in the renaissance of copper catalysis, and the Ullmann, Goldberg, Rosenmund–von Braun, Hurtley, and other related copper-mediated processes can now be conducted under much milder conditions together with dramatically enhanced yield. Recent advances in the arylation of N-, O-, and S- as well as P-nucleophiles with aryl halides will be described in Chapters 1, 2, and 3, respectively.

In addition to the development of improved protocols for the arylation of various heteronucleophiles, the introduction of chelating ligands also considerably expanded the scope of the original procedures and allowed for the use of new reaction partners for the introduction of vinyl, alkyne, and allene functional groups. These copper-mediated syntheses of heterosubstituted alkenes, alkynes, and allenes will be overviewed in Chapter 5. Copper-catalyzed aromatic versions of the Finkelstein reaction have also been recently reported: they proceed with remarkable efficiencies and will be covered in Chapter 6.

These recent developments also had a strong impact on the development of new and improved protocols for the formation of C–C bonds, and a variety of robust, mild, and reliable procedures have been designed with the assistance of copper complexes.

The classical Hurtley-type transformation as well as the tremendous progresses that have been reported recently in this central C-arylation reaction of CH-acid derivatives will be presented in Chapter 8, with an emphasis on the importance of the different types of ligands, mechanisms, and synthetic applications. The copper-catalyzed synthesis of nitriles from aryl halides is then covered in Chapter 9, including discussions on the different catalytic systems and sources of cyanide that have been greatly improved in the past two decades.

A central reaction in the realm of copper-catalyzed and copper-mediated transformations is of course the aryl–aryl coupling that has continuously evolved into an efficient and practical tool with multiple applications. Chapter 10 is devoted to this vibrant field of research that has recently witnessed a shift of paradigm with even more direct reactions such as the copper-catalyzed direct C–H arylations. The next three chapters, Chapters 11, 12, and 13, have captured relevant advances in some of the most practical and efficient copper-catalyzed transformations that have clearly modified our conception of carbon–carbon bond disconnections: the alkenylation and alkynylation of aryl, alkynyl, and vinyl derivatives.

To close this section dedicated to the formation of carbon–carbon bonds, the central role of copper catalysts in methods aiming at the introduction of the trifluoromethyl moiety will be exposed. Indeed, the occurrence of the trifluoromethyl motif in pharmaceutical and agrochemical active ingredients

is well recognized and has triggered the development of new methodologies for its selective introduction. The copper-catalyzed trifluoromethylation reaction and copper-mediated oxidative processes are efficient strategies in this context and are developed in detail in Chapter 14.

Besides substrate modifications and the design of more efficient catalytic systems, the drawbacks associated with the harsh conditions of the classical Ullmann–Goldberg and related transformations also motivated the search for alternative reagents to aryl halides that would enable the arylation of heteronucleophiles under milder reaction conditions. In 1998, independent back-to-back publications by Chan,^[20] Evans,^[21] and Lam^[22] revolutionized the arylation of heteronucleophiles as they reported generally applicable protocols for the copper-mediated oxidative cross coupling with arylboronic acids at room temperature. This reaction is now widely applied and other reagents such as organobismuth, -lead, -stannanes, -siloxanes, or hypervalent iodonium salts have also been used. They are sometimes more reactive than aryl halides, allow overcoming some limitations met with these reagents, or are just more readily available in some cases. Advances in copper-mediated cross coupling using these reagents are covered in Chapter 4 as well as the heteroarylation involving masked nucleophiles and the direct introduction of a heteronucleophile from a C–H bond. The use of these alternative reagents for the alkenylation/alkynylation of heteronucleophiles will be overviewed in Chapter 5.

If the modern copper-catalyzed Ullmann and Goldberg cross-coupling reactions have recently evolved as reliable and efficient methods for the construction of products relevant to both industry and academic settings, their mechanisms have been studied only recently. Mechanistic insights into these reactions, which will be crucial for the development of even more efficient catalytic systems, will be described in Chapter 7.

MODERN COPPER-MEDIATED CROSS-COUPLING REACTIONS AND THEIR IMPACT IN ORGANIC CHEMISTRY

We have all heard that we have arrived at a point where it is possible to make almost any conceivable chemical structure by a rational approach, using the large toolbox of synthetic methods available today and that there was basically nothing left to do in organic synthesis. The impact that had the renaissance of copper catalysis over the past 10 years is one of the most striking examples that this is just as stupid as it seems. Many long-standing synthetic problems have been solved using copper-mediated cross-coupling technologies, which had an important impact on the way chemists design their synthetic routes. These transformations have found remarkable applications in heterocyclic synthesis, which are detailed in Chapter 15, or in natural product synthesis, as shown in Chapters 16–18, an area where chemical transformations are clearly challenged and pushed way beyond their limits. Copper-mediated

cross-coupling reactions have also been readily integrated in process chemistry where remarkably efficient procedures have been developed on the basis of copper catalysis as shown by examples collected in Chapter 19.

Copper is cheap, at least for the moment, but high catalyst loading is still required in most cases and catalyst recovery is therefore a quite important issue. Reusable catalysts for copper-mediated cross-coupling reactions under heterogeneous conditions have also been developed recently and will be over-viewed in Chapter 20.

As a concluding remark, all advances reported in the last decade in copper-mediated cross-coupling reactions are truly remarkable and had strong impact, in a relatively short period of time, in organic synthesis. Many of the transformations presented in this book are already classics in organic synthesis and, in many areas, copper really showed the way.

REFERENCES

- [1] Meyer, K. H. *Helv. Chim. Acta* **1940**, 23, 93–100.
- [2] Ullmann, F.; Bielecki, J. *Ber. Dtsch. Chem. Ges.* **1901**, 34, 2174–2185.
- [3] Ullmann, F. *Ber. Dtsch. Chem. Ges.* **1903**, 36, 2382–2384.
- [4] Goldberg, I. *Ber. Dtsch. Chem. Ges.* **1906**, 39, 1691–1692.
- [5] Ullmann, F.; Sponagel, P. *Ber. Dtsch. Chem. Ges.* **1905**, 38, 2211–2212.
- [6] Mowry, D. T. *Chem. Rev.* **1948**, 42, 189–283.
- [7] Ellis, G. P.; Romney-Alexander, T. M. *Chem. Rev.* **1987**, 87, 779–794.
- [8] Farbwerke vormals Meister, Lucius & Brüning, assignee. Verfahren zur Darstellung von Anthrachinon- α -nitrilen. German patent 271,790. 1913 February 18.
- [9] (a) Rosenmund, K. W.; Struck, E. *Ber. Dtsch. Chem. Ges.* **1919**, 52, 1749–1746. (b) Rosenmund, K. W.; Harms, H. *Ber. Dtsch. Chem. Ges.* **1920**, 53, 2226–2240.
- [10] (a) de Diesbach, H. *Helv. Chim. Acta* **1923**, 6, 539–548. (b) de Diesbach, H.; Schmidt, V.; Decker, E. *Helv. Chim. Acta* **1923**, 6, 548–549.
- [11] Newman, M. S. *J. Am. Chem. Soc.* **1937**, 59, 2472.
- [12] Pongratz, A. *Monatsh. Chem.* **1927**, 48, 585–591.
- [13] von Braun, J.; Manz, G. *Liebigs Ann. Chem.* **1931**, 488, 111–126.
- [14] Hurtley, W. R. H. *J. Chem. Soc.* **1929**, 1870–1873.
- [15] (a) Nicolaou, K. C.; Boddy, C. N. C.; Natarajan, S.; Yue, T.-Y.; Li, H.; Bräse, S.; Ramanjulu, J. M. *J. Am. Chem. Soc.* **1997**, 119, 3421–3422. (b) Nicolaou, K. C.; Li, H.; Boddy, C. N. C.; Ramanjulu, J. M.; Yue, T.-Y.; Natarajan, S.; Chu, X.-J.; Bräse, S.; Rübsam, F. *Chem. Eur. J.* **1999**, 5, 2584–2601. (c) Nicolaou, K. C.; Boddy, C. N. C.; Li, H.; Koumbis, A. E.; Hughues, R.; Natarajan, S.; Jain, N. F.; Ramanjulu, J. M.; Bräse, S.; Solomon, M. E. *Chem. Eur. J.* **1999**, 5, 2602–2621. (d) Nicolaou, K. C.; Koumbis, A. E.; Takayanagi, M.; Natarajan, S.; Jain, N. F.; Bando, T.; Li, H.; Hughues, R. *Chem. Eur. J.* **1999**, 5, 2622–2647. (e) Nicolaou, K. C.; Mitchell, H. J.; Jain, N. F.; Bando, T.; Hughes, R.; Winssinger, N.; Natarajan, S.; Koumbis, A. E. *Chem. Eur. J.*

- 1999**, 5, 2648–2667. (f) Nicolaou, K. C.; Boddy, C. N. C. *J. Am. Chem. Soc.* **2002**, *124*, 10451–10455.
- [16] Paine, A. J. *J. Am. Chem. Soc.* **1987**, *109*, 1496–1502.
- [17] Bryant, R. J., inventor; Sterwin A.-G., assignee. Substitution of aromatic organic compounds. G. B. patent 2,089,672. 1982 June 30.
- [18] Capdevielle, P.; Maumy, M. *Tetrahedron Lett.* **1993**, *34*, 1007–1010.
- [19] Goodbrand, H. B.; Hu, N.-X. *J. Org. Chem.* **1999**, *64*, 670–674.
- [20] Chan, D. M. T.; Monaco, K. L.; Wang, R.-P.; Winter, M. P. *Tetrahedron Lett.* **1998**, *39*, 2933–2936.
- [21] Evans, D. A.; Katz, J. L.; West, T. R. *Tetrahedron Lett.* **1998**, *39*, 2937–2940.
- [22] Lam, P. Y. S.; Clark, C. G.; Saubern, S.; Adams, J.; Winters, M. P.; Chan, D. M. T.; Combs, A. *Tetrahedron Lett.* **1998**, *39*, 2941–2944.

CONTRIBUTORS

Carole Alayrac, Laboratoire de Chimie Moléculaire et Thioorganique, UMR CNRS 6507, INC3M, FR3038, ENSICAEN & Université de Caen Basse-Normandie, 6 boulevard du Maréchal Juin, 14050 Caen, France

Irina P. Beletskaya, Moscow M.V. Lomonosov State University, Chemical Faculty, Department of Organic Chemistry, Leninskie gory 1, Moscow 119991, Russian Federation

Matthias Beller, Leibniz-Institut für Katalyse e.V. an der Universität Rostock, Albert-Einstein-Strasse 29a, 18059 Rostock, Germany

Nicolas Blanchard, Université de Strasbourg, Ecole Européenne de Chimie, Polymères et Matériaux, Laboratoire de Chimie Moléculaire Associé au CNRS, 25 rue Becquerel 67087 Strasbourg, France

Alicia Casitas, QBIS Group, Departament de Química, Universitat de Girona, Campus de Montilivi, 17071 Girona, Spain

Morgan Donnard, Université de Strasbourg, Faculté de Pharmacie, Laboratoire d'Innovation Thérapeutique Associé au CNRS, 74 route du Rhin—CS60024, 67401 Illkirch Cedex, France

Gwilherm Evano, Laboratoire de Chimie Organique, Service de Chimie et PhysicoChimie Organiques, Université Libre de Bruxelles, Avenue F. D. Roosevelt 50, CP160/06, B-1050 Brussels, Belgium

Alexey Yu. Fedorov, N.I. Lobachevsky Nizhny Novgorod State University, Chemical Faculty, Department of Organic Chemistry, Gagarin avenue 23, 603950 Nizhny Novgorod, Russian Federation

Hua Fu, Key Laboratory of Bio-organic Phosphorus Chemistry and Chemical Biology, Department of Chemistry, Tsinghua University, Beijing 100084, P. R. China

Annie-Claude Gaumont, Laboratoire de Chimie Moléculaire et Thioorganique, UMR CNRS 6507, INC3M, FR3038, ENSICAEN & Université de Caen Basse-Normandie, 6 boulevard du Maréchal Juin, 14050 Caen, France

Céline Guissart, Laboratoire de Chimie Organique, Service de Chimie et PhysicoChimie Organiques, Université Libre de Bruxelles, Avenue F. D. Roosevelt 50, CP160/06, B-1050 Brussels, Belgium

Ruimao Hua, Department of Chemistry, Tsinghua University, Beijing 100084, P. R. China

Yongwen Jiang, State Key Laboratory of Bio-organic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R. China

Kévin Jouvin, Institut Lavoisier de Versailles, Université de Versailles Saint-Quentin-en-Yvelines, 45 avenue des Etats Unis, 78035 Versailles Cedex, France

Klaus Kunz, Bayer Crop Science AG, Alfred-Nobel-Strasse 50, 40789 Monheim, Germany

Jihoon Lee, Department of Chemistry and Center for Chemical, Methodology and Library Development, Metcalf Center for Science and Engineering, 590 Commonwealth Avenue, Boston University, Boston, MA 02215, USA

Hao Li, Department of Chemistry, Atwood Hall, 1515 Dickey Drive, Emory University, Atlanta, GA, 30322-2210, USA

Jin-Heng Li, State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, P. R. China

Lanny S. Liebeskind, Department of Chemistry, Atwood Hall, 1515 Dickey Drive, Emory University, Atlanta, GA, 30322-2210, USA

Songbai Liu, Department of Food Science and Nutrition, Zhejiang University, 388 Yuhangtang Road, Hangzhou, Zhejiang 310058, P. R. China

Norbert Lui, Bayer Crop Science AG, Alfred-Nobel-Strasse 50, 40789 Monheim, Germany

Dawei Ma, State Key Laboratory of Bio-organic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R. China

Luc Neuville, Centre de Recherche de Gif, Laboratoire International Associé, Institut de Chimie des Substances Naturelles, CNRS, 91198 Gif-sur-Yvette Cedex, France

James S. Panek, Department of Chemistry and Center for Chemical, Methodology and Library Development, Metcalf Center for Science and

Engineering, 590 Commonwealth Avenue, Boston University, Boston, MA, 02215, USA

Doron Pappo, Department of Chemistry, Ben-Gurion University, P.O. Box 653, Be'er-Sheva 84105, Israel

Xavi Ribas, QBIS Group, Departament de Química, Universitat de Girona, Campus de Montilivi, 17071 Girona, Spain

Thomas Schareina, Leibniz-Institut für Katalyse e.V. an der Universität Rostock, Albert-Einstein-Strasse 29a, 18059 Rostock, Germany

Ren-Jie Song, State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, P. R. China

Marc Taillefer, Institut Charles Gerhardt Montpellier, ENSCM, 8 rue de l'Ecole Normale, 34296 Montpellier Cedex 5, France

Cédric Theunissen, Laboratoire de Chimie Organique, Service de Chimie et PhysicoChimie Organiques, Université Libre de Bruxelles, Avenue F. D. Roosevelt 50, CP160/06, B-1050 Brussels, Belgium

Anis Tlili, Institut Charles Gerhardt Montpellier, ENSCM, 8 rue de l'Ecole Normale, 34296 Montpellier Cedex 5, France

Changfeng Wan, Hefei National Laboratory for Physical Sciences at Microscale, CAS Key Laboratory of Soft Matter Chemistry and Department of Chemistry, University of Science and Technology of China, Hefei, Anhui, 230026, P. R. China

Ye Wang, Hefei National Laboratory for Physical Sciences at Microscale, CAS Key Laboratory of Soft Matter Chemistry and Department of Chemistry, University of Science and Technology of China, Hefei, Anhui, 230026, P. R. China

Zhiyong Wang, Hefei National Laboratory for Physical Sciences at Microscale, CAS Key Laboratory of Soft Matter Chemistry and Department of Chemistry, University of Science and Technology of China, Hefei, Anhui, 230026, P. R. China

Yoshihiko Yamamoto, Department of Basic Medicinal Sciences, Graduate School of Pharmaceutical Sciences, Nagoya University, Chikusa, Nagoya 464-8601, Japan

Daoshan Yang, Key Laboratory of Bio-organic Phosphorus Chemistry and Chemical Biology, Department of Chemistry, Tsinghua University, Beijing 100084, P. R. China