

Recent Developments in Copper-Catalyzed Aerobic Oxidative Coupling Reactions Using Oxygen as Oxidant


ZHE JIA

MICHIGAN STATE UNIVERSITY
DEPARTMENT OF CHEMISTRY

2/16/2011

Transition Metal Catalyzed Aerobic Oxidative Coupling Reaction

2



- Activate two different C-H bonds or H-Heteroatom bonds
- Couple them via a transition metal catalyst
- Carry out the reaction in air or O₂, with O₂ as the oxidant


Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

Gligorich, K. M. Sigman, M. S. *Angew. Chem. Int. Ed.* **2006**, 45, 6612-6615.

Gligorich, K. M. Sigman, M. S. *Chem. Commun.*, **2009**, 3854-3867.

Copper Catalyzed Aerobic Oxidative Coupling Reaction


3



Why copper?

- Copper is relatively abundant in the earth's crust.
- Copper can bind and activate dioxygen well that perform a variety of critical biological functions.

Glaser Coupling (1869)



Gamez, P.; Aubel, P. G.; Driessen, W. L.; Reedijk, J. *Chem. Soc. Rev.*, **2001**, 30, 376-385.

Lewis, E. A.; Tolman, W. B. *Chem. Rev.*, **2004**, 104, 1047-1076.

Glaser, C. *Ber. Dtsch. Chem. Ges.* **1869**, 2, 422.

Glaser, C. *Ann. Chem. Pharm.* **1870**, 154, 137.

Pros and Cons of Oxidative Coupling Reaction

4

Attractions:

- ❖ Direct C-H functionalization
- ❖ High atom economy efficiency
- ❖ H₂O is generated as by-product environmentally friendly

Challenges:

- ❖ Homocoupling of the two nucleophiles
- ❖ Direct reaction of the nucleophiles with the oxidant


Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.


Stahl, S. S. *Science*, **2005**, 309, 1824-1826.

Gligorich, K. M.; Sigman, M. S. *Chem. Commun.*, **2009**, 3854-3867.


Transition Metal Catalyzed Coupling Reaction Pattern

5



 Nucleophile
  Electrophile

M = Fe, Co, Ni, Cu, Pd, Ru, Rh




Models for Coupling Reaction Bond Construction

6

➤ Classic Coupling Model





➤ Oxidative Coupling Model



To Combine Two Nucleophiles Together

7





Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

Gligorich, K. M.; Sigman, M. S. *Angew. Chem. Int. Ed.* **2006**, 45, 6612-6615.

To Combine Two Nucleophiles Together

8



Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

Gligorich, K. M.; Sigman, M. S. *Angew. Chem. Int. Ed.* **2006**, 45, 6612-6615.

Why O₂ as Oxidant?

9

O₂ is the quintessential oxidant for chemical synthesis!

- Huge abundance in nature
- Available at virtually no cost
- Produces no environmentally hazardous by-products


Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

Stahl, S. S. *Science* **2005**, 309, 1824-1826.


Metalloenzyme-catalyzed Aerobic Oxidation

10

a) Oxygenase Pathway



b) Oxidase Pathway



- In the **oxygenase pathway (a)**, the substrate oxidation involves oxygen-atom transfer from molecular oxygen.
- The **oxidase pathway (b)**, utilizes molecular oxygen as a electron/proton acceptor in the oxidation of the substrate.

Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.


Stahl, S. S. *Science* **2005**, 309, 1824-1826.

Examples of Oxygenase and Oxidase


11

C-N Functionalization of Terminal Alkynes

Stahl – Oxidase Pathway



Jiao – Dioxygenase Pathway




Hamada, T.; Ye, X.; Stahl, S. S. *J. Am. Chem. Soc.*, **2008**, 130, 833–835.

Zhang, C.; Jiao, N. *J. Am. Chem. Soc.*, **2010**, 132, 28–29.

Metal Catalyzed Oxidase Reactions

12




Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

<http://www.chem.wisc.edu/content/investigation-fundamental-reactions-between-palladium-and-molecular-oxygen>

Metal Catalyzed Oxidase Reactions

13




Stahl, S. S. *Angew. Chem. Int. Ed.* **2004**, 43, 3400 – 3420.

<http://www.chem.wisc.edu/content/investigation-fundamental-reactions-between-palladium-and-molecular-oxygen>

Outline

14

- Cu-Catalyzed Oxidative **Homo-Coupling** Reaction




- Cu-Catalyzed Oxidative **Hetero-Coupling** Reaction

- C-C Bond Formation
- C-N Bond Formation
- C-P Bond Formation


Cu Catalyzed Aerobic Oxidative C-C Bond Homocoupling Reaction

15

Glaser Coupling (1869)



One Possible Mechanism Pathway:




Glaser, C. *Ber. Dtsch. Chem. Ges.* **1869**, 2, 422.

Glaser, C. *Ann. Chem. Pharm.* **1870**, 154, 137.


Arene Dimerization

16

Aromatic Glaser-Hay Reaction




Copper-catalyzed, deprotonative dimerization of arenes by employing O₂ as the terminal oxidant.



Aromatic Glaser-Hay Reaction

17

Previous Work



Do, H.-Q.; Daugulis, O. *J. Am. Chem. Soc.* **2007**, 129, 12404-12405.


Do, H.-Q.; Daugulis, O. *J. Am. Chem. Soc.* **2008**, 130, 1128.-1129

Do, H.-Q.; Khan, R. M. K.; Daugulis, O. *J. Am. Chem. Soc.* **2008**, 130, 15185-15192.

Aromatic Glaser-Hay Reaction


18

Inspiration from Previous Work



Initial Study on Aromatic Glaser-Hay Reaction

19




A less polarized C-metal bond in the intermediate is needed!

Solution

Formed by ArLi intermediate with O_2 or ArCu with hydroxide derived from water.

Substrate Scope of Aromatic Glaser-Hay Reaction

20



- Hindered Zn and Mg amide bases were used
- Less acidic the arene H, the stronger base was needed




Base

Dicyclohexylamine :
iPrMgCl•LiCl : ZnCl₂
(1.1 : 1 : 0.25)

iPrMgCl•LiCl :
tetramethylpiperidine : ZnCl₂
(1 : 1.1 : 0.5)

iPrMgCl•LiCl :
tetramethylpiperidine
(1.1 : 1)

Product




Outline

21

- Cu-Catalyzed Oxidative Homo-Coupling Reaction
- Cu-Catalyzed Oxidative **Hetero-Coupling** Reaction

- **C-C** Bond Formation




- C-N Bond Formation

- C-P Bond Formation

Cu-Catalyzed Direct Aerobic Alkynylation of Arenes with Terminal Alkynes


22

Direct Alkynylation




The first example of direct alkynylation of an aromatic C-H bond with terminal alkynes.

...but limited to polyfluoroarenes!




Alkynylation of Arenes with Terminal Alkynes

23



Sonogashira (1975)





Armin de Meijere, François Diederich, *Metal-Catalyzed Cross-Coupling Reactions, 2nd, Completely Revised and Enlarged Edition*, Wiley-VCH, Weinheim, **2004**

Wei, Y.; Zhao, H.; Kan, J.; Su, W. *J. Am. Chem. Soc.*, **2010**, 132, 2522–2523.

Alkynylation of Arenes with Terminal Alkynes

24





Kobayashi, K.; Arisawa, M.; Yamaguchi, M. *J. Am. Chem. Soc.* **2002**, *124*, 8528-8529.

Amemiya, R.; Fujii, A.; Yamaguchi, M. *Tetrahedron Lett.* **2004**, *45*, 4333-4335.

Seregin, I. V.; Ryabova, V.; Gevorgyan, V. *J. Am. Chem. Soc.* **2007**, *129*, 7742-7744.


Cu-Catalyzed Direct Aerobic Alkynylation of Arenes with Terminal Alkynes

25



Cu-Catalyzed Aerobic Alkynylation of Pentafluorobenzene with Phenylacetylene

26




Entry	Reaction Condition (equiv. of reagents)	Solvent	% Yield 3 (4/5)
1	CuCl ₂ (20 mol%), O ₂ (1atm), NaHCO ₃ /K ₃ PO ₄ (2equiv.)	DMSO, 70°C	undetectable (>90/-)
2	CuCl ₂ (20 mol%), O ₂ (1atm), tBuOLi (2equiv.)	DMSO, 40°C	11 (57/5)
3	CuCl ₂ (20 mol%), O ₂ (1atm), tBuOLi (2equiv.), 1,10-phenanthroline (0.2equiv.)	DMSO, 40°C	49 (38/4)
4	CuCl ₂ (20 mol%), O ₂ (1atm), tBuOLi (2equiv.), 1,10-phenanthroline (0.2equiv.) DDQ (0.15equiv.)	DMSO, 40°C	72 (24/7)
5	CuCl ₂ (30 mol%), O ₂ (1atm), tBuOLi (3equiv.), 1,10-phenanthroline (0.3 equiv.) DDQ (0.15equiv.)	DMSO, 40°C	85 (14/13)

Wei, Y.; Zhao, H.; Kan, J.; Su, W.; *J. Am. Chem. Soc.*, **2010**, 132, 2522–2523.

Proposed Mechanism of Cu-Catalyzed Direct Aerobic Alkynylation

27


Cross-Coupling Pathway



Proposed Mechanism of Cu-Catalyzed Direct Aerobic Alkynylation

28


Cross-Coupling Pathway



Proposed Mechanism of Cu-Catalyzed Direct Aerobic Alkynylation








29

Homo-Coupling and Cross-Coupling Pathway








Substrate Scope of Cu-Catalyzed Direct Aerobic Alkynylation

30





Substrate Scope of Cu-Catalyzed Direct Aerobic Alkynylation

31



Substrate Scope of Cu-Catalyzed Direct Aerobic Alkynylation

32



Both tri- and difluoroarenes are unreactive.

Cu Mediated Aerobic Oxidative Trifluoromethylation of Terminal Alkynes

33

The first example of a copper-mediated oxidative trifluoromethylation.



- In situ generated CuCF_3
- Direct oxidative coupling of terminal alkynes


34




Initial Study on Aerobic Oxidative Trifluoromethylation

35

Under dry air



Under O₂




Reactive CuCF₃
was quenched by
high concentration
of O₂

Chu, L.; Qing, F. *J. Am. Chem. Soc.*, **2010**, 132, 7262–7263.

Wiemers, D. M.; Burton, D. J. *J. Am. Chem. Soc.*, **1986**, 108, 832–834.


Plausible Reaction Pathways for Aerobic Oxidative Trifluoromethylation

36










Plausible Reaction Pathways for Aerobic Oxidative Trifluoromethylation

37



Substrate Scope of Aerobic Oxidative Trifluoromethylation

38




Outline

39

- Cu-Catalyzed Oxidative Homo-Coupling Reaction
- Cu-Catalyzed Oxidative **Hetero-Coupling** Reaction

- C-C Bond Formation

- **C-N** Bond Formation




- C-P Bond Formation


Amidation of Terminal Alkynes

40

Coupling via Alkynyl Halide



Direct Coupling of Terminal Alkynes




Hamada, T.; Ye, X.; Stahl, S. *J. Am. Chem. Soc.*, **2008**, 130, 833–835.

DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, 110, 5064–5106.

Cu Catalyzed Aerobic Oxidative Amidation of Terminal Alkynes

41




Entry	Reaction Condition (equiv. of reagents)	Solvent	% Yield 3 (4/5)
1	CuCl ₂ (2), Cs ₂ CO ₃ (2), 1 equiv. of 2	DMSO 70°C	26 (42/17)
2	CuCl ₂ (2), Cs ₂ CO ₃ (2), 5 equiv. of 2	DMSO 70°C	89 (4/-)
3	CuCl ₂ (0.2), Cs ₂ CO ₃ (2), 5 equiv. of 2	DMSO 70°C	Trace (19/-)
4	CuCl ₂ (0.2), Na ₂ CO ₃ (2), Pyridine (2), 5 equiv. of 2	DMSO 70°C	90 (4/-)
5	CuCl ₂ (0.2), Na ₂ CO ₃ (2), Pyridine (2), 1 equiv. of 2	Toluene 70°C	69 (16/4)

Hamada, T.; Ye, X.; Stahl, S. *J. Am. Chem. Soc.*, **2008**, 130, 833–835.


Mechanistic Study of Cu Catalyzed Aerobic Oxidative Amidation of Terminal Alkynes

42





Proposed Mechanism of Cu Catalyzed Aerobic Oxidative Amidation of Terminal Alkynes

43



Cu Catalyzed Direct Amination of Benzothiazoles and Benzoxazoles


44



Outline


45

- Cu-Catalyzed Oxidative Homo-Coupling Reaction
- Cu-Catalyzed Oxidative **Hetero-Coupling** Reaction
 - C-C Bond Formation
 - C-N Bond Formation
 - **C-P** Bond Formation




Cu Catalyzed Aerobic Phosphonation of sp C

46




A new methodology for catalytically constructing C-P bonds:



Cu Catalyzed Aerobic Phosphonation of sp C

47

Under dry air




Gao, Y.; Zhao, Y.; Zhou, Y.; Han, L. *J. Am. Chem. Soc.*, **2009**, 131, 7956–7957.

Nakamura, E.; Mori, S. *Angew. Chem. Int. Ed.* **2000**, 39, 3750–3771.

Cu Catalyzed Aerobic Phosphonation of sp C

48

Under N₂




Gao, Y.; Zhao, Y.; Zhou, Y.; Han, L. *J. Am. Chem. Soc.*, **2009**, 131, 7956–7957.

Nakamura, E.; Mori, S. *Angew. Chem. Int. Ed.* **2000**, 39, 3750-3771.

Niu, M.; Fu, H.; Jiang, Y.; Zhao, Y. *Chem. Commun.* **2007**, 272-274.


Proposed Reaction Path Cu Catalyzed Aerobic Phosphonation of sp C

49






Proposed Reaction Path Cu Catalyzed Aerobic Phosphonation of sp C

50





Substrate Scope for Cu Catalyzed Aerobic Phosphonation of sp C

51




Cu Catalyzed Aerobic Phosphonation of sp^3 C

52





Mechanistic Study of Cu Catalyzed Aerobic Phosphonation of sp^3 C

53




Aryl-protected amines is an important feature.



Aryl group can stabilize the oxidized form of the tertiary amine.

Mechanistic Study of Cu Catalyzed Aerobic Phosphonation of sp^3 C

54



A molecular oxygen uptake experiment shows


- a total consumption of 1 equiv. of O_2 for the generation of 1 equiv. of desired product **3**;
- 1 equiv. of **1** and 1 equiv. of **2** consumes 0.5 equiv. of O_2 .



O_2 is involved in the oxidation of excess **2** to methyl phosphate


Possible Mechanism Pathway for Cu Catalyzed Aerobic Phosphonation of sp^3 C

55



Substrate Scope of Cu Catalyzed Aerobic Phosphonation of sp^3 C

56



Entry	Amine	Phosphite	Product	% Yield
1				79 (95)
2	 PMP = p-methoxyphenyl			73 (75)
3	 OMP = o-methoxyphenyl			69 (85)

Summary

57

- Copper-catalyzed aerobic oxidative coupling reactions enable direct C-H functionalization, and can be a powerful tool for constructing C-C bonds and C-heteroatom bonds.
- Reactions are conducted in air or an O₂ atmosphere, with O₂ as a stoichiometric oxidant. H₂O or H₂O₂ is generated as a by-product.
- However, this chemistry is still in its infancy. The substrate scope is very limited for some reactions.

Acknowledgement

58

- Dr. Baker
- Dr. Maleczka
- Dr. Smith, Dr. Wulff
- Baker's Group: Gina, Hui, Heyi, Quanxuan, Wen, Yiding, Greg, Salinda, Cat and Kayla
- Li, Hong, Yong, Chunjuan and Hao
- Xin, Ipek, Herbert, Wenjun and Jason

THANK YOU!