The model were using and what it tells ud

Lee Carter

The Lee-carter model was introduced in 1992 by Ronald Lee and Lawrence Carter in their paper "Modeling and Forecasting U.S. Mortality". It revolutionized mortality forecasting by providing a data-driven approach in lieu of the expert-led methods that were previously dominant. Prior to its development, official forcasts often underestimated the gains in human life expectancy, creating a pressing need for more accurate projections to asses the solvency of pension systems and social security programs.

The models core innovation was its ability to reduce the matrix of log age specific death rates into a single time varying mortality index K_t an age specific sensitivity parameter b_x. and a baseline mortality rate a_x. To achieve this, Lee and Carter applied Singular Value Decomposition (SVD) to extract the characteristic pattern of mortality improvement over time. Unfortunately, the solution to the SVD is not unique, so they imposed two constraints: the sum of b_x across ages must equal 1, and the sum of K_t across time must equal 0. This allowed them to identify a unique solution that captured the dominant trend in mortality improvement. Then by only taking the first singular value and its corresponding vectors in a_x and b_x, they reduce the complexity of the data while retaining the most important information. This serves as compression of the data reducing the number of parameters needed to model mortality trends but keeping the essential features of the data. Following this, the time-varying mortality index K_t is modeled as a random walk with drift, allowing for the creation of objective prediction intervals that provided a quantifiable measure of uncertainty. This was a significant departure from previous "best-guess" forecasts, which lacked any formal way to assess the range of possible outcomes.

The simplicity and robustness of the Lee-Carter model allowed it to outperform contemporary models in backtesting, closely tracking actual mortality data while older models significantly under-predicted life expectancy gains. Its ability to capture the linear trend of log-mortality decline across the 20th century made it particularly effective for long-term forecasting. Unlike earlier parametric models that focused on the biological shape of the mortality curve at a single point in time, the Lee-Carter model prioritized the dynamic movement of that curve over decades.

Thanks to the nature of the Lee-Carter model extensions and variations have been developed to address its limitations and improve its accuracy. For example, the LC-ER (Extended Rotation) method was developed to account for "rotation" in mortality decline, which addresses implausible age-pattern artifacts in vanilla LC projections. Additionally, the stmomo R package, whose implementation we will be using later, provides a flexible framework for mortality forecasting using the Lee-Carter model as well as several other models, allowing users to fit the model to various data types and generate forecasts while assessing uncertainty.

The model and its extensions have been used in various applications, including forecasting breast cancer mortality in China and Pakistan, where it outperformed simpler models for the screen-age/late-onset population (ages 50–84), revealing rising mortality in Pakistan versus declining rates in China, which was used to inform targeted healthcare interventions. The UN employs the LC model to forecast global mortality trends, especially for long-term projections up to 2100. For high-life-expectancy countries like Japan, the model was extended to account for "rotation" in mortality decline (slowing improvements at young ages and accelerating declines at older ages). This adjustment, termed the LC-ER (Extended Rotation) method, addressed implausible age-pattern artifacts in vanilla LC projections (**booth_baselini_carmada_2023?**)

Now, when talking about the implementation we will be using, Stochastic Mortality Modeling (stMoMo) is an R package that provides a comprehensive framework for fitting and forecasting mortality models, including the Lee-Carter model. The stMoMo package allows users to easily fit the Lee-Carter model to their mortality data, generate forecasts, and assess uncertainty through confidence intervals. It also includes various extensions of the Lee-Carter model, such as the LC-ER method, which can be used to improve the accuracy of forecasts for high-life-expectancy countries. The package is designed to be user-friendly and flexible, making it accessible for researchers and practitioners in the field of mortality forecasting. For out use case, we will be using the base Lee-Carter model and using its initial fit to bootstrap its parameters to then simulate numerous forecasts to create a distribution of possible outcomes. For this we need to set several parameter. First would be the link function, which determines the random component associated with the mortality model. The default link function is the log link, this assumes that the deaths follow a Poisson distribution, which is appropriate for count data like mortality. Secondly, we need to set the restraints for the model to ensure identifiability. For this we will set it as "sum". This ensures that $\Sigma_t k_t = 0$. With these two we can now create the seed fit for the bootstrapping process. The bootstrap itself requires its own set of parameters, these being the number of bootstrap samples, the approach to be applied and the type of death to resample. After several cycles, we determined that 100 samples generated properly distributed residuals, but for robustness, we choose 500 to provide a more robust fit and still iterate at an acceptable pace. Following this we proceed to the simulations. The needed parameters for this are the number of simulations which we will set as the same as the number of bootstraps, the years to project set at 50. The jump choice, whether to use the fitted end or the actual end, we will choose the actual end in the data. Lastly for the simulation is the method of the projection of kt. Our alternatives are multivariate random

walk with drift and an N independent arima models. Lee and Carter stated they are equivalent so we will use the random walk.

Once all this is completed, we have all our needed products. Starting off with the ax parameter, this is our result.



Here, we can see the drastic fall of the baseline log mortality after the first year of life. This is followed by the steady increase as one ages and the telltale male mortality hump starting at the teen years.

Next we have the bx parameter, which is our age specific sensitivity to changes in the mortality index kt.



Here we can see the sensitivity of the mortality index to changes in mortality. The highest sensitivity is at early life. This is due to the monumental work done on the island in the latter half of the 20th century, better sanitation, better nutrition, and better healthcare. The high point is followed by an significant drop for both sexes at the teen years. Women then transition to a steady decline in sensitivity but males continue to drop until near zero at the age of 20 finally making a slow climb through to meet the female sensitivity at around 80 years of age.

Finally we have the kt parameter, which is our time-varying mortality index.



The index corresponds to the average level of mortality for that year. There are fluctuations in the index, a spike in the mid 1960s, The effect of HIV in the mid 1990s but the overall trend is a steady decline. The index starts off higher for women than men but they close the gap by the mid 1970 and start pulling apart as time passes leading to an slow growing gap, but both share the same general shape.

Now that we have the initial fit, we can move on to the simulations. We provide the initial fit to the bootstrap method, which resamples the data to create new sets of parameters. These new parameters come with their own residuals to evaluate which we use to determine the number of bootstraps needed to create a robust distribution. Earlier, we stated that 500 samples were sufficient and these are the corresponding residuals.

AX residuals



Table 1: Bx parameter residuals of the bootstrap.

Age	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
0	0.0098	(-0.0358 - 0.0485)	0.0843	0.0043	(-0.0297 - 0.0281)	0.0578	0.685
10	0.0213	(-0.0729 - 0.1204)	0.1933	0.0436	(-0.0597 - 0.1439)	0.2035	1.053
20	-0.0008	(-0.0446 - 0.0405)	0.0852	0.0121	(-0.0555 - 0.0818)	0.1373	1.612
30	0.0043	(-0.0425 - 0.0389)	0.0815	0.0073	(-0.0379 - 0.0402)	0.0781	0.959
40	-0.0033	(-0.0294 - 0.0346)	0.0640	0.0057	(-0.0318 - 0.0543)	0.0861	1.346
50	-0.0006	(-0.0281 - 0.0299)	0.0581	0.0060	(-0.0173 - 0.0338)	0.0512	0.882
60	-0.0017	(-0.0263 - 0.0156)	0.0419	0.0035	(-0.0229 - 0.0209)	0.0438	1.045

Age	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
70	-0.0024	(-0.0173 - 0.0179)	0.0352	-0.0036	(-0.0164 - 0.0146)	0.0310	0.880
80	-0.0001	(-0.0148 - 0.0188)	0.0336	0.0021	(-0.0106 - 0.0145)	0.0251	0.747

Here we have the ax residuals of the bootstrap. It presents a large deviance spike around childhood. For males this tapers quickly when entering the teen years and continues to narrow through adulthood and the senior years. Women on the other hand present a slower narrowing into adulthood and further slows down past the 40 year mark. Looking at the quantiles, at the median there's a clear skew towards opposite sides of zero between the sexes and the range between the 95% quantiles between men and woman are withing 20% of each other. In short both share the same general shape of uncertainty with the males having a slight offset towards the negative. This finally leads to males having a lower baseline level of log mortality than females

BX residuals



Table 2: Bx parameter residuals of the bootstrap.

Age	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
0	-8e-04	(-0.005 - 0.0064)	0.0114	0.0002	(-0.0024 - 0.0031)	0.0055	0.482
10	0e+00	(-0.017 - 0.0158)	0.0328	-0.0017	(-0.0083 - 0.0038)	0.0121	0.368
20	-1e-04	(-0.0066 - 0.0058)	0.0124	0.0004	(-0.0043 - 0.0063)	0.0105	0.852
30	7e-04	(-0.0067 - 0.0058)	0.0125	0.0003	(-0.0054 - 0.0043)	0.0097	0.772
40	6e-04	(-0.0042 - 0.0066)	0.0108	-0.0001	(-0.0044 - 0.0034)	0.0078	0.724
50	1e-04	(-0.0044 - 0.0039)	0.0083	-0.0001	(-0.0024 - 0.0039)	0.0063	0.759
60	4e-04	(-0.0026 - 0.0027)	0.0052	0.0004	(-0.0016 - 0.0028)	0.0044	0.833
70	4e-04	(-0.002 - 0.0031)	0.0051	0.0001	(-0.0016 - 0.0022)	0.0038	0.743
80	9e-04	(-0.0026 - 0.0038)	0.0064	0.0005	(-9e-04 - 0.002)	0.0029	0.447

Moving on to the bx parameters bootstrap residuals, in many ways it mirrors the ones from the ax parameter, with a large deviance spike around childhood that tapers off as age progresses. However, unlike the ax residuials, we see a clear difference in the range of the residuals between sexes. For males, the residuals are much more spread out than the female counterpart, with the 95% range ratio being ranging between 0.50 and 0.94, meaning that the male sensitivity to changes in the mortality index is much more volatile than the female sensitivity.

KT residuals



Table 3: Bx parameter residuals of the bootstrap.

Year	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
1950	-0.0466	(-0.5437 - 0.5573)	1.1010	-0.0744	(-0.7538 - 0.4017)	1.1555	1.050
1960	-0.0547	(-0.7197 - 0.535)	1.2548	-0.0203	(-0.6676 - 1.0128)	1.6804	1.339
1970	-0.0888	(-0.7766 - 0.7084)	1.4850	0.0083	(-1.1084 - 0.8412)	1.9495	1.313
1980	-0.1857	(-1.1007 - 0.9779)	2.0785	-0.0782	(-1.7967 - 1.0985)	2.8953	1.393
1990	-0.3443	(-1.686 - 0.8762)	2.5622	-0.0035	(-1.3126 - 1.302)	2.6146	1.020
2000	0.1126	(-1.0921 - 1.5416)	2.6337	0.1011	(-1.5743 - 1.2266)	2.8009	1.064
2010	0.5677	(-1.2499 - 1.9255)	3.1754	0.0893	(-1.0906 - 1.5513)	2.6420	0.832

Year	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
2020	0.0134	(-1.4942 - 1.7533)	3.2475	0.2578	(-1.1727 - 1.4896)	2.6623	0.820

Moving on to the residuals for the kt variable, we see that unlike both other parameters, kt slowly fans out over time. We also see the medians are likewise, slowly climbing as time progresses unlike the two other parameters and this is consistent for both sexes. Meanwhile the range ratio between the sexes is shifting first skewed towards females but now centers around 0.92.

Now, that we have the bootstrap parameters, we can now do the simulations. The simulations are done by taking the bootstrapped parameters and simulating the kt variable forward in time using a random walk with drift. This allows us to create a distribution of possible future mortality scenarios based on the uncertainty in the parameters. The results of the simulations can be used to create confidence intervals for the future mortality rates. The parameters that define the simulations are the number of simulations, the years to project, the jump choice, and the method of projection for kt. For our case, we set the number of simulations to 500, the same as the bootstrap samples, the years to project, 50 in our case, the jump choice, the actual end in our data. The method of projection for kt, multivariate random walk with drift.

Simulated kt



Table 4: Kt simulations 2022-2050.

Year	Median Male	Range Male	Range Difference Male	Median Female	Range Female	Range Difference Female	Range Ratio
2030	-20.5919	(-22.0042 - -7.51)	14.4941	-28.7410	(-29.8972 - -14.2961)	15.6011	1.076
2040	-26.5211	(-27.7995 - -7.768)	20.0315	-37.2430	(-39.0038 - -15.726)	23.2778	1.162
2050	-31.6553	(-33.6041 - -9.0678)	24.5363	-44.5686	(-46.8984 - -18.4907)	28.4077	1.158
2060	-36.3478	(-38.5131 - -10.361)	28.1520	-51.3027	(-53.8427 - -21.5232)	32.3195	1.148
2070	-41.1745	(-43.5647 - -11.9991)	31.5656	-58.3640	(-61.113 - -24.9683)	36.1447	1.145

Now, looking at the projections for the kt variable, we see that both medians fall at a steady rate over time. As expected of the random walk both ranges spread out over time, but the

range ratio is steady between sexes and roughly 1.1. Something of note is the behavior of the projected median. It almost falls to the lower bound for both sexes, which is a sign of a skewed distribution.

Death rates

Now, with the kt projections, we can now calculate the projected mortality rates for each age and sex by using the ax and bx parameters. The projected mortality rates can be calculated using the formula: log(mortality_rate) = ax + bx * kt. This allows us to create a distribution of possible future mortality rates based on the uncertainty in the parameters and the projections of kt. The projected mortality rates can be used to create confidence intervals for the future mortality rates.

Table 5: Kt simulations 2022-2050.

age	year	central	lower	upper	type	sex
20	2000	-7.7218	NA	NA	historic	female
20	2025	-8.0941	-7.7588	-8.4120	forecast	female
20	2050	-8.9213	-8.0298	-9.8497	forecast	female
40	2000	-6.3572	NA	NA	historic	female
40	2025	-6.6798	-6.4657	-6.8784	forecast	female
40	2050	-7.2041	-6.6341	-7.7776	forecast	female
60	2000	-4.8466	NA	NA	historic	female
60	2025	-5.1118	-4.9598	-5.2565	forecast	female
60	2050	-5.4883	-5.0817	-5.8932	forecast	female
20	2000	-6.0309	NA	NA	historic	male
20	2025	-6.1839	-6.1643	-6.2130	forecast	male
20	2050	-6.1958	-6.1137	-6.3102	forecast	male
40	2000	-5.4020	NA	NA	historic	male
40	2025	-5.3775	-5.2496	-5.5105	forecast	male
40	2050	-5.5914	-5.2732	-5.9461	forecast	male
60	2000	-4.1723	NA	NA	historic	male
60	2025	-4.3596	-4.2324	-4.4901	forecast	male
60	2050	-4.5745	-4.2560	-4.8997	forecast	male

Here we finally see one of our final products, the death rates. Using the median projection and dividing by age group and sex, we see that minors in both sets are projected to have a continued accelerated mortality decline through the projected period thogu their mortality is highly eratic especially by the end of our historic data. Adult women are also projected to have a continued decline in mortality but at a slower pace than minors, while mortality is projected to plateau for young adult. Those that survive to late adulthood are projected to continue their decline in mortality but at a much slower pace than that of women. Finally for both senior groups we see a continued decline in mortality with women projected to have a much faster decline. In general, female morality is projected to decline faster with considerably large gaps between the age groups while male mortality is projected to decline at a much slower pace with smaller gaps between the age groups.

Now for our final products. Proper life tables for both sexes and the life expectancy at birth projected through the entire period.

Table 6: Kt simulations 2022-2050.

sex	year	age	n	qx	lx	dx	Lx	Tx	ex	type
female	2000	0	1	0.0091	1e+05	907	99156.09	8063352	80.6335	historic
female	2010	0	1	0.0066	1e+05	660	99381.47	8285586	82.8559	historic
female	2020	0	1	0.0077	1e+05	771	99279.76	8540751	85.4075	historic
female	2030	0	1	0.0045	1e+05	446	99579.43	8782896	87.8290	central

sex	year	age	n	qx	lx	dx	Lx	Tx	ex	type
female	2040	0	1	0.0030	1e+05	300	99715.92	9010147	90.1015	central
female	2050	0	1	0.0020	1e+05	197	99812.91	9252561	92.5256	central
female	2060	0	1	0.0013	1e+05	131	99875.35	9495096	94.9510	central
female	2070	0	1	0.0008	1e+05	83	99920.92	9777405	97.7740	central
male	2000	0	1	0.0108	1e+05	1083	98997.54	7270175	72.7017	historic
male	2010	0	1	0.0085	1e+05	854	99204.16	7537447	75.3745	historic
male	2020	0	1	0.0065	1e+05	653	99387.89	7711881	77.1188	historic
male	2030	0	1	0.0054	1e+05	544	99488.46	7782675	77.8268	central
male	2040	0	1	0.0034	1e+05	339	99679.35	7955056	79.5506	central
male	2050	0	1	0.0022	1e+05	222	99789.32	8114791	81.1479	central
male	2060	0	1	0.0015	1e+05	146	99861.14	8278732	82.7873	central
male	2070	0	1	0.0009	1e+05	93	99911.42	8456161	84.5616	central



Seeing the life expectancy at birth projections, we see that the historic data shows a steady increase in life expectancy. The projected values agree with this trend, showing a continued increase in life expectancy of about 2.4 years per decade for women and 1.7 years per decade for men. The median of the projection reaches 97 years for women and 84 for men. The confidence intervals on the otherhand present a curious picture. The projection intervals are large enough to warrant a closer look, in particular, the upper bound of the female life expectancy projection presents exponential growth and is projected to reach 107 years by 2070

and male life expectancy reaches 95 year. The lower bound for both seem much more plausible, with the female projection reaching 89 years and the male 76.

References