

Effective Dynamics and Matter Coupling

Abstract

The effective dynamics of the Æther-flow Interpretation of Relativity are specified by the Einstein–Hilbert action with ordinary matter and an optional cosmological constant. This action is adopted, not derived from the unresolved source structure. We fix the units of the action and stress tensor, derive the bulk field equation, distinguish total from separate component conservation, and state the assumptions behind particle, light, and clock descriptions. The weak-field calculation recovers Poisson’s equation and leading Newtonian acceleration in the stated regime. These results define a comparison target for source research. A separate recovery criterion requires independently specified source laws, an explicit reconstruction, and control of any approximation or added assumption. The present calculation establishes consequences of the general-relativistic action, not a source origin of that action or evidence of artificial-intelligence research performance.

1 What is being specified

An interpretation becomes a predictive physical model only after its fields, equations, and measurement rules have been fixed. In this framework, the effective gravitational choice is explicit: retain general relativity (GR), rather than introduce an additional low-energy Æther-flow field. The purpose of this paper is to state that choice in a form that can be varied, checked, and used in calculations.

The distinction from the source proposal is essential. Foundations uses $\mathcal{A}$ for a smooth four-dimensional source arena and Φ_{src} for an unresolved source-order or evolution structure. Neither is identified here with physical spacetime or an observer velocity. The effective fields instead live on a separate manifold M and are governed by the action below. No inference from the source assumptions to that action is claimed.

The project seeks a source construction that reproduces GR rather than changes its predictions. This paper supplies the independently adopted target dynamics for that investigation. It also explains the effective flow interpretation, but an interpretation of a chosen action is not a derivation of that action. If a source candidate fails to reproduce the target, retaining GR as a reference does not repair or validate that candidate.

2 Conventions and model specification

The spacetime (M, g) is a smooth, time-oriented four-dimensional Lorentzian manifold, with signature $(-+++)$ and Levi-Civita connection ∇ . Greek indices range over 0, 1, 2, 3. The curvature convention is

$$[\nabla_\mu, \nabla_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma, \quad R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}. \quad (1)$$

Coordinates in the covariant action have dimensions of length; in a time-adapted chart $x^0 = ct$. When a metric is instead written with t in seconds, the displayed dt and factors of c implement that coordinate change. Tensor equations do not depend on the chosen chart.

The matter fields are denoted collectively by ψ . Their action must be specified for an actual prediction. “Ordinary matter” here means a matter model with the appropriate locally Lorentz-invariant laws and standard coupling to g ; it does not mean every arbitrary Lagrangian containing a metric. Stability, characteristics, and interactions of a chosen matter sector require their own assessment.

The intended source account would recover this shared metric coupling rather than add a direct force between matter and an Æther-flow wind. Its absence from the adopted action is explicit; its absence from a future source reduction must be demonstrated. Coupling matter only to one metric is not, by itself, a proof that the full theory has no observable preferred structure. Source-dependent gravitational dynamics or restrictions on solutions could still change the comparison. The reconstruction must therefore account for the metric dynamics and admissible solutions as well as the matter coupling.

The free constants include $G > 0$ and a constant Λ . Initial and boundary conditions, topology, and the domain of the solution must also be supplied for a particular physical problem. They are not selected by the phrase exact closure.

3 Action and bulk field equation

The gravitational benchmark is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi]. \quad (2)$$

Setting $\Lambda = 0$ recovers the Einstein–Hilbert form

$$S_{\text{eff}}|_{\Lambda=0} = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} R + S_m[g, \psi]. \quad (3)$$

Both S_m and the gravitational term have units of action. Since d^4x has dimensions L^4 and R has dimensions L^{-2} in this coordinate convention, the factor c^3/G provides the required normalization.

Define the physical stress–energy tensor by

$$T_{\mu\nu} := -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}. \quad (4)$$

The factor of c is required when $x^0 = ct$ and S_m is a physical action. An equivalent convention could rescale the matter action, but then its variational definition would need to say so. With (4),

$$\delta S_m = -\frac{1}{2c} \int_M d^4x \sqrt{-g} T_{\mu\nu} \delta g^{\mu\nu} \quad (5)$$

on the matter equations, apart from boundary terms associated with the chosen variations.

For compactly supported metric variations, the bulk variation of the full action is

$$\delta S_{\text{eff}} = \int_M d^4x \sqrt{-g} \left[\frac{c^3}{16\pi G} (G_{\mu\nu} + \Lambda g_{\mu\nu}) - \frac{T_{\mu\nu}}{2c} \right] \delta g^{\mu\nu}. \quad (6)$$

Stationarity for arbitrary such variations gives

$$\boxed{G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}}. \quad (7)$$

For $\Lambda = 0$, this becomes

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (8)$$

The calculation is the standard metric variation of GR, with the unit convention made explicit (Carroll, 1997). It establishes what follows from (2), not why a source ontology should produce that action.

A finite-boundary problem needs an appropriate boundary action and boundary data. The Einstein–Hilbert integrand alone contains boundary variations involving derivatives of δg . Compact support avoids that issue for the bulk derivation; it is not a substitute for specifying boundary terms when the physical problem requires them.

4 Cosmological constant and other dark-energy models

A cosmological constant can be kept on the geometric side of (7), or moved to the matter side as the specific tensor

$$T_{\mu\nu}^{(\Lambda)} = -\frac{\Lambda c^4}{8\pi G} g_{\mu\nu}. \quad (9)$$

These are exactly equivalent descriptions of the same constant term, not two independent sources to be counted together. In perfect-fluid language, with ρ denoting mass-equivalent density,

$$\rho_\Lambda = \frac{\Lambda c^2}{8\pi G}, \quad p_\Lambda = -\rho_\Lambda c^2. \quad (10)$$

Metric compatibility makes (9) covariantly conserved for constant Λ .

An arbitrary dark-energy field or fluid is not equivalent to (9). It may remain within GR as part of S_m , but its equations, perturbations, and initial data then belong to a different matter model. A phenomenological relation $p_{\text{DE}} = w(a)\rho_{\text{DE}}c^2$ specifies a homogeneous background relation only; it need not determine sound speeds, anisotropic stress, or interactions. Those details cannot be replaced by the statement that the stress tensor is conserved.

The cosmological discussion uses the supernova evidence for accelerated expansion and a conventional Λ CDM benchmark (Perlmutter et al., 1999; Planck Collaboration, 2020; Riess et al., 1998). Here these are historical reference points. Neither an added Λ nor a chosen dark-energy stress tensor derives GR from $\mathcal{A}$ and Φ_{src} , or identifies the physical origin of vacuum energy.

5 Matter equations and conservation

Variation with respect to the matter variables supplies their own equations of motion. On a solution, the contracted Bianchi identity and metric compatibility give

$$\nabla^\mu T_{\mu\nu} = 0. \quad (11)$$

This is local covariant conservation of total nongravitational stress–energy. It does not by itself define a globally conserved total energy in every spacetime, nor does it provide a local tensor for gravitational energy. Global charges require suitable symmetries or boundary structure (Arnowitt et al., 1962; Carroll, 1997).

For components $T_{\mu\nu}^{(A)}$, interactions can exchange energy and momentum:

$$\nabla^\mu T_{\mu\nu}^{(A)} = Q_\nu^{(A)}, \quad \sum_A Q_\nu^{(A)} = 0. \quad (12)$$

Separate conservation is an additional no-exchange condition, not a consequence of total conservation alone. For example, a separately modeled dark-energy component must either satisfy that condition or supply its

exchange law.

For a perfect fluid comoving with a normalized four-velocity u , the convention used here is

$$T_{\mu\nu} = \left(\rho + \frac{p}{c^2}\right) u_\mu u_\nu + p g_{\mu\nu}, \quad u^\mu u_\mu = -c^2. \quad (13)$$

Its locally measured energy density is ρc^2 . This convention prevents the Friedmann and Raychaudhuri equations from mixing mass density with energy density.

6 Metric, trajectories, and measurements

The metric assigns a tangent vector X its causal type:

$$g(X, X) < 0 \text{ timelike}, \quad g(X, X) = 0 \text{ null}, \quad g(X, X) > 0 \text{ spacelike}, \quad (14)$$

with the zero vector excluded when classifying null directions. A time orientation selects the future timelike cone without supplying a preferred observer congruence.

An ideal clock following a timelike curve measures

$$d\tau^2 = -\frac{1}{c^2} g_{\mu\nu} dx^\mu dx^\nu. \quad (15)$$

The action $S_p = -mc^2 \int d\tau$ for a neutral structureless test particle gives

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} = 0. \quad (16)$$

This description neglects external forces, spin, finite size, and significant self-gravity. A charged particle or a supported observer need not follow a geodesic.

For standard Maxwell matter in vacuum, the leading geometric-optics limit gives

$$g_{\mu\nu} k^\mu k^\nu = 0, \quad k^\nu \nabla_\nu k^\mu = 0, \quad (17)$$

with an affine parametrization. Finite wavelengths, a material medium, or modified electromagnetic dynamics require a separate analysis. There is no second optical metric postulated here.

At an event p , a local inertial chart satisfies

$$g_{ab}(p) = \eta_{ab}, \quad \partial_c g_{ab}(p) = 0. \quad (18)$$

Curvature remains in the second derivatives. The local special-relativistic description is exact at the tangent-space level, while a finite laboratory has curvature corrections of relative order $\ell^2/\mathcal{R}^2$ when its characteristic size ℓ is small compared with a curvature radius $\mathcal{R}$.

7 Weak-field limit

Consider an isolated, slowly moving source with negligible pressure at leading order, no leading anisotropic stress, and weak nearly static fields. Let

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll 1. \quad (19)$$

This section uses $\Lambda = 0$, or a region where $|\Lambda|L^2$ is negligible relative to the weak Newtonian potential $\epsilon = \sup |\Phi|/c^2$. Boundary conditions select the usual decaying isolated-source solution.

The leading Einstein equation gives

$$\nabla^2 \Phi = 4\pi G \rho. \quad (20)$$

In an isotropic weak-field chart,

$$h_{00} = -\frac{2\Phi}{c^2}, \quad h_{ij} = -\frac{2\Phi}{c^2} \delta_{ij}, \quad (21)$$

at leading order. Thus the metric through first order is

$$ds^2 \simeq -\left(1 + \frac{2\Phi}{c^2}\right) c^2 dt^2 + \left(1 - \frac{2\Phi}{c^2}\right) \delta_{ij} dx^i dx^j. \quad (22)$$

The omitted dimensionless diagonal metric components are $O(\epsilon^2)$ in the static expansion; velocity-dependent components enter at their appropriate post-Newtonian orders. Comparing the spatial coefficient with $1 - 2\gamma\Phi/c^2$ gives

$$\gamma = 1. \quad (23)$$

This is the familiar GR parameter in this weak isolated-source regime, not an independent constraint obtained from Φ_{src} .

The leading connection is $\Gamma^i_{00} = \partial_i \Phi/c^2$. Since $dx^0/dt = c$, the slow-motion geodesic equation gives

$$\frac{d^2 x^i}{dt^2} = -\partial_i \Phi + O(c^2 \epsilon^2/L), \quad (24)$$

where $v^2/c^2 = O(\epsilon)$ and derivatives vary on scale L . The estimate has acceleration units. It summarizes the size of terms omitted in the stated quasi-static regime; it is not a substitute for a full post-Newtonian calculation when those terms are needed.

8 What the flow language adds, and what it does not

An observer congruence can be introduced after a solution (M, g, ψ) has been chosen. Its expansion, shear, vorticity, and acceleration describe the neighboring observer worldlines. These are useful ways to organize cosmological expansion, static support, and relative motion. Geometry develops them explicitly.

There is no separate variation with respect to that interpretive congruence in (2). If u is a matter-fluid velocity, its behavior follows from the specified matter model. If it denotes a family of measuring observers, its choice describes the measurement setup. Neither role supplies a new gravitational field equation or fixes the meaning of Φ_{src} .

No source evolution law is specified. For an action-based proposal, a future expression such as

$$S_{\text{src}}[q] = \int_{\mathcal{A}} \mathcal{L}_{\text{src}}[q] \quad (25)$$

would require a defined source configuration q and an integrable top-degree form or density $\mathcal{L}_{\text{src}}$, together with a domain, variations, and any boundary terms. None of these is adopted merely by displaying the schematic integral. In particular, a source measure or metric cannot be supplied without acknowledgment to make it look complete.

9 A non-circular dynamics-recovery task

The source research task is different from repeating the variation above. It must start from independently specified source states and laws, identify how an effective metric and matter variables are obtained, and show that their induced equations meet a declared GR target. An action-based construction may derive an effective action; a construction based on evolution equations or constraints may instead establish equivalent dynamics. Neither route is selected by the untyped Φ_{src} .

For an action-based route, the argument must track the measure, allowed variations, boundary terms, symmetry assumptions, and normalization. If eliminating source variables or taking a limit is involved, the regime and discarded terms must be stated. Terms that become small in a controlled limit cannot be called identically absent. A nonlinear completion is not established by agreement only at the quadratic or Newtonian level.

The same discipline applies to matter. A task may assume a specified matter sector and ask only for gravitational recovery, but it must not then claim to have derived universal matter coupling. Constants may be left as parameters if that is the stated target. The contribution lies in what the source argument actually fixes, not in renaming freely chosen inputs.

In a fixed-assumption AI audit, importing (2) as a new source premise invalidates a claim to have derived that action from the original source data. In open construction, the same addition can be recorded as a hypothesis, but it does not advance that particular explanatory step. A valid negative result would identify a contradiction with a required dynamical property or a counterexample to a precisely stated implication. Failure to find an action after a finite search is unresolved, not a no-go theorem.

10 Exact closure and remaining questions

The effective gravitational equations are fixed by the adopted action. Their equality to the corresponding GR equations is exact, not only weak-field. Observational equality further requires the same matter sector, constants, solution class, and experimental interpretation. A source-based selection rule or new matter coupling would need a fresh comparison.

Retaining this benchmark when a speculative extension fails is a reasonable comparison policy. It does not turn the source ontology into a verified theory or guarantee the consistency of later additions. Likewise, the standard conservation law and geodesic limit do not constrain undefined source degrees of freedom.

The central result of this paper is therefore a correctly normalized effective model and its representative limits. It supplies the equations needed for the companion analyses, while leaving the source origin of spacetime, gravity, matter, clocks, and the constants G and Λ unresolved. The source-research task is to justify an effective dynamics from separately stated source laws, with added assumptions and approximation errors exposed. Reproducing the benchmark variation is a control calculation; it is not evidence that this explanatory step has been completed.

References

- Arnowitt, R., Deser, S., & Misner, C. W. (1962). The dynamics of general relativity. In L. Witten (Ed.), *Gravitation: An introduction to current research* (pp. 227–265). Wiley. <https://arxiv.org/abs/gr-qc/0405109>
- Carroll, S. M. (1997). *Lecture notes on general relativity*. arXiv: gr-qc/9712019. <https://arxiv.org/abs/gr-qc/9712019>
- Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., Deustua, S., Fabbro, S., Goobar, A., Groom, D. E., Hook, I. M., Kim, A. G., Kim, M. Y., Lee, J. C., Nunes, N. J., Pain, R., Pennypacker, C. R.,

- Quimby, R., Lidman, C., . . . Couch, W. J. (1999). Measurements of Ω and Λ from 42 high-redshift supernovae. *The Astrophysical Journal*, 517, 565–586. <https://doi.org/10.1086/307221>
- Planck Collaboration. (2020). Planck 2018 results. VI. cosmological parameters. *Astronomy & Astrophysics*, 641, Article A6. <https://doi.org/10.1051/0004-6361/201833910>
- Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., Gilliland, R. L., Hogan, C. J., Jha, S., Kirshner, R. P., Leibundgut, B., Phillips, M. M., Reiss, D., Schmidt, B. P., Schommer, R. A., Smith, R. C., Spyromilio, J., Stubbs, C., Suntzeff, N. B., & Tonry, J. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116, 1009–1038. <https://doi.org/10.1086/300499>