

Æther-flow Ontology

and Relativistic Framework

Contents

1	Ontology: Foundations	1
1.1	The question and the present answer	1
1.2	Physical intent and intrinsic ordered motion	1
1.3	The source assumptions	2
1.4	Source arena and physical spacetime	3
1.5	What “experienced space” would need to mean	3
1.6	S-time: order, duration, and experience	3
1.7	The adopted relativistic benchmark	4
1.8	Expansion, light, and the limits of interpretation	5
1.9	How a later source proposal should be assessed	5
1.10	Research purpose and the assumption boundary	5
1.11	Recovery targets and negative conclusions	6
1.12	Conclusion	7
2	Effective Dynamics and Matter Coupling	8
2.1	What is being specified	8
2.2	Conventions and model specification	8
2.3	Action and bulk field equation	9
2.4	Cosmological constant and other dark-energy models	10
2.5	Matter equations and conservation	10
2.6	Metric, trajectories, and measurements	11
2.7	Weak-field limit	11
2.8	What the flow language adds, and what it does not	12
2.9	A non-circular dynamics-recovery task	13
2.10	Exact closure and remaining questions	13
3	Gravitational Degrees of Freedom and Perturbative Consistency	14
3.1	The consistency question	14
3.2	Action and admissible backgrounds	14
3.3	Linearization and gauge symmetry	15
3.4	Scalar, vector, and tensor perturbations	16
3.5	Hamiltonian degree-of-freedom count	16
3.6	Quadratic radiative action and its implications	17
3.7	What does not follow from the flat calculation	17
3.8	Source consistency and scoped obstruction tasks	18
3.9	Conclusion	19
4	Relativistic Recovery and Operational Predictions	20
4.1	Recovery as a statement about the effective model	20
4.2	Exact relation to the corresponding GR model	20
4.3	Weak-field gravity	21
4.4	Light and causal structure	22

4.5	Static gravitational redshift and clocks	22
4.6	Homogeneous cosmology	23
4.7	Local special relativity and spatial measurement	24
4.8	From comparison calculations to recovery evidence	25
4.9	Conclusion	25
5	Observer Congruences and the Geometry of Flow	27
5.1	Why a congruence is useful	27
5.2	Spacetime, normalization, and observer choice	27
5.3	Local rest space and relative velocity	28
5.4	Expansion, shear, vorticity, and acceleration	28
5.5	Raychaudhuri evolution	29
5.6	Cosmological expansion and defocusing	30
5.7	Static support and gravitational redshift	31
5.8	Vorticity and frame dragging are distinct	32
5.9	Curvature, tidal acceleration, and wave strain	33
5.10	Local special relativity	34
5.11	A geometry-first source milestone	34
5.12	Interpretive scope and conclusion	35
6	The Exact-GR Benchmark	36
6.1	Purpose and source assumptions	36
6.2	Definition of the effective benchmark	36
6.3	Meaning of exact closure	37
6.4	Clocks, light, and local relativity	37
6.5	Cosmological completion	38
6.6	The restricted weak-field result	38
6.7	Observer flow within GR	39
6.8	The open source derivation	39
6.9	Research outcomes and AI-assisted use	39
6.10	Benchmark policy and conclusion	40
7	Overview of the Æther-flow Framework	41
7.1	Two questions, two levels of description	41
7.2	Common effective statement	41
7.3	What each article contributes	42
7.4	Results and shared limits	43
7.5	Relation to other approaches	43
7.6	Use of the benchmark	43
7.7	A bounded feasibility sequence	44
7.8	Conclusion	44
8	Æther-flow as an Interpretation of General Relativity	45
8.1	The proposal and its central distinction	45
8.2	What is assumed about the source	45
8.3	A definite effective gravitational model	46
8.4	What an observer-based flow description can explain	48

8.5	Explanatory value and empirical limits	48
8.6	Relationship to other gravitational approaches	49
8.7	A dual physics and AI research objective	49
8.8	Task specification and the meaning of success	50
8.9	Proposed evaluation protocol	51
8.10	Initial feasibility study and stopping conditions	52
8.11	Conclusion	52
References		54

Ontology: Foundations

Abstract. The *Æther* / *Æther-flow* framework asks whether spatial and temporal relations can be recovered from intrinsic ordered motion in a proposed four-dimensional source structure, rather than assumed at the outset. Its present assumptions specify a smooth four-dimensional manifold but leave the mathematical type of the source flow unresolved. General relativity is adopted separately as the quantitative gravitational benchmark. This paper defines the source vocabulary, separates interpretive aims from mathematical results, and sets the conditions for assessing a future reconstruction. The research has two purposes: investigate a source explanation of relativity and evaluate whether an artificial-intelligence research system can produce independently checkable deductions or scoped negative results. A counterexample to entailment, an obstruction within a specified model class, and an unresolved search have different meanings. No source-to-spacetime derivation or general impossibility theorem is established here.

1.1 The question and the present answer

The *Æther* / *Æther-flow* framework is motivated by a question about physical explanation: could the spacetime relations described by relativity arise from a more basic organization, rather than being the starting point of the account? The words *Æther* and *Æther-flow* express that aim. They do not, by themselves, specify the objects or laws needed to answer it.

The project uses this source question as both a physics investigation and a proposed AI-assisted research case study. Its physical aim is a deeper account of GR, not a departure from its predictions introduced for its own sake. GR agreement is the recovery target against which source candidates are tested, not a guaranteed outcome. A valid argument against a specified candidate is therefore an acceptable research result.

The present framework takes a conservative position on the measurable physics. It adopts general relativity (GR), with a single Lorentzian metric and specified ordinary matter, as the effective gravitational model. The source proposal is kept separate. This separation allows familiar relativistic calculations to be used without implying that the proposed source already explains their origin.

There are consequently two different questions. One is whether the adopted effective model reproduces the observations that the same GR model reproduces. Conditional on identical matter, parameters, solutions, and measurement rules, it does. The other is whether the source assumptions produce that effective model. No construction establishing the second statement is supplied here. The distinction is substantive: agreement obtained by adopting a theory is not evidence that its equations have been derived from another theory.

This paper establishes the vocabulary and the scope of that distinction. Dynamics states the effective action; Consistency examines its local gravitational degrees of freedom; Relativistic Recovery derives representative measurement relations; Geometry explains the observer-congruence language used to interpret them. Those companion analyses concern GR unless a source-to-target connection is explicitly supplied.

1.2 Physical intent and intrinsic ordered motion

The working ontological hypothesis is that the *Æther* is a physical four-dimensional substrate and that *Æther-flow* is its intrinsic ordered motion. Here *substrate* means a proposed underlying physical reality, not an independently observed material. The intended explanation would connect its organization to the

spacetime geometry measured through clocks, distances, and light signals. The smooth manifold specified below supplies a mathematical starting point; it does not establish the physical existence of that substrate.

Observers are conceived as part of the same underlying reality, with access to local relations rather than a view from outside it. “Experienced space” names the proposed account of local three-dimensional spatial measurements. “S-time” names the intended connection between ordered change and temporal experience. These remain working hypotheses. Neither an embedded spatial slice nor a calibrated clock functional follows merely from giving them names.

The flow is not intended as a wind through an already existing three-dimensional space. Rather, the research asks whether intrinsic source relations can generate the metric, the structure that assigns spacetime intervals and causal directions. This reverses the explanatory order of ordinary motion described using a metric and a clock. A mathematical definition of source motion must therefore state its ordering or parameter structure without silently treating that structure as already measured physical time.

This distinction preserves the physical intent without treating it as an accomplished mechanism. Calling the source four-dimensional does not supply its laws, explain how observers are represented, or establish agreement with GR. Those are reconstruction requirements. The effective model below is the comparison target, not evidence that the proposed source already produces it.

It is also important not to confuse this use of Æther with Einstein-aether theory. The latter introduces a dynamical unit timelike vector with an action and additional coupled modes (Jacobson & Mattingly, 2004; Jacobson & Mattingly, 2001). The effective model here contains no corresponding independent gravitational vector. An observer’s four-velocity will appear in the geometric description, but choosing observers is not the same operation as adding a field equation for a new physical substance.

1.3 The source assumptions

Definition 1.1 (Source arena). *The symbol $\mathcal{A}$ denotes a smooth four-dimensional manifold. Its manifold structure and dimension are assumed. No particular topology is selected by this statement, and no Lorentzian metric, causal order, physical clock, or matter content is assigned to it.*

The source text describes these assumptions as *primitive debt*: they are structure used at the beginning, not structure recovered by a derivation. Here that phrase means only “assumed rather than explained.” A later proposal may justify or replace these assumptions, but they cannot currently be advertised as consequences of Æther-flow.

Definition 1.2 (Unspecified source structure). *Φ_{src} is a placeholder for a possible source-order or evolution structure. Its mathematical type has not been fixed. It is not yet a map, vector field, differential operator, stochastic process, group action, or physical-time parameter.*

The distinction between a placeholder and a mathematical flow matters. A conventional flow might be specified by maps Φ_λ obeying a composition law and defined on a parameter domain. Writing such maps here would add assumptions about that domain, composition, and perhaps invertibility. The source definition explicitly leaves those choices unresolved. We therefore retain Φ_{src} without assigning it a convenient but unsupported type.

The source package can be summarized as

$$\mathcal{A} \text{ smooth, } \dim \mathcal{A} = 4, \quad \Phi_{\text{src}} \text{ not yet typed.} \quad (1.1)$$

This is a statement of the current setup, not a closed source theory. It supplies no state space for evolving

degrees of freedom, no admissible variations, no dynamics, no observables, and no map to clocks or detectors. These omissions constrain what can be inferred; they do not establish that every future specification will fail.

1.4 Source arena and physical spacetime

Let M denote the effective spacetime manifold and $g_{\mu\nu}$ its metric. Writing (M, g) separately from $\mathcal{A}$ prevents an ambiguity present in informal descriptions of the ontology. Both arenas may be four-dimensional, but equal dimension does not identify them. Nor does smoothness supply a metric or a causal cone.

A source-to-target reconstruction would need to explain how some independently defined source data determine, approximate, or otherwise represent (M, g) and its operational content. Whether this requires a map between manifolds, a map between state spaces, a quotient, or a limiting construction depends on the source model. No one of these possibilities is adopted here.

It follows that a familiar GR congruence u^μ cannot simply be declared equal to Φ_{src} . The former is a well-defined vector field on an already Lorentzian spacetime; the latter has neither a type nor a target map. The geometric use of the word flow here is therefore an effective description of observer motion, not a construction of source motion.

A further distinction concerns reality claims. Selecting $\mathcal{A}$ as a mathematical arena does not establish that a physical substrate with that structure exists. Its proposed physical role is an ontological hypothesis. The effective metric, in contrast, already has an operational role within the adopted model: it enters the rules used to compare clocks, trajectories, and light signals. A future bridge must connect these roles rather than relying on the common use of geometric language.

1.5 What “experienced space” would need to mean

The phrase *experienced spatial slice* originally expressed the aim of explaining why observers describe local measurements using three spatial directions. GR already provides a precise local construction for that purpose. Given a future-directed four-velocity normalized by $u^\mu u_\mu = -c^2$, define

$$h_{\mu\nu} = g_{\mu\nu} + \frac{u_\mu u_\nu}{c^2}. \quad (1.2)$$

The projector $h^\mu{}_\nu$ selects vectors orthogonal to u^μ . Its image is a three-dimensional local rest space. This is an observer-dependent tangent-space statement, not a new source derivation (Ellis & van Elst, 1999).

Three different ideas must not be merged. A rest space specifies local spatial directions. A spacelike hypersurface organizes events into a spatial surface when the required integrability conditions hold. A visual scene is reconstructed from received signals and need not correspond to events simultaneous in that rest frame. The ontology has not shown that any of these is an embedded slice of $\mathcal{A}$.

For a smooth timelike congruence, vanishing vorticity gives local hypersurface orthogonality. A global slicing needs additional global conditions. The present framework does not assume that every observer family defines a global three-space, and it does not derive a source embedding $\Sigma \hookrightarrow \mathcal{A}$. A future account of experienced space must state which of these operational or geometric targets it aims to reconstruct.

1.6 S-time: order, duration, and experience

S-time is retained as the name of an interpretive aim: relating temporal experience to the organization of change. No S-time functional is defined by the present source package. In particular, no monotonicity law or

calibration identifies source order with measured duration.

Order and duration are different. An order may distinguish which events can precede others without assigning a number of seconds to an interval. In relativity, proper time assigns duration along a timelike trajectory using the metric. A thermodynamic arrow introduces further questions about irreversible behavior and boundary conditions. Subjective temporal experience raises questions not settled by the mere existence of a clock variable. The source manuscripts contain no derivation connecting all these notions.

The effective model therefore uses ordinary proper time. For a timelike worldline Γ ,

$$\tau[\Gamma] = \frac{1}{c} \int_{\Gamma} \sqrt{-g_{\mu\nu} dx^{\mu} dx^{\nu}}. \quad (1.3)$$

This operational prescription is adopted from GR (Carroll, 1997). Calling it an expression of S-time is an interpretation only until a source construction reproduces the same clock readings and their dependence on trajectory. No universal preferred clock or source time is inferred from the notation.

1.7 The adopted relativistic benchmark

1.7.1 Fields, action, and conventions

The effective arena is a time-oriented four-dimensional Lorentzian spacetime with signature $(-+++)$ and Levi-Civita connection. Greek indices run from 0 to 3. In the covariant action, coordinates are length-valued, with $x^0 = ct$; S_m has the units of physical action. The benchmark is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad (1.4)$$

where $G > 0$, Λ is constant, and ψ denotes specified matter fields. Taking $\Lambda = 0$ gives the original noncosmological specialization. Matter is coupled to the same metric, with ordinary locally Lorentz-invariant matter laws when the standard light and clock interpretations are used.

With the physical-action stress tensor

$$T_{\mu\nu} = -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}, \quad (1.5)$$

the bulk field equation is

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (1.6)$$

Dynamics gives the variation and its boundary assumptions. No source variable occurs in this action as a new propagating gravitational field.

1.7.2 Operational content and equivalence

The effective metric supplies the causal classification of tangent directions. Standard Maxwell light in vacuum geometric optics follows null directions,

$$g_{\mu\nu} k^{\mu} k^{\nu} = 0, \quad (1.7)$$

and ideal clocks measure (1.3). Neutral structureless test bodies follow timelike geodesics when nongravitational forces and finite-size effects can be neglected. These are model assumptions and standard consequences, not new laws induced by an already defined source flow.

The term *exact closure* names the decision to fix this effective gravitational model to GR. For a specified matter model, the two descriptions have identical observables only when they also use the same constants, admissible solutions, initial and boundary data, and measurement rules. If a source interpretation imposed an additional restriction on possible solutions, that restriction would have to enter the comparison. Equality of the printed metric equation alone would no longer suffice.

The action with unspecified S_m thus defines a family of possible GR models. It does not select one cosmological history or one laboratory outcome. Its purpose is to provide a reference against which a completed source model could be compared.

1.8 Expansion, light, and the limits of interpretation

The motivating ontology includes expansion, light propagation, and temporal change. In the adopted benchmark, cosmological expansion is described by a changing FLRW scale factor and the expansion of comoving worldlines. No external container is needed for that description. Light propagation and gravitational redshift follow the same metric and observer measurements developed in Relativistic Recovery.

These facts provide a vocabulary for the proposed interpretation, not evidence that Φ_{src} produces them. A comoving expansion scalar $\theta = 3H$, for example, is calculated after a spacetime metric and congruence have been supplied. It cannot serve as the missing source dynamics without reversing the intended explanatory direction.

The same caution applies to metaphor. Static support, rotational observer motion, and wave-induced deformation can all be described using congruences, but no single image of a vortex exhausts gravity. Geometry makes these distinctions quantitative. Its quantities depend on the chosen observers as well as the metric and are not separate manifestations of an independently established source field.

1.9 How a later source proposal should be assessed

A useful source development would first define the objects left open here. It would then state a particular reconstruction claim and show which source assumptions imply it. The claim might concern a causal structure, an operational clock, a metric, or a field equation; these are different levels of achievement. Recovering one does not automatically recover the others.

A successful bridge must avoid importing its intended conclusion as a disguised premise. Starting with a Lorentzian source metric and identifying it with g may define a model, but it would not derive Lorentzian geometry from the present metric-free source assumptions. Similarly, postulating the Einstein–Hilbert action under a different name is not a derivation of that action. Such choices can be legitimate assumptions if they are acknowledged as assumptions.

If an optional extension fails a consistency or observational test, the independent GR benchmark remains a reference. This is a research policy, not a theorem protecting all source hypotheses. A failure may provide evidence against the source assumptions responsible for it. Conversely, absence of a bridge in these manuscripts is not a proof that no bridge can be constructed.

1.10 Research purpose and the assumption boundary

The project seeks to determine whether a mathematically specified development of the ontology can recover GR, or whether a stated reconstruction claim can be ruled out within a defined scope. A negative result is an admissible scientific outcome. The project also provides a proposed research case study for artificial

intelligence (AI): can a system formulate precise questions, track its premises, and produce arguments that withstand independent checking? These objectives are related but must be assessed separately.

Sys4AI is the intended agentic research framework. In this role, it would coordinate AI agents, research tasks, records of assumptions, and verification work. That intended use does not supply a physical law or establish the capability of an implemented system. The manuscripts report no completed Sys4AI evaluation runs. The protocol in the synthesis article specifies evidence that such runs would need to produce.

A research task begins with a fixed version of the source assumptions. It must identify the mathematical type of every object used, the admissible source states or histories, and the rule connecting those objects to the proposed target. Smoothness and four-dimensionality remain assumed unless that task explicitly replaces them. Choosing a type for Φ_{src} , a topology, an evolution rule, a measure, or a metric is an addition to the present specification, not a deduction from the word flow.

There are two distinct modes of work. In a *fixed-assumption audit*, the agent may analyze the stated premises but may not change them to obtain the desired conclusion. In *open construction*, the agent may propose new definitions and physical assumptions. Each addition must be recorded, motivated, and included in the final theorem statement. Open construction is part of the research objective; it is not permission to describe a result obtained from strengthened premises as a deduction from the original ontology alone.

The first mathematical deliverable should therefore be a source specification and a precise recovery claim, together with a nonempty example or an appropriate existence argument. An inconsistent premise set cannot earn recovery credit through a vacuous implication. Where consistency or existence remains unproved, that dependency stays explicit. A conditional theorem can still be reported as conditional.

1.11 Recovery targets and negative conclusions

A recovery task must name its target before its outcome is assessed. A causal structure describes which events can influence others. A metric assigns spacetime intervals; an operational clock rule relates those intervals to measured duration. Gravitational dynamics specifies how geometry responds to matter. Matter coupling and measurement rules determine which quantities an experiment would observe. Progress on one target does not establish the others.

The task must also identify its domain. A local construction, a special family of solutions, an approximate limiting theory, and exact recovery of a specified GR solution class are different claims. A limited theorem can be valuable without being relabeled as full GR. A derivation of every matter field or the numerical values of every constant is not required unless those are included in the stated claim; externally supplied inputs must nevertheless remain visible.

Four outcomes need separate labels. A *verified derivation* establishes the declared target from the declared premises. A *counterexample to entailment* satisfies the premises but violates the target, showing that those premises do not force that conclusion. A *scoped obstruction* proves that no permitted reconstruction achieves the target within the specified model class and conditions. An *unresolved search* supplies neither the requested derivation nor a valid negative argument.

Non-entailment is weaker than incompatibility. A set of premises may admit a completion that reproduces GR and another that does not. The existence of the latter refutes a universal recovery claim, but not the existence of the former. Conversely, finding one GR-compatible completion does not show that the original ontology selects it uniquely. A proof excluding all admissible completions must quantify over the class actually defined, not over unspecified future meanings of Φ_{src} .

These distinctions set the present status. The source specification lacks the definitions needed for the

intended reconstruction; this paper does not supply a countermodel to a completed source theory or a theorem excluding every completion. Recognizing that gap is an audit finding. New research progress requires a further checkable result about an explicitly stated development of the proposal.

1.12 Conclusion

The present ontology consists of a selected smooth four-dimensional source arena, an untyped *Æther-flow* placeholder, and an interpretive aim concerning spatial and temporal organization. Its quantitative effective model is GR with specified ordinary matter. The geometric and operational results developed in the companion papers belong to that adopted model.

This division preserves the source question without overstating its answer. The existing assumptions do not establish a physical substrate, a source clock, or a derivation of Einsteinian dynamics. They define where further explanation is sought. A future result must supply the missing source structure and demonstrate a specific bridge; until then, exact agreement with GR is a property of the adopted benchmark rather than independent confirmation of the ontology. The research question remains open in a specific sense: the present source package is not a completed theory from which the target can be decided. The next useful result is a theorem or counterexample about a precisely defined source development. A valid scoped negative finding can advance that inquiry and the assessment of AI reasoning, without being a universal verdict on every possible completion.

Effective Dynamics and Matter Coupling

Abstract. The effective dynamics of the $\mathcal{A}$ ether-flow Interpretation of Relativity are specified by the Einstein–Hilbert action with ordinary matter and an optional cosmological constant. This action is adopted, not derived from the unresolved source structure. We fix the units of the action and stress tensor, derive the bulk field equation, distinguish total from separate component conservation, and state the assumptions behind particle, light, and clock descriptions. The weak-field calculation recovers Poisson’s equation and leading Newtonian acceleration in the stated regime. These results define a comparison target for source research. A separate recovery criterion requires independently specified source laws, an explicit reconstruction, and control of any approximation or added assumption. The present calculation establishes consequences of the general-relativistic action, not a source origin of that action or evidence of artificial-intelligence research performance.

2.1 What is being specified

An interpretation becomes a predictive physical model only after its fields, equations, and measurement rules have been fixed. In this framework, the effective gravitational choice is explicit: retain general relativity (GR), rather than introduce an additional low-energy $\mathcal{A}$ ether-flow field. The purpose of this paper is to state that choice in a form that can be varied, checked, and used in calculations.

The distinction from the source proposal is essential. Foundations uses $\mathcal{A}$ for a smooth four-dimensional source arena and Φ_{src} for an unresolved source-order or evolution structure. Neither is identified here with physical spacetime or an observer velocity. The effective fields instead live on a separate manifold M and are governed by the action below. No inference from the source assumptions to that action is claimed.

The project seeks a source construction that reproduces GR rather than changes its predictions. This paper supplies the independently adopted target dynamics for that investigation. It also explains the effective flow interpretation, but an interpretation of a chosen action is not a derivation of that action. If a source candidate fails to reproduce the target, retaining GR as a reference does not repair or validate that candidate.

2.2 Conventions and model specification

The spacetime (M, g) is a smooth, time-oriented four-dimensional Lorentzian manifold, with signature $(-+++)$ and Levi-Civita connection ∇ . Greek indices range over 0, 1, 2, 3. The curvature convention is

$$[\nabla_\mu, \nabla_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma, \quad R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}. \quad (2.1)$$

Coordinates in the covariant action have dimensions of length; in a time-adapted chart $x^0 = ct$. When a metric is instead written with t in seconds, the displayed dt and factors of c implement that coordinate change. Tensor equations do not depend on the chosen chart.

The matter fields are denoted collectively by ψ . Their action must be specified for an actual prediction. “Ordinary matter” here means a matter model with the appropriate locally Lorentz-invariant laws and standard coupling to g ; it does not mean every arbitrary Lagrangian containing a metric. Stability, characteristics, and interactions of a chosen matter sector require their own assessment.

The intended source account would recover this shared metric coupling rather than add a direct force between matter and an Æther-flow wind. Its absence from the adopted action is explicit; its absence from a future source reduction must be demonstrated. Coupling matter only to one metric is not, by itself, a proof that the full theory has no observable preferred structure. Source-dependent gravitational dynamics or restrictions on solutions could still change the comparison. The reconstruction must therefore account for the metric dynamics and admissible solutions as well as the matter coupling.

The free constants include $G > 0$ and a constant Λ . Initial and boundary conditions, topology, and the domain of the solution must also be supplied for a particular physical problem. They are not selected by the phrase exact closure.

2.3 Action and bulk field equation

The gravitational benchmark is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi]. \quad (2.2)$$

Setting $\Lambda = 0$ recovers the Einstein–Hilbert form

$$S_{\text{eff}}|_{\Lambda=0} = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} R + S_m[g, \psi]. \quad (2.3)$$

Both S_m and the gravitational term have units of action. Since d^4x has dimensions L^4 and R has dimensions L^{-2} in this coordinate convention, the factor c^3/G provides the required normalization.

Define the physical stress–energy tensor by

$$T_{\mu\nu} := -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}. \quad (2.4)$$

The factor of c is required when $x^0 = ct$ and S_m is a physical action. An equivalent convention could rescale the matter action, but then its variational definition would need to say so. With (2.4),

$$\delta S_m = -\frac{1}{2c} \int_M d^4x \sqrt{-g} T_{\mu\nu} \delta g^{\mu\nu} \quad (2.5)$$

on the matter equations, apart from boundary terms associated with the chosen variations.

For compactly supported metric variations, the bulk variation of the full action is

$$\delta S_{\text{eff}} = \int_M d^4x \sqrt{-g} \left[\frac{c^3}{16\pi G} (G_{\mu\nu} + \Lambda g_{\mu\nu}) - \frac{T_{\mu\nu}}{2c} \right] \delta g^{\mu\nu}. \quad (2.6)$$

Stationarity for arbitrary such variations gives

$$\boxed{G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}}. \quad (2.7)$$

For $\Lambda = 0$, this becomes

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (2.8)$$

The calculation is the standard metric variation of GR, with the unit convention made explicit (Carroll, 1997). It establishes what follows from (2.2), not why a source ontology should produce that action.

A finite-boundary problem needs an appropriate boundary action and boundary data. The Einstein–Hilbert integrand alone contains boundary variations involving derivatives of δg . Compact support avoids that issue for the bulk derivation; it is not a substitute for specifying boundary terms when the physical problem requires them.

2.4 Cosmological constant and other dark-energy models

A cosmological constant can be kept on the geometric side of (2.7), or moved to the matter side as the specific tensor

$$T_{\mu\nu}^{(\Lambda)} = -\frac{\Lambda c^4}{8\pi G} g_{\mu\nu}. \quad (2.9)$$

These are exactly equivalent descriptions of the same constant term, not two independent sources to be counted together. In perfect-fluid language, with ρ denoting mass-equivalent density,

$$\rho_\Lambda = \frac{\Lambda c^2}{8\pi G}, \quad p_\Lambda = -\rho_\Lambda c^2. \quad (2.10)$$

Metric compatibility makes (2.9) covariantly conserved for constant Λ .

An arbitrary dark-energy field or fluid is not equivalent to (2.9). It may remain within GR as part of S_m , but its equations, perturbations, and initial data then belong to a different matter model. A phenomenological relation $p_{\text{DE}} = w(a)\rho_{\text{DE}}c^2$ specifies a homogeneous background relation only; it need not determine sound speeds, anisotropic stress, or interactions. Those details cannot be replaced by the statement that the stress tensor is conserved.

The cosmological discussion uses the supernova evidence for accelerated expansion and a conventional Λ CDM benchmark (Perlmutter et al., 1999; Planck Collaboration, 2020; Riess et al., 1998). Here these are historical reference points. Neither an added Λ nor a chosen dark-energy stress tensor derives GR from $\mathcal{A}$ and Φ_{src} , or identifies the physical origin of vacuum energy.

2.5 Matter equations and conservation

Variation with respect to the matter variables supplies their own equations of motion. On a solution, the contracted Bianchi identity and metric compatibility give

$$\nabla^\mu T_{\mu\nu} = 0. \quad (2.11)$$

This is local covariant conservation of total nongravitational stress–energy. It does not by itself define a globally conserved total energy in every spacetime, nor does it provide a local tensor for gravitational energy. Global charges require suitable symmetries or boundary structure (Arnowitt et al., 1962; Carroll, 1997).

For components $T_{\mu\nu}^{(A)}$, interactions can exchange energy and momentum:

$$\nabla^\mu T_{\mu\nu}^{(A)} = Q_\nu^{(A)}, \quad \sum_A Q_\nu^{(A)} = 0. \quad (2.12)$$

Separate conservation is an additional no-exchange condition, not a consequence of total conservation alone. For example, a separately modeled dark-energy component must either satisfy that condition or supply its

exchange law.

For a perfect fluid comoving with a normalized four-velocity u , the convention used here is

$$T_{\mu\nu} = \left(\rho + \frac{p}{c^2}\right) u_\mu u_\nu + p g_{\mu\nu}, \quad u^\mu u_\mu = -c^2. \quad (2.13)$$

Its locally measured energy density is ρc^2 . This convention prevents the Friedmann and Raychaudhuri equations from mixing mass density with energy density.

2.6 Metric, trajectories, and measurements

The metric assigns a tangent vector X its causal type:

$$g(X, X) < 0 \text{ timelike}, \quad g(X, X) = 0 \text{ null}, \quad g(X, X) > 0 \text{ spacelike}, \quad (2.14)$$

with the zero vector excluded when classifying null directions. A time orientation selects the future timelike cone without supplying a preferred observer congruence.

An ideal clock following a timelike curve measures

$$d\tau^2 = -\frac{1}{c^2} g_{\mu\nu} dx^\mu dx^\nu. \quad (2.15)$$

The action $S_p = -mc^2 \int d\tau$ for a neutral structureless test particle gives

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} = 0. \quad (2.16)$$

This description neglects external forces, spin, finite size, and significant self-gravity. A charged particle or a supported observer need not follow a geodesic.

For standard Maxwell matter in vacuum, the leading geometric-optics limit gives

$$g_{\mu\nu} k^\mu k^\nu = 0, \quad k^\nu \nabla_\nu k^\mu = 0, \quad (2.17)$$

with an affine parametrization. Finite wavelengths, a material medium, or modified electromagnetic dynamics require a separate analysis. There is no second optical metric postulated here.

At an event p , a local inertial chart satisfies

$$g_{ab}(p) = \eta_{ab}, \quad \partial_c g_{ab}(p) = 0. \quad (2.18)$$

Curvature remains in the second derivatives. The local special-relativistic description is exact at the tangent-space level, while a finite laboratory has curvature corrections of relative order $\ell^2/\mathcal{R}^2$ when its characteristic size ℓ is small compared with a curvature radius $\mathcal{R}$.

2.7 Weak-field limit

Consider an isolated, slowly moving source with negligible pressure at leading order, no leading anisotropic stress, and weak nearly static fields. Let

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll 1. \quad (2.19)$$

This section uses $\Lambda = 0$, or a region where $|\Lambda|L^2$ is negligible relative to the weak Newtonian potential $\epsilon = \sup |\Phi|/c^2$. Boundary conditions select the usual decaying isolated-source solution.

The leading Einstein equation gives

$$\nabla^2 \Phi = 4\pi G \rho. \quad (2.20)$$

In an isotropic weak-field chart,

$$h_{00} = -\frac{2\Phi}{c^2}, \quad h_{ij} = -\frac{2\Phi}{c^2} \delta_{ij}, \quad (2.21)$$

at leading order. Thus the metric through first order is

$$ds^2 \simeq -\left(1 + \frac{2\Phi}{c^2}\right) c^2 dt^2 + \left(1 - \frac{2\Phi}{c^2}\right) \delta_{ij} dx^i dx^j. \quad (2.22)$$

The omitted dimensionless diagonal metric components are $O(\epsilon^2)$ in the static expansion; velocity-dependent components enter at their appropriate post-Newtonian orders. Comparing the spatial coefficient with $1 - 2\gamma\Phi/c^2$ gives

$$\gamma = 1. \quad (2.23)$$

This is the familiar GR parameter in this weak isolated-source regime, not an independent constraint obtained from Φ_{src} .

The leading connection is $\Gamma^i_{00} = \partial_i \Phi/c^2$. Since $dx^0/dt = c$, the slow-motion geodesic equation gives

$$\frac{d^2 x^i}{dt^2} = -\partial_i \Phi + O(c^2 \epsilon^2/L), \quad (2.24)$$

where $v^2/c^2 = O(\epsilon)$ and derivatives vary on scale L . The estimate has acceleration units. It summarizes the size of terms omitted in the stated quasi-static regime; it is not a substitute for a full post-Newtonian calculation when those terms are needed.

2.8 What the flow language adds, and what it does not

An observer congruence can be introduced after a solution (M, g, ψ) has been chosen. Its expansion, shear, vorticity, and acceleration describe the neighboring observer worldlines. These are useful ways to organize cosmological expansion, static support, and relative motion. Geometry develops them explicitly.

There is no separate variation with respect to that interpretive congruence in (2.2). If u is a matter-fluid velocity, its behavior follows from the specified matter model. If it denotes a family of measuring observers, its choice describes the measurement setup. Neither role supplies a new gravitational field equation or fixes the meaning of Φ_{src} .

No source evolution law is specified. For an action-based proposal, a future expression such as

$$S_{\text{src}}[q] = \int_{\mathcal{A}} \mathcal{L}_{\text{src}}[q] \quad (2.25)$$

would require a defined source configuration q and an integrable top-degree form or density $\mathcal{L}_{\text{src}}$, together with a domain, variations, and any boundary terms. None of these is adopted merely by displaying the schematic integral. In particular, a source measure or metric cannot be supplied without acknowledgment to make it look complete.

2.9 A non-circular dynamics-recovery task

The source research task is different from repeating the variation above. It must start from independently specified source states and laws, identify how an effective metric and matter variables are obtained, and show that their induced equations meet a declared GR target. An action-based construction may derive an effective action; a construction based on evolution equations or constraints may instead establish equivalent dynamics. Neither route is selected by the untyped Φ_{src} .

For an action-based route, the argument must track the measure, allowed variations, boundary terms, symmetry assumptions, and normalization. If eliminating source variables or taking a limit is involved, the regime and discarded terms must be stated. Terms that become small in a controlled limit cannot be called identically absent. A nonlinear completion is not established by agreement only at the quadratic or Newtonian level.

The same discipline applies to matter. A task may assume a specified matter sector and ask only for gravitational recovery, but it must not then claim to have derived universal matter coupling. Constants may be left as parameters if that is the stated target. The contribution lies in what the source argument actually fixes, not in renaming freely chosen inputs.

In a fixed-assumption AI audit, importing (2.2) as a new source premise invalidates a claim to have derived that action from the original source data. In open construction, the same addition can be recorded as a hypothesis, but it does not advance that particular explanatory step. A valid negative result would identify a contradiction with a required dynamical property or a counterexample to a precisely stated implication. Failure to find an action after a finite search is unresolved, not a no-go theorem.

2.10 Exact closure and remaining questions

The effective gravitational equations are fixed by the adopted action. Their equality to the corresponding GR equations is exact, not only weak-field. Observational equality further requires the same matter sector, constants, solution class, and experimental interpretation. A source-based selection rule or new matter coupling would need a fresh comparison.

Retaining this benchmark when a speculative extension fails is a reasonable comparison policy. It does not turn the source ontology into a verified theory or guarantee the consistency of later additions. Likewise, the standard conservation law and geodesic limit do not constrain undefined source degrees of freedom.

The central result of this paper is therefore a correctly normalized effective model and its representative limits. It supplies the equations needed for the companion analyses, while leaving the source origin of spacetime, gravity, matter, clocks, and the constants G and Λ unresolved. The source-research task is to justify an effective dynamics from separately stated source laws, with added assumptions and approximation errors exposed. Reproducing the benchmark variation is a control calculation; it is not evidence that this explanatory step has been completed.

Gravitational Degrees of Freedom and Perturbative Consistency

Abstract. The effective gravitational sector of the $\mathcal{A}$ ether-flow Interpretation of Relativity is general relativity. We display its Ricci-flat linearized equation, the scalar–vector–tensor decomposition on Minkowski spacetime, and the Hamiltonian count of two local gravitational configuration degrees of freedom. For radiative vacuum perturbations about Minkowski spacetime, the reduced transverse-traceless action has positive quadratic energy for positive Newton’s constant and a massless dispersion relation. These are restricted checks of the adopted metric theory, not stability results for every background or for an untyped source proposal. The paper also specifies how a future source-consistency task should distinguish a verified counterexample, a class-wide obstruction, and an unsuccessful search. Those criteria support evaluation of artificial-intelligence-assisted reasoning without treating a negative physical result as a failure of valid reasoning.

3.1 The consistency question

A claim that a gravitational theory is “healthy” can mean several different things. It may mean that no physical mode has a negative kinetic term, that the equations admit a well-posed local evolution problem, that a particular background has no growing perturbations, or that solutions avoid singularities. These are not interchangeable claims. This paper identifies which of them follow from the effective model adopted here.

The adopted gravitational action is the Einstein–Hilbert action. No independent vector, scalar, or other $\mathcal{A}$ ether-flow gravitational field is added. Consequently, the following calculation is a check of the standard effective GR sector rather than a new source-level consistency proof. The source objects $\mathcal{A}$ and Φ_{src} are not yet defined by the state space, action, or evolution law needed for such a proof.

The main calculation concerns vacuum radiative perturbations about Minkowski spacetime. A more general Ricci-flat background is included to display how curvature enters the linearized equation. Matter and cosmological backgrounds are discussed as separate cases whose stability cannot be inferred from the flat calculation alone.

3.2 Action and admissible backgrounds

We use signature $(- + + +)$, length-valued covariant coordinates with $x^0 = ct$, and

$$[\nabla_\mu, \nabla_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma. \quad (3.1)$$

The effective action is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad (3.2)$$

where $G > 0$ and the physical-action stress tensor is $T_{\mu\nu} = -(2c/\sqrt{-g})\delta S_m/\delta g^{\mu\nu}$. Its field equation is

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (3.3)$$

A perturbation background must solve this equation together with the background matter equations. An arbitrary smooth metric is not automatically an admissible background for a fixed matter model.

For the explicit flat-vacuum calculation, take $\Lambda = 0$, no background matter, and $\bar{g}_{\mu\nu} = \eta_{\mu\nu}$. For the displayed curved vacuum equation, take $\bar{R}_{\mu\nu} = 0$. Initial-value discussions are restricted to suitable globally hyperbolic regions with appropriate data. This is an assumption about the evolution problem, not a claim that every GR spacetime has a global Cauchy surface.

3.3 Linearization and gauge symmetry

Write

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}, \quad \psi = \bar{\psi} + \delta\psi, \quad (3.4)$$

and retain terms linear in the perturbations. The general metric equation becomes

$$\delta G_{\mu\nu}[h] + \Lambda h_{\mu\nu} = \frac{8\pi G}{c^4} \delta T_{\mu\nu}[h, \delta\psi]. \quad (3.5)$$

The linearized matter equations must accompany it whenever the background contains matter.

With an active infinitesimal diffeomorphism convention, the gauge transformation is

$$h_{\mu\nu} \mapsto h_{\mu\nu} + 2\bar{\nabla}_{(\mu}\xi_{\nu)}, \quad \delta\psi \mapsto \delta\psi + \mathcal{L}_\xi \bar{\psi}. \quad (3.6)$$

The sign would reverse under the opposite coordinate convention, but all transformed quantities must use the same convention. Gauge redundancy expresses different mathematical descriptions of the same perturbative geometry and matter configuration. It is not a new physical degree of freedom.

Define the trace and trace reversal by

$$h = \bar{g}^{\mu\nu} h_{\mu\nu}, \quad \tilde{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2} \bar{g}_{\mu\nu} h. \quad (3.7)$$

In de Donder gauge,

$$\bar{\nabla}^\mu \tilde{h}_{\mu\nu} = 0, \quad (3.8)$$

the vacuum equation on a Ricci-flat background is

$$\bar{\nabla}^\rho \bar{\nabla}_\rho \tilde{h}_{\mu\nu} + 2\bar{R}_{\mu\rho\nu\sigma} \tilde{h}^{\rho\sigma} = 0. \quad (3.9)$$

The curvature term is important: local wave propagation does not imply that perturbations on every curved background behave like flat plane waves. Equation (3.9) is not the unchanged perturbation equation for a general matter or nonzero- Λ background (Carroll, 1997; Gourgoulhon, 2007).

For Minkowski vacuum, it reduces to

$$\square \tilde{h}_{\mu\nu} = 0, \quad \square = -\frac{1}{c^2} \partial_t^2 + \nabla^2. \quad (3.10)$$

Residual gauge transformations preserving de Donder gauge satisfy $\square \xi_\mu = 0$ in this flat case. The metric has ten symmetric components, but neither those components nor the four gauge conditions alone are a count of physical radiative modes. The constraints and remaining gauge freedom must also be taken into account.

3.4 Scalar, vector, and tensor perturbations

On Minkowski spacetime, decompose the spatial dependence using the Euclidean metric δ_{ij} :

$$h_{00} = -2\phi, \quad h_{0i} = S_i + \partial_i B, \quad (3.11)$$

$$h_{ij} = h_{ij}^{\text{TT}} + \partial_{(i} F_{j)} + \left(\partial_i \partial_j - \frac{1}{3} \delta_{ij} \nabla^2 \right) E + \frac{1}{3} \delta_{ij} H. \quad (3.12)$$

The fields obey

$$\partial^i S_i = \partial^i F_i = 0, \quad \partial^i h_{ij}^{\text{TT}} = 0, \quad \delta^{ij} h_{ij}^{\text{TT}} = 0. \quad (3.13)$$

Parentheses denote symmetrization with weight 1/2. The quantities ϕ, B, E, H are scalar components of this decomposition, not the source variable Φ_{src} or the Newtonian potential Φ used elsewhere. Their units follow from the derivatives in (3.11)–(3.12).

The decomposition requires appropriate spatial boundary conditions or a Fourier description; global zero modes and boundary degrees of freedom need separate treatment. In the radiative vacuum sector, the constraints and gauge freedom permit transverse-traceless (TT) gauge:

$$h_{00} = 0, \quad h_{0i} = 0, \quad \partial^i h_{ij} = 0, \quad \delta^{ij} h_{ij} = 0. \quad (3.14)$$

For a wave moving in the z direction, the independent components can be written

$$h_{ij}^{\text{TT}} = \begin{pmatrix} h_+ & h_\times & 0 \\ h_\times & -h_+ & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (3.15)$$

The two functions are the two usual gravitational-wave polarizations. No third amplitude is supplied by Φ_{src} , because that source placeholder does not enter the effective action as a field.

This radiative statement should not be misread as saying that all scalar or vector metric perturbations are physically meaningless. In a sourced problem they can describe constrained gravitational fields and contribute to observables. Cosmological matter perturbations also have their own physical degrees of freedom. Absence of an additional vacuum scalar graviton is a narrower claim.

3.5 Hamiltonian degree-of-freedom count

The Arnowitt–Deser–Misner (ADM) formulation separates the spatial metric, lapse, and shift (Arnowitt et al., 1962). On the ordinary regular local constraint surface, the six components of the spatial metric γ_{ij} and their six conjugate momenta π^{ij} provide twelve phase-space variables per point. The lapse and shift enforce the Hamiltonian and three momentum constraints rather than supplying propagating metric modes.

Each of these four first-class constraints removes one phase-space variable through the constraint and one through gauge equivalence. The reduction is therefore

$$2 \times 4 = 8, \quad (3.16)$$

leaving

$$12 - 8 = 4 \quad (3.17)$$

local physical phase-space dimensions, or

$$4/2 = 2 \quad (3.18)$$

local gravitational configuration degrees of freedom. This count agrees with the TT wave analysis.

The count concerns the gravitational bulk sector. It does not count independent matter modes, global moduli, or boundary observables, and it presumes the usual nondegenerate metric and regular constraint structure. It is also not a stability theorem: two physical modes can evolve stably or unstably depending on the background and problem.

3.6 Quadratic radiative action and its implications

For weak vacuum perturbations about Minkowski spacetime, after the constraints and gauge have been handled and suitable boundary terms discarded, the TT action is

$$S_{\text{TT}}^{(2)} = \frac{c^3}{64\pi G} \int d^4x \left[\frac{1}{c^2} \dot{h}_{ij}^{\text{TT}} \dot{h}_{ij}^{\text{TT}} - \partial_k h_{ij}^{\text{TT}} \partial_k h_{ij}^{\text{TT}} \right], \quad (3.19)$$

where the dot denotes ∂_t and $d^4x = c dt d^3x$. Equivalently, the coefficient of $\dot{h}^2 - c^2(\nabla h)^2$ in the time–space integral is $c^2/(64\pi G)$.

The associated quadratic energy is

$$H_{\text{TT}}^{(2)} = \frac{c^2}{64\pi G} \int d^3x \left[\dot{h}_{ij}^{\text{TT}} \dot{h}_{ij}^{\text{TT}} + c^2 \partial_k h_{ij}^{\text{TT}} \partial_k h_{ij}^{\text{TT}} \right]. \quad (3.20)$$

For $G > 0$, both contributions are nonnegative. This establishes absence of a negative-kinetic-energy radiative graviton in this flat, reduced quadratic sector. It is not a positive local gravitational-energy tensor on an arbitrary spacetime.

Variation gives $\ddot{h}_{ij}^{\text{TT}} - c^2 \nabla^2 h_{ij}^{\text{TT}} = 0$. A Fourier mode proportional to $e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}$ therefore satisfies

$$\omega^2 = c^2 |\mathbf{k}|^2. \quad (3.21)$$

There is no mass term, no tachyonic mass in this dispersion relation, and no wrong-sign spatial-gradient term. Finite-wavenumber vacuum plane waves are oscillatory. Possible zero modes and boundary conditions remain part of the particular perturbation problem.

3.7 What does not follow from the flat calculation

3.7.1 Local evolution and causal structure

After a suitable gauge choice, the Einstein equations have a hyperbolic evolution formulation, subject to constraints and suitable matter equations (Gourgoulhon, 2007). Its principal part controls local propagation and the initial-value problem. Establishing this structure is not the same as proving that every allowed perturbation remains small indefinitely. A hyperbolic equation can have growing solutions, and a locally well-posed evolution can develop singular behavior.

On curved backgrounds, lower-order curvature terms, the matter sector, boundaries, and global geometry matter. Stability must be attached to a specified background and a specified class of perturbations. The Minkowski calculation does not establish nonlinear stability, global existence, or the absence of caustics and

spacetime singularities for all GR solutions.

3.7.2 No additional preferred-structure sector

The absence of an independent Æther-flow field in (3.2) means that this action introduces no additional gravitational vector or scalar mode. That differs from a theory containing a dynamical aether vector, for which extra coupled modes must be studied (Jacobson & Mattingly, 2004). Merely writing a normalized observer velocity in a measurement description does not alter the count.

This statement does not automatically extend to a future source theory. Such a theory might reconstruct only the existing modes, introduce additional ones, or fail to define consistent evolution. Since its dynamical variables are not fixed here, none of those outcomes has been calculated.

The intended recovery keeps the effective gravitational content of the stated GR model, with no additional observable force field or second operative metric. This is a requirement on a source construction, not an inferred property of Φ_{src} . Any additional source variables must have their physical role and effective consequences assessed. A candidate that requires an observable extra mode in the claimed exact-GR regime has not met that target; the two-mode count of the independently adopted GR action cannot resolve the failure.

3.8 Source consistency and scoped obstruction tasks

A source-consistency task must identify its mathematical variables, allowed configurations, evolution or constraint laws, and proposed reconstruction. It must also name the property being checked. Absence of a negative kinetic term, a well-posed initial-value problem, and agreement with a target mode count are separate properties. The effective GR results above cannot be transferred to source variables that have not been defined.

A negative result needs a witness or a proof with a stated scope. For a counterexample to a universal recovery claim, the witness must satisfy the source assumptions and fail the required conclusion under the specified reconstruction. One failed choice of reconstruction does not exclude every other permitted map. A class-wide obstruction must address all maps allowed by the task, or state the narrower subclass it actually excludes.

A useful diagnostic example would be a proposed source model whose laws force every reconstructed metric to be flat. If the declared target includes curved GR solutions, a proof of that restriction would rule out full recovery for that model and reconstruction class. This is a hypothetical example of the required logic, not a property proved for the present ontology. A model restricted to flat spacetime might still meet a narrower target that explicitly asks only for flat-space kinematics.

The evaluation record must separate physical and reasoning outcomes. A valid obstruction is a negative answer for the specified source proposal and a successful piece of reasoning. An invalid obstruction, an unsupported assertion of consistency, and an unsuccessful search are different outcomes. Exhausting a computation budget does not prove that a reconstruction is impossible.

Checking should be independent of the original argument where practical. Algebraic manipulations may be checked with symbolic software; a proof assistant may verify a formal statement; another researcher may inspect the assumptions and physical interpretation. Each check must identify its scope. A machine-checked theorem about an inaccurate formalization does not settle the intended physics, and a second agent's agreement alone is not a proof. No source obstruction or source-consistency result is claimed by these methodological requirements.

3.9 Conclusion

The adopted effective gravitational sector has the standard local GR gauge structure and two radiative gravitational degrees of freedom. In the stated Minkowski vacuum regime, its reduced quadratic action has positive kinetic energy for $G > 0$ and the massless dispersion relation $\omega^2 = c^2|\mathbf{k}|^2$. The framework adds no propagating Æther-flow gravitational field to that action.

These are useful bounded consistency results. They do not establish the stability of every relativistic background, the health of arbitrary matter, or the consistency of the unresolved source ontology. A future claim in any of those directions needs the corresponding model and argument rather than an extension of the flat-vacuum conclusion by wording alone. For source research, the appropriate outcome is a checked statement about a declared model and property. A verified obstruction can count as successful reasoning; a finite unsuccessful search must remain unresolved. Neither outcome is supplied by the flat effective-sector calculation alone.

Relativistic Recovery and Operational Predictions

Abstract. The effective $\mathcal{A}$ ether-flow model adopts general relativity, so its predictions agree with the corresponding relativistic model when matter, constants, allowed solutions, and measurement procedures match. This paper develops representative calculations of Newtonian acceleration, vacuum light propagation, gravitational redshift, proper time, and homogeneous cosmology. Each result is attached to its approximation and operational assumptions. These calculations provide comparison targets for a future source theory; they do not construct a metric or clock from the ontology. A recovery claim must distinguish agreement for a selected solution from recovery of a declared solution class, and approximate agreement from exact equivalence. The resulting task specification permits checkable partial progress while preventing finite tests or inherited calculations from being presented as a full source derivation.

4.1 Recovery as a statement about the effective model

There are two meanings of recovering relativity. A proposed modification might approach GR in a limit. A source theory might also produce a relativistic spacetime and its equations from other variables. Neither is demonstrated merely by adopting the Einstein–Hilbert action. Here, “recovery” refers to obtaining familiar operational results from an effective model already fixed to GR.

This is still worth making explicit. A field equation is not the whole content of a prediction: one must choose a matter model, a solution, and a procedure for comparing clocks or signals. The calculations below make those choices visible in simple regimes. They also provide concrete targets for the open source program.

The source arena $\mathcal{A}$ is not identified with physical spacetime M , and the placeholder Φ_{src} remains untyped. No source clock, causal order, or reconstruction map is used in the following derivations. The physical interpretation of the source proposal must therefore remain separate from the established meaning of the effective observables.

The motivating physical picture treats observers as local participants in a deeper ordered source. Their distances and elapsed times are the intended outputs of a reconstruction, not measurements by an observer outside that source. To make this picture operational, a candidate must represent the measuring systems and recover the specified clock and signal comparisons. Calling spatial measurements a local slice or duration S-time does not yet provide that construction.

4.2 Exact relation to the corresponding GR model

Use signature $(-+++)$ and length-valued coordinates $x^0 = ct$ in covariant action integrals. The effective action and equation are

$$S_{\text{eff}} = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad (4.1)$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad T_{\mu\nu} = -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}. \quad (4.2)$$

The matter action uses the same operative metric. Its fields and interactions must be specified, with standard locally Lorentz-invariant matter assumed for the usual clock and light interpretation.

Let $\mathcal{D}$ denote matching constants, initial and boundary data, and a corresponding physical solution; let $\mathcal{P}$ denote the same measurement procedure. Then

$$O_{\text{Æther-flow}}(\mathcal{D}, \mathcal{P}) = O_{\text{GR}}(\mathcal{D}, \mathcal{P}) \quad (4.3)$$

provided the effective solution class and observable definitions also coincide and the source interpretation introduces no additional restriction or coupling. The equality follows because the two sides evaluate the same effective model. It is not a new empirical fit or a proof of source emergence.

An unspecified S_m leaves a family of models rather than one prediction. Likewise, a hidden condition selecting different solutions would have to be included in (4.3); equality of the Einstein equations alone would not establish the stated equivalence.

4.3 Weak-field gravity

4.3.1 Newtonian potential and spatial curvature

For an isolated slowly moving source, assume negligible leading pressure and anisotropic stress, weak quasi-static fields, and boundary conditions selecting the usual isolated-source solution. In this section $\Lambda = 0$, or its contribution is negligible on the region's scale L . Write

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll 1. \quad (4.4)$$

The leading field equation gives

$$\nabla^2 \Phi = 4\pi G \rho, \quad (4.5)$$

where ρ is the mass density and Φ has dimensions of velocity squared. In an isotropic chart, the metric through first order is

$$ds^2 \simeq - \left(1 + \frac{2\Phi}{c^2}\right) c^2 dt^2 + \left(1 - \frac{2\gamma\Phi}{c^2}\right) \delta_{ij} dx^i dx^j, \quad (4.6)$$

with

$$\gamma = 1. \quad (4.7)$$

The parameter γ measures the coefficient of spatial curvature relative to the Newtonian potential in this weak-field parametrization. Its value here is inherited from GR, not computed from an independent source-flow law (Carroll, 1997).

For a localized positive mass M , the exterior potential is $\Phi = -GM/r$ at leading order. Both the temporal and spatial parts of (4.6) matter for light propagation. A model that matched only the slow-motion acceleration would not thereby have shown the same light deflection or travel-time delay. In the present benchmark those comparisons agree because the full effective metric and light law agree.

4.3.2 Slow test-particle motion

The leading connection in length-valued time coordinates is $\Gamma^i_{00} = \partial_i \Phi / c^2$. The geodesic equation for a neutral structureless test particle gives

$$\frac{d^2 x^i}{dt^2} = -\partial_i \Phi + O(c^2 \epsilon^2 / L), \quad (4.8)$$

where $\epsilon = \sup |\Phi|/c^2 \ll 1$, $v^2/c^2 = O(\epsilon)$, and fields vary on spatial scale L . The remainder is an acceleration estimate under those assumptions, not a complete post-Newtonian expression. The diagonal metric corrections omitted in the static approximation are dimensionless $O(\epsilon^2)$ terms.

The acceleration is attractive for the isolated potential just specified. A static supported observer instead needs a proper acceleration approximately $+\nabla\Phi$. Distinguishing support from free fall is important when interpreting the lapse gradient as a flow-related quantity.

4.4 Light and causal structure

For standard Maxwell theory in vacuum, the geometric-optics wave covector is locally the gradient of a phase. At leading order it satisfies

$$g^{\mu\nu} k_\mu k_\nu = 0. \quad (4.9)$$

Because $\nabla_\mu k_\nu = \nabla_\nu k_\mu$, differentiation of $k_\mu k^\mu = 0$ gives

$$k^\nu \nabla_\nu k^\mu = 0 \quad (4.10)$$

with the natural affine parametrization. Thus the same metric determines the vacuum light cone and its ray trajectories. These statements are restricted to the specified geometric-optics regime, not arbitrary propagation through a dispersive medium.

At a point p , local inertial coordinates satisfy

$$g_{ab}(p) = \eta_{ab}, \quad \partial_c g_{ab}(p) = 0. \quad (4.11)$$

In the associated local orthonormal frame the null condition is

$$-c^2 dt_{\text{loc}}^2 + d\ell_{\text{loc}}^2 = 0, \quad (4.12)$$

so the measured vacuum light speed is

$$\frac{d\ell_{\text{loc}}}{dt_{\text{loc}}} = c. \quad (4.13)$$

A coordinate speed in an extended curved chart can differ from c . That is not a second local light speed or evidence for propagation through a material Æther wind. Over a finite region of scale ℓ , curvature corrections are controlled by $\ell^2/\mathcal{R}^2$ when that ratio is small; they cannot generally be removed throughout the region by changing coordinates. Making the region smaller reduces these finite-size corrections; it does not assert that the curvature tensor itself vanishes. The local special-relativistic statement concerns the tangent-space metric, not a source mechanism that switches curvature off.

4.5 Static gravitational redshift and clocks

Consider a static region with positive lapse $N(\mathbf{x})$:

$$ds^2 = -N(\mathbf{x})^2 c^2 dt^2 + \gamma_{ij}(\mathbf{x}) dx^i dx^j. \quad (4.14)$$

The coordinates here use t in seconds. For an observer fixed at $\mathbf{x}$,

$$d\tau = N(\mathbf{x})dt, \quad u = N^{-1}\partial_t. \quad (4.15)$$

The Killing vector $K = \partial_t$ generates the time-translation symmetry. Along a null geodesic, $-k_\mu K^\mu$ is conserved. The frequency measured by a static observer is proportional to $-k_\mu u^\mu$, and therefore to $1/N$. For a signal emitted at E and received at R ,

$$\frac{\nu_R}{\nu_E} = \frac{N_E}{N_R}. \quad (4.16)$$

The formula compares specified static observers and the same freely propagating signal. Moving emitters or receivers introduce additional Doppler factors.

In the weak field,

$$N = 1 + \frac{\Phi}{c^2} + O(\epsilon^2), \quad (4.17)$$

which gives

$$\frac{\nu_R - \nu_E}{\nu_E} = \frac{\Phi_E - \Phi_R}{c^2} + O(\epsilon^2). \quad (4.18)$$

Light climbing from a lower potential is redshifted relative to the receiver's clock. This result follows from static symmetry and local frequency measurement, not from a separately specified source clock.

More generally, an ideal clock records

$$\tau[\Gamma] = \frac{1}{c} \int_{\Gamma} \sqrt{-g_{\mu\nu} dx^\mu dx^\nu}. \quad (4.19)$$

Comparing different worldlines can therefore produce different elapsed times. That path dependence is part of the adopted relativistic model. The current ontology has not derived a source functional whose calibrated value reproduces (4.19).

4.6 Homogeneous cosmology

4.6.1 Geometry and expansion

For the homogeneous and isotropic benchmark, use the Friedmann–Lemaître–Robertson–Walker (FLRW) metric

$$ds^2 = -c^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right]. \quad (4.20)$$

Here a has dimensions of length, r is dimensionless, $k = 0, \pm 1$, and $d\Omega^2$ is the unit-sphere metric. Cosmic time t is proper time along the comoving congruence. Define

$$H = \frac{\dot{a}}{a}. \quad (4.21)$$

For $u = \partial_t$, the congruence expansion is

$$\theta = \nabla_\mu u^\mu = 3H. \quad (4.22)$$

The result measures a fractional volume-change rate: a comoving volume scales as a^3 . It does not require expansion into an external physical space.

4.6.2 Friedmann equations and acceleration

Let ρ and p denote the total mass-equivalent density and pressure of non- Λ matter in a comoving perfect-fluid description. Einstein's equation gives

$$H^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}, \quad (4.23)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{c^2}\right) + \frac{\Lambda c^2}{3}. \quad (4.24)$$

The continuity equation is

$$\dot{\rho} + 3H\left(\rho + \frac{p}{c^2}\right) = 0. \quad (4.25)$$

These equations, with an equation of state and initial data, specify the homogeneous model (Carroll, 1997; Ellis & van Elst, 1999).

Positive Λ contributes toward acceleration, but net $\ddot{a} > 0$ requires the positive term to exceed the total matter focusing term. Equivalently, moving Λ into the fluid side gives a vacuum component with $p_\Lambda = -\rho_\Lambda c^2$. It must not be counted on both sides. A component with sufficiently negative pressure contributes toward defocusing; it need not dominate the total at every epoch.

One may instead specify a dynamical dark-energy model with

$$p_{\text{DE}} = w(a)\rho_{\text{DE}}c^2. \quad (4.26)$$

That is not generally equivalent to constant Λ . The background relation must be supplemented by conservation or exchange equations and, for perturbative predictions, a matter model or suitable perturbation closure. An arbitrary conserved tensor is not a complete physical specification.

Supernova observations and Planck results motivate the accelerated cosmological models considered here (Perlmutter et al., 1999; Planck Collaboration, 2020; Riess et al., 1998). The present calculations inherit those models' equations; they contain no new fit and provide no evidence that dark energy originates in Φ_{src} .

4.6.3 Cosmological redshift

For comoving emission and reception in FLRW, radial null propagation and the comparison of successive wave crests give

$$1 + z = \frac{\nu_E}{\nu_R} = \frac{a(t_R)}{a(t_E)}. \quad (4.27)$$

This redshift has a different geometric setting from the static-lapse result. Both are evaluations of frequency relative to specified observers along a null signal. Their common operational basis is the metric and matter law, not an uncalibrated source-order parameter.

4.7 Local special relativity and spatial measurement

In a local inertial frame, a massive clock with measured speed $v < c$ satisfies

$$d\tau = dt_{\text{loc}}\sqrt{1 - v^2/c^2}. \quad (4.28)$$

The tangent-space Lorentz symmetry and the local null cone remain the usual ones. Introducing an observer family for interpretation does not designate it as the preferred family for the laws of physics. A future source model that selected an observable preferred frame would need to demonstrate how its consequences satisfy or depart from the intended benchmark.

Relative to a normalized observer u , local spatial projections use

$$h_{\mu\nu} = g_{\mu\nu} + \frac{u_\mu u_\nu}{c^2}. \quad (4.29)$$

This projector supplies a three-dimensional rest space at each event. It does not automatically define a global simultaneous space, and it is not a derivation of an experiential slice inside $\mathcal{A}$. The term S-time likewise remains an interpretive target. The measured duration appearing in this paper is ordinary τ , with no additional source-time law assumed.

4.8 From comparison calculations to recovery evidence

A source-recovery task should specify its output in terms of the physical structures used in these calculations. Recovering a metric with a null cone is narrower than recovering Maxwell propagation on that cone. Assigning a parameter to source histories is narrower than recovering proper time for ideal clocks. Reproducing Newtonian acceleration is narrower than reproducing the same weak-field spatial geometry, redshift, and light propagation. The omitted structures remain explicit dependencies.

For a selected source model, the reconstruction must identify which source states correspond to which effective states, how source transformations affect the physical description, and how the claimed observables are obtained. A full equivalence claim over a declared domain requires more than mapping some source solutions into GR solutions. It must explain whether the target GR solutions in that domain are represented and whether unwanted physical alternatives or additional restrictions remain. Redundant descriptions may be identified when their equivalence has been established.

Approximate recovery needs a stated comparison, a control parameter or scale, and an error estimate for the quantities claimed to agree. Exact recovery requires the corresponding statement without an unaccounted approximation. Agreement on a finite collection of numerical examples is evidence for those examples, not a proof covering all allowed initial data or all spacetime geometries.

These distinctions also determine AI credit. Reproducing the calculations in this paper is a useful control task for technical competence. Merely restating them or identifying their already documented source gap is not a new source result. A research contribution must add a verifiable implication, construction, or counterexample beyond the supplied exposition. An independently checked partial target can be reported as partial success; it must not be scored as full GR recovery.

4.9 Conclusion

The effective framework reproduces the stated weak-field, optical, clock, and cosmological relations because it uses the same relativistic equations and measurement rules as the corresponding GR model. The calculations show the role of pressure assumptions, observer motion, symmetry, length scales, and matter specification; those conditions are part of each result.

They also clarify the remaining task. A source theory would have to reproduce not only a tensor equation but the operational structures that turn its solutions into predictions. No such source-to-metric, source-to-matter, or source-to-clock map has been established by adopting GR. The present benchmark provides explicit

comparison targets without conflating them with a completed derivation. A future recovery claim should therefore be reported at the level actually established: a selected example, a controlled approximation, a declared solution class, or a full operational equivalence within a stated domain. The proposed AI assessment uses the same boundary between demonstrated and intended results.

Observer Congruences and the Geometry of Flow

Abstract. A timelike observer congruence gives flow language a precise meaning within general relativity. Its expansion, shear, vorticity, and acceleration describe different aspects of neighboring observer motion, not an independently established underlying substance. We develop the rest-space decomposition and Raychaudhuri equation, apply them to homogeneous expansion and static support, and show why frame dragging need not coincide with congruence vorticity. Tidal curvature, deformation rate, and wave strain are distinguished. These effective geometric results define possible reconstruction targets for source research. A geometry-first task must specify its source data and map, distinguish local from global claims, and account for every imported metric or observer structure. Neither the source variable nor its relation to spacetime is determined by the congruence calculations.

5.1 Why a congruence is useful

Flow language can express several different physical ideas. A fluid can expand, a family of supported observers can accelerate, neighboring freely falling bodies can acquire relative motion, and a rotating family can fail to define orthogonal spatial hypersurfaces. General relativity (GR) distinguishes these situations mathematically. Treating all of them as a literal swirl would obscure those distinctions.

A *congruence* is a smooth family of nonintersecting worldlines on a region of spacetime. A timelike congruence describes a family of observers, or the motion of a suitable continuous matter model. Its geometric quantities give a disciplined effective meaning to the word flow. This is standard relativistic kinematics (Ellis & van Elst, 1999; Raychaudhuri, 1955); the present use does not introduce an additional gravitational field.

The source hypothesis seeks a physical origin for the geometry accessible to local observers. Foundations assumes a smooth source manifold $\mathcal{A}$ and leaves Φ_{src} undefined. No source-to-metric or source-to-congruence map is given. The calculations below supply a geometric dictionary within GR, not a derivation from the source.

5.2 Spacetime, normalization, and observer choice

Use signature $(- + + +)$ and the Levi-Civita connection. Our curvature convention is

$$[\nabla_\mu, \nabla_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma, \quad R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}. \quad (5.1)$$

The congruence is represented by a future-directed four-velocity

$$u^\mu = \frac{dx^\mu}{d\tau}, \quad g_{\mu\nu}u^\mu u^\nu = -c^2. \quad (5.2)$$

In length-valued coordinates, u has velocity dimensions. When an example uses t in seconds, its time component is correspondingly $u^t = dt/d\tau$; this does not change the normalization.

The effective field equation is

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (5.3)$$

The kinematic identities developed first do not require (5.3). That equation enters when curvature is related to matter or a particular gravitational solution is chosen.

There is no unique congruence assigned to every metric. Comoving cosmological observers, static supported observers, and freely falling detectors are different choices made for different questions. If a matter model supplies a velocity, that velocity has a physical interpretation within the model; it is not automatically a universally preferred frame for all laws. Kinematic scalars are coordinate-invariant once u is fixed, but generally change when the observer family changes.

5.3 Local rest space and relative velocity

The rest-space metric and projector are

$$h_{\mu\nu} = g_{\mu\nu} + \frac{u_\mu u_\nu}{c^2}, \quad h^\mu{}_\nu = \delta^\mu{}_\nu + \frac{u^\mu u_\nu}{c^2}. \quad (5.4)$$

Normalization gives

$$h_{\mu\nu} u^\nu = 0, \quad h^\mu{}_\alpha h^\alpha{}_\nu = h^\mu{}_\nu, \quad h^\mu{}_\mu = 3. \quad (5.5)$$

Thus h selects the three-dimensional positive-definite subspace orthogonal to u at each event. It is not, without further conditions, the metric of a global family of spatial slices.

Every tangent vector decomposes as

$$X^\mu = -\frac{u_\nu X^\nu}{c^2} u^\mu + h^\mu{}_\nu X^\nu. \quad (5.6)$$

The first term is its temporal component relative to the observer; the second is spatial. For another *normalized future-directed four-velocity* w , this can be written

$$w^\mu = \Gamma(u^\mu + v^\mu), \quad u_\mu v^\mu = 0. \quad (5.7)$$

Using $w^2 = -c^2$ gives

$$\Gamma = -\frac{u_\mu w^\mu}{c^2} = \frac{1}{\sqrt{1 - v^2/c^2}}, \quad v^2 = h_{\mu\nu} v^\mu v^\nu. \quad (5.8)$$

The normalization of w is necessary; an arbitrary rescaled timelike tangent does not give the same Lorentz factor. Equations (5.7)–(5.8) are the usual local special-relativistic relative-velocity relations, expressed covariantly.

5.4 Expansion, shear, vorticity, and acceleration

For a scalar f , spatial differentiation is

$$D_\mu f = h_\mu{}^\alpha \nabla_\alpha f. \quad (5.9)$$

For a spatial covector X_ν , every free index must be projected:

$$D_\mu X_\nu = h_\mu{}^\alpha h_\nu{}^\beta \nabla_\alpha X_\beta. \quad (5.10)$$

The same rule applies to higher-rank tensors. Projecting only the derivative index need not leave the other indices spatial.

The observer acceleration and expansion scalar are

$$a_\mu = u^\nu \nabla_\nu u_\mu, \quad (5.11)$$

$$\theta = \nabla_\mu u^\mu. \quad (5.12)$$

The projected derivative $B_{\mu\nu} = h_\mu^\alpha h_\nu^\beta \nabla_\alpha u_\beta$ splits into its trace, symmetric trace-free part, and antisymmetric part:

$$\sigma_{\mu\nu} = B_{(\mu\nu)} - \frac{1}{3}\theta h_{\mu\nu}, \quad (5.13)$$

$$\omega_{\mu\nu} = B_{[\mu\nu]}. \quad (5.14)$$

Symmetrization and antisymmetrization each include the factor 1/2. These definitions imply

$$a_\mu u^\mu = 0, \quad \sigma_{\mu\nu} u^\nu = \omega_{\mu\nu} u^\nu = 0, \quad \sigma^\mu{}_\mu = 0. \quad (5.15)$$

The full derivative is therefore

$$\boxed{\nabla_\mu u_\nu = \frac{1}{3}\theta h_{\mu\nu} + \sigma_{\mu\nu} + \omega_{\mu\nu} - \frac{u_\mu a_\nu}{c^2}.} \quad (5.16)$$

The acceleration term accounts for the part not contained in the spatial derivative. Contracting the first index with u^μ recovers a_ν , which checks its sign and normalization.

Expansion describes fractional local volume change; shear describes anisotropic deformation rate; vorticity describes local twisting of the observer directions; acceleration measures departure from geodesic motion. In a length-valued orthonormal frame, θ , σ , and ω have dimensions T^{-1} , while a has dimensions L/T^2 . These quantities describe (g, u) together. They are not four independent gravitational substances, nor an invariant partition of all gravitational phenomena.

5.5 Raychaudhuri evolution

5.5.1 Derivation with the stated normalization

The evolution of θ follows from differentiating the acceleration and using the Ricci identity:

$$\nabla_\mu a^\mu = (\nabla_\mu u^\nu)(\nabla_\nu u^\mu) + u^\nu \nabla_\mu \nabla_\nu u^\mu \quad (5.17)$$

$$= (\nabla_\mu u^\nu)(\nabla_\nu u^\mu) + u^\nu \nabla_\nu \theta + R_{\mu\nu} u^\mu u^\nu. \quad (5.18)$$

The contraction of (5.16) gives

$$(\nabla_\mu u^\nu)(\nabla_\nu u^\mu) = \frac{1}{3}\theta^2 + \sigma_{\mu\nu}\sigma^{\mu\nu} - \omega_{\mu\nu}\omega^{\mu\nu}. \quad (5.19)$$

The acceleration cross terms vanish by orthogonality. Solving for the derivative of θ yields

$$\boxed{u^\mu \nabla_\mu \theta = -\frac{1}{3}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} + \nabla_\mu a^\mu - R_{\mu\nu} u^\mu u^\nu.} \quad (5.20)$$

This is the Raychaudhuri identity for the definitions used here (Ellis & van Elst, 1999; Raychaudhuri, 1955). Every term has dimensions T^{-2} . There is no additional factor c^{-2} on the Ricci contraction because the four-velocity already has norm $-c^2$.

The acceleration term is a *full spacetime divergence*. An alternative form uses

$$\nabla_\mu a^\mu = D_\mu a^\mu + \frac{a_\mu a^\mu}{c^2}. \quad (5.21)$$

Using this projected divergence introduces the positive acceleration-square term. Adding it while retaining the full divergence would double count the contribution.

5.5.2 Focusing and the meaning of rotation

For geodesic observers $a = 0$. For an irrotational geodesic congruence, the vorticity term also vanishes. If $R_{\mu\nu}u^\mu u^\nu \geq 0$, (5.20) gives $\dot{\theta} \leq -\theta^2/3$. This is a conditional focusing statement, not a universal claim that every observer congruence must contract.

Frobenius provides the local integrability criterion

$$\omega_{\mu\nu} = 0 \iff u_{[\mu} \nabla_{\nu]} u_\rho = 0 \iff \text{local hypersurface orthogonality.} \quad (5.22)$$

The result assumes a smooth nonvanishing timelike field. It does not, by itself, supply a global foliation. Nonzero vorticity obstructs orthogonal local slices for that observer family, not the existence of all possible spatial slicings of the spacetime.

5.6 Cosmological expansion and defocusing

For a small bundle of neighboring worldlines, let V denote the infinitesimal volume measured in the local rest space. Its fractional evolution is

$$\frac{1}{V} \frac{dV}{d\tau} = \theta. \quad (5.23)$$

The statement concerns local measured volume and requires no embedding in a larger container.

For comoving observers in a Friedmann–Lemaître–Robertson–Walker (FLRW) spacetime,

$$ds^2 = -c^2 dt^2 + a(t)^2 d\Sigma_k^2, \quad u = \partial_t, \quad (5.24)$$

where a is a length and $d\Sigma_k^2$ is a dimensionless constant-curvature spatial metric. The determinant gives

$$\theta = \frac{1}{\sqrt{-g}} \partial_t \sqrt{-g} = 3 \frac{\dot{a}}{a} = 3H, \quad a_\mu = \sigma_{\mu\nu} = \omega_{\mu\nu} = 0. \quad (5.25)$$

Thus an expanding cosmology need not involve vorticity. It is the volume-changing part of this particular congruence's motion.

For a comoving perfect fluid with non- Λ mass-equivalent density ρ and pressure p , Einstein's equation implies

$$R_{\mu\nu}u^\mu u^\nu = 4\pi G \left(\rho + \frac{3p}{c^2} \right) - \Lambda c^2. \quad (5.26)$$

Substituting into (5.20) gives

$$\dot{\theta} = -\frac{1}{3}\theta^2 - 4\pi G \left(\rho + \frac{3p}{c^2} \right) + \Lambda c^2, \quad (5.27)$$

and therefore

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2} \right) + \frac{\Lambda c^2}{3}. \quad (5.28)$$

The same result is obtained from the Friedmann equations in Relativistic Recovery. Positive Λ opposes focusing, but produces net acceleration only when the total right-hand side is positive. A negative-pressure component contributes similarly only according to its actual stress tensor and matter dynamics.

A constant Λ can be represented by $T_{\mu\nu}^{(\Lambda)} = -\Lambda c^4 g_{\mu\nu} / (8\pi G)$. A general dark-energy fluid is not interchangeable with this tensor. The geometric calculation here does not identify either component with Φ_{src} or derive its magnitude.

5.7 Static support and gravitational redshift

5.7.1 The lapse gradient is a proper acceleration

For a static region with $N > 0$,

$$ds^2 = -N(\mathbf{x})^2 c^2 dt^2 + \gamma_{ij}(\mathbf{x}) dx^i dx^j, \quad (5.29)$$

the fixed-position observers have

$$u = N^{-1} \partial_t. \quad (5.30)$$

Their spatial metric is time-independent and they are hypersurface normal, so

$$\theta = 0, \quad \sigma_{\mu\nu} = 0, \quad \omega_{\mu\nu} = 0. \quad (5.31)$$

Nevertheless, direct evaluation of $u^\nu \nabla_\nu u_\mu$ gives

$$a_\mu = c^2 D_\mu \ln N. \quad (5.32)$$

This is already the observer's physical four-acceleration; it must not be multiplied by another c^2 . The $c = 1$ lapse relation is common in the 3 + 1 literature (Gourgoulhon, 2010).

In a weak field with $\epsilon = \sup |\Phi|/c^2 \ll 1$ and variation scale L ,

$$N = 1 + \frac{\Phi}{c^2} + O(\epsilon^2), \quad (5.33)$$

so

$$a_i = \partial_i \Phi + O(c^2 \epsilon^2 / L). \quad (5.34)$$

For the isolated Newtonian potential $\Phi = -GM/r$, this is outward support, while the leading free-fall acceleration is inward. The locally measured magnitude is

$$a_{\text{proper}} = \sqrt{\gamma^{ij} a_i a_j}. \quad (5.35)$$

Static observers therefore illustrate why gravity cannot be identified with vorticity or expansion. They can have neither while living in curved spacetime.

5.7.2 An explicit check of Raychaudhuri

Take the diagnostic metric

$$ds^2 = -c^2 N(x)^2 dt^2 + dx^2 + dy^2 + dz^2. \quad (5.36)$$

Its static congruence has

$$a^x = c^2 \frac{N'}{N}, \quad \nabla_\mu a^\mu = c^2 \frac{N''}{N}, \quad R_{\mu\nu} u^\mu u^\nu = c^2 \frac{N''}{N}. \quad (5.37)$$

The two terms cancel in (5.20), as required by $\theta = \sigma = \omega = 0$. For a lapse linear in x , the curvature vanishes but the static observers can still accelerate; for a nonlinear lapse, the divergence and curvature terms need not vanish separately. This separates observer acceleration from spacetime curvature and checks the normalization without appealing to a source model.

5.7.3 Clock comparison

Static proper time satisfies

$$d\tau = N dt. \quad (5.38)$$

For a freely propagating null signal, the Killing energy associated with ∂_t is constant. Its locally measured frequency is proportional to $1/N$, so

$$\frac{\nu_R}{\nu_E} = \frac{N_E}{N_R}, \quad (5.39)$$

for static emitter E and receiver R . In the weak field,

$$\frac{\nu_R - \nu_E}{\nu_E} = \frac{\Phi_E - \Phi_R}{c^2} + O(\epsilon^2). \quad (5.40)$$

The lapse therefore enters both support acceleration and static clock comparison. That common geometric role is a valid interpretive connection. It does not show that a source flow has produced either effect.

5.8 Vorticity and frame dragging are distinct

The terms rotation, twist, and frame dragging should not be used as synonyms. Vorticity is a property of a chosen congruence. Frame-dragging measurements concern relations between specified observers, gyroscopes, and the geometry of a rotating gravitating system. The choice of reference frame and, often, asymptotic conditions matter.

A concrete example is a circular stationary-axisymmetric metric in a region where the following slicing is valid:

$$ds^2 = -N^2 c^2 dt^2 + \gamma_{\phi\phi} (d\phi - \Omega_Z dt)^2 + \gamma_{AB} dx^A dx^B. \quad (5.41)$$

Here A, B label the two meridional coordinates, $N > 0$, and all coefficients are independent of t and ϕ . Consider

$$u = N^{-1} (\partial_t + \Omega_Z \partial_\phi). \quad (5.42)$$

The corresponding covector is

$$u_b = -N c^2 dt, \quad u_\phi = 0, \quad \Omega_Z = -\frac{g_{t\phi}}{g_{\phi\phi}}. \quad (5.43)$$

These are zero-angular-momentum observers (ZAMOs). Because $u_b \wedge du_b = 0$, their vorticity vanishes. They are the locally nonrotating, hypersurface-normal observers described in Gourgoulhon (2010). Their coordinate angular velocity Ω_Z can nonetheless be nonzero in a rotating gravitational field relative to an appropriate asymptotic reference.

Thus frame dragging cannot be universally identified with nonzero $\omega_{\mu\nu}$. An off-diagonal metric component alone is also insufficient: rotating coordinates in flat spacetime can create one. Conversely, a physically rotating family in Minkowski spacetime can have nonzero vorticity without any gravitational curvature. A proposed flow account must state which observers and which observable are involved rather than assigning all rotational phenomena to one tensor.

5.9 Curvature, tidal acceleration, and wave strain

5.9.1 The electric Weyl tensor

Define the electric part of the Weyl tensor relative to u by

$$E_{\mu\nu} = \frac{1}{c^2} C_{\mu\alpha\nu\beta} u^\alpha u^\beta. \quad (5.44)$$

It is spatial, symmetric, and trace-free by the algebraic Weyl identities, whether or not matter is present. Its orthonormal components have dimensions L^{-2} . It describes the Weyl part of the tidal field seen by these observers, not every contribution to geodesic deviation.

For neighboring freely falling worldlines with an infinitesimal connecting vector ξ commuting with their common geodesic tangent, geodesic deviation is

$$\frac{D^2 \xi^\mu}{D\tau^2} = -R^\mu{}_{\alpha\nu\beta} u^\alpha \xi^\nu u^\beta. \quad (5.45)$$

The sign follows the stated curvature convention and index ordering. In a Ricci-flat region, $R = C$, so for spatial ξ ,

$$\boxed{\frac{D^2 \xi^\mu}{D\tau^2} = -c^2 E^\mu{}_\nu \xi^\nu.} \quad (5.46)$$

The factor of c^2 converts curvature times separation into acceleration (Carroll, 1997). In a weak static vacuum field, $E_{ij} \approx \partial_i \partial_j \Phi / c^2$, which reproduces the Newtonian relative acceleration $-\partial_i \partial_j \Phi \xi^j$.

Vacuum with $\Lambda \neq 0$ is not Ricci-flat. For an Einstein space $R_{\mu\nu} = \Lambda g_{\mu\nu}$, the spatial tidal equation becomes

$$\frac{D^2 \xi^\mu}{D\tau^2} = -c^2 E^\mu{}_\nu \xi^\nu + \frac{\Lambda c^2}{3} \xi^\mu. \quad (5.47)$$

De Sitter spacetime has zero Weyl tensor but nonzero isotropic relative acceleration. Matter supplies further Ricci contributions. These examples show why an expression involving only $E_{\mu\nu}$ needs a stated regime.

5.9.2 Shear, strain, and a weak gravitational wave

Shear is a rate of anisotropic deformation, not the curvature tensor itself. The distinction becomes explicit in a weak vacuum gravitational wave written in transverse–traceless (TT) coordinates:

$$ds^2 = -c^2 dt^2 + [\delta_{ij} + h_{ij}^{\text{TT}}(t, \mathbf{x})] dx^i dx^j, \quad \partial^i h_{ij}^{\text{TT}} = 0, \quad \delta^{ij} h_{ij}^{\text{TT}} = 0. \quad (5.48)$$

Let $\varepsilon_h \ll 1$ bound the dimensionless wave amplitude, let T be its temporal variation scale, and assume the corresponding derivative bounds. The estimates below refer to that fixed scale, with $|\xi|$ the local spatial separation. To first order, the fixed-spatial-coordinate worldlines define a freely falling congruence $u = \partial_t$. Its projected velocity gradient is $B_{ij} = \frac{1}{2}\partial_t g_{ij}$. Hence

$$\sigma_{ij} = \frac{1}{2}\dot{h}_{ij}^{\text{TT}} + O(\varepsilon_h^2/T), \quad \theta = O(\varepsilon_h^2/T). \quad (5.49)$$

The curvature calculation gives

$$E_{ij} = -\frac{1}{2c^2}\ddot{h}_{ij}^{\text{TT}} + O(\varepsilon_h^2/(c^2T^2)), \quad (5.50)$$

so a small spatial separation obeys

$$\frac{D^2 \xi^i}{D\tau^2} = \frac{1}{2}\ddot{h}_{ij}^{\text{TT}}\xi^j + O(\varepsilon_h^2|\xi|/T^2), \quad (5.51)$$

at leading order about the central observer. The dimensionless metric perturbation represents strain, its first time derivative gives shear in this congruence, and its second derivative gives tidal acceleration per separation.

These relations explain the connection without identifying the quantities. Arbitrary shear need not indicate radiation, and static congruences can have zero shear despite nonzero tidal curvature. The weak-wave example assumes linear perturbations, the specified observer family, and separations small enough for the local geodesic-deviation description.

5.10 Local special relativity

At an event p , one can choose a local inertial frame with

$$g_{ab}(p) = \eta_{ab}, \quad \partial_c g_{ab}(p) = 0, \quad (5.52)$$

$$u^a(p) = (c, 0, 0, 0). \quad (5.53)$$

The local line element is consequently

$$ds^2 = -c^2 dt_{\text{loc}}^2 + dx_{\text{loc}}^2 + dy_{\text{loc}}^2 + dz_{\text{loc}}^2 \quad (5.54)$$

at the tangent-space level. Relative velocities obey (5.8). A local inertial frame does not remove curvature. Reducing laboratory size reduces finite-region corrections, not curvature itself. Nothing here selects a physical *Æther* rest frame.

5.11 A geometry-first source milestone

The observer construction suggests a limited first research target: specify source data and determine whether they support a well-defined local causal geometry or clock relation. This target does not require a simultaneous derivation of Einstein's field equation. It does require an explicit map from the independently specified source objects to the proposed geometric object, together with a statement of which features are assumed and which follow.

If a candidate constructs a metric, its dimension, regularity, nondegeneracy, and signature must be established on the claimed domain. A source metric that is assumed to be Lorentzian cannot then be counted

as a derivation of Lorentzian signature. If only causal directions are recovered, the remaining interval scale and calibration of physical clocks must be addressed separately. If an observer field is selected, its role must be stated rather than inferred from the name *Æther-flow*.

Local and global claims also differ. The rest-space projector above is local; a global family of spatial hypersurfaces requires additional conditions. A source task aimed at experienced space must say whether it concerns a local measurement space, a foliation, or a signal-based reconstruction. Those alternatives cannot be merged by calling each a spatial slice.

The static-support, flat rotating-observer, and zero-vorticity frame-dragging examples provide diagnostic checks against overbroad interpretations. They test proposed identifications among observer quantities and curvature, not the existence of all possible source reconstructions. An AI system should be able to explain which identification a counterexample defeats and which wider questions it leaves open. The next source-level milestone is a construction or obstruction with that same precision, not another change of notation for (g, u) .

5.12 Interpretive scope and conclusion

The congruence description makes the flow vocabulary mathematically usable: expansion measures volume change, acceleration distinguishes support from free fall, vorticity tests hypersurface orthogonality, and shear measures anisotropic deformation rate. Curvature controls tidal relations and enters the evolution of these quantities. Their roles overlap in particular situations but are not interchangeable.

This is an account of observers and geometry within the adopted GR model. It adds no independent gravitational degree of freedom and no empirical prediction beyond the corresponding effective model. It also supplies no source-to-congruence map, source clock, or proof that physical spacetime is a slice of $\mathcal{A}$. Those claims require source structure not given here. The geometric results remain useful precisely because their observer choice, normalization, and domain are explicit. A source-level geometry task can now use these distinctions as explicit targets and diagnostic checks. Its result must identify the assumptions and reconstruction responsible, whether it is a partial construction or an obstruction. The present effective geometry supplies neither a source success nor a source impossibility theorem.

The Exact-GR Benchmark

A compact statement of adoption, interpretation, and open derivation

Abstract. The Æther-flow Interpretation of Relativity fixes its effective gravitational benchmark to general relativity while leaving its source structure incomplete. A smooth four-dimensional source manifold is assumed, but a source evolution law and a reconstruction of spacetime are not specified. This note states the effective action, its operational assumptions, and the conditions for observational equivalence. It separates those adopted results from the research objective of deriving relativity or establishing a scoped obstruction from a defined source proposal. It also summarizes how that objective can serve as an artificial-intelligence research case study. Verified deductions and valid negative results can both demonstrate successful reasoning; an unsuccessful search cannot establish physical impossibility. Exact closure remains a benchmark choice, not a completed source derivation or a validated evaluation result.

6.1 Purpose and source assumptions

The central claim can be stated directly: use general relativity (GR) for measurable gravitational physics, while investigating whether a separate source structure can explain why that effective description applies. The first part specifies the effective model. The second is a research question.

At source level, $\mathcal{A}$ denotes a smooth four-dimensional manifold, not physical spacetime. The symbol Φ_{src} stands for a possible source-order or evolution structure whose mathematical type is unresolved. No parameter domain, composition law, generator, stochastic rule, gauge interpretation, or physical clock is assigned to it. The source assumptions therefore do not yet constitute a dynamical theory.

The working hypothesis is intrinsic ordered motion in a four-dimensional source, with observers inside it. Experienced space and S-time name the intended spatial and temporal account. The aim is to generate the metric, not a wind within space. This construction is not supplied, and the observer velocity u^μ is not identified with Φ_{src} .

6.2 Definition of the effective benchmark

Let (M, g) be a time-oriented Lorentzian spacetime with signature $(- + + +)$ and Levi-Civita connection. The covariant action uses length-valued coordinates, with $x^0 = ct$, and a physical matter action S_m . The gravitational model is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad G > 0. \quad (6.1)$$

Λ is constant; ψ denotes specified ordinary matter fields coupled to the same metric. The noncosmological specialization is

$$S_{\text{eff}}|_{\Lambda=0} = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} R + S_m[g, \psi]. \quad (6.2)$$

With the physical normalization

$$T_{\mu\nu} = -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}, \quad (6.3)$$

the bulk metric equation is

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (6.4)$$

For $\Lambda = 0$, it reduces to the original benchmark equation

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (6.5)$$

Compactly supported variations suffice for this bulk statement. A finite-boundary variational problem must include its boundary terms and data. Dynamics provides the explicit variation; the construction is the standard one for GR (Carroll, 1997).

Ordinary matter is a substantive assumption. The action symbol does not, by itself, guarantee healthy propagation for every conceivable choice of S_m . The familiar optical and clock interpretations require the standard locally Lorentz-invariant matter models used for those measurements. The action must be specified, not merely named, before numerical predictions follow.

6.3 Meaning of exact closure

Exact closure fixes the effective gravitational equations to GR throughout their stated classical domain, not only a Newtonian limit. It specifies the target a source construction must recover; it does not provide that construction or complete the source explanation, matter physics, or boundary conditions.

For full observational equivalence, compare models with the same matter action, constants, admissible solution class, initial and boundary data, and measurement rules. If the source interpretation introduces no additional coupling or restriction, corresponding observables are identical. That conclusion follows from the identity of the effective model, not from a derivation using $\mathcal{A}$ and Φ_{src} .

A source rule that admitted only a subset of GR solutions would alter this statement. So would a new matter coupling, clock rule, or light-propagation law. These possibilities cannot be declared observationally absent solely because the displayed Einstein equation is unchanged. They must either be excluded as part of the benchmark definition, as here, or explicitly compared.

6.4 Clocks, light, and local relativity

The metric defines causal directions. For standard Maxwell light in vacuum geometric optics,

$$g_{\mu\nu} k^\mu k^\nu = 0, \quad (6.6)$$

and the rays are affinely parametrized null geodesics. Ideal clocks along timelike paths satisfy

$$d\tau^2 = -\frac{1}{c^2} g_{\mu\nu} dx^\mu dx^\nu. \quad (6.7)$$

Neutral structureless test particles follow timelike geodesics in the absence of nongravitational forces and relevant finite-size effects. These domain conditions matter: a supported observer, a charged particle in a field, or light in a medium requires a more specific description.

At an event, a local orthonormal frame gives the ordinary Minkowski tangent-space metric and local Lorentz transformations. Curvature remains over finite regions. A congruence chosen to discuss measurements does not select a new preferred frame for the effective laws.

S-time has not been calibrated to (6.7). An account of source order would still need to explain how operational durations, causal relations, and their observer dependence arise. The familiar clock formula is a benchmark that such an account must match, not a substitute for it.

6.5 Cosmological completion

A constant Λ may be written as the vacuum stress tensor

$$T_{\mu\nu}^{(\Lambda)} = -\frac{\Lambda c^4}{8\pi G} g_{\mu\nu}, \quad p_\Lambda = -\rho_\Lambda c^2, \quad \rho_\Lambda = \frac{\Lambda c^2}{8\pi G}. \quad (6.8)$$

Moving this tensor to the matter side is an exact rewriting of the same model. It should not also be retained as an independent duplicate on the geometric side.

A generic dark-energy field or fluid is different. Its action, equation of state, perturbations, and possible exchanges with other components must be specified. Covariant conservation alone does not make its stress tensor equivalent to (6.8). In homogeneous cosmology, positive Λ contributes toward acceleration; the total sign of $\ddot{a}$ still depends on the other densities and pressures.

The source manuscripts' supernova and Planck references motivate consideration of a conventional accelerated-expansion benchmark (Perlmutter et al., 1999; Planck Collaboration, 2020; Riess et al., 1998). They neither establish the proposed ontology nor derive the value of Λ . This note reports no new observational analysis.

6.6 The restricted weak-field result

For an isolated, slowly moving source with negligible leading pressure and anisotropic stress, the ordinary quasi-static weak-field solution in an isotropic chart is

$$ds^2 \simeq -\left(1 + \frac{2\Phi}{c^2}\right) c^2 dt^2 + \left(1 - \frac{2\gamma\Phi}{c^2}\right) d\mathbf{x}^2, \quad (6.9)$$

where $\nabla^2\Phi = 4\pi G\rho$ and

$$\gamma = 1. \quad (6.10)$$

The approximation assumes $|\Phi|/c^2 \ll 1$ and a negligible cosmological-constant contribution on the local scale. The relation is obtained from the adopted GR equations. It is not evidence that an independent source mechanism has already recovered both metric potentials.

In the same scaling regime, with $\epsilon = \sup|\Phi|/c^2$, $v^2/c^2 = O(\epsilon)$, and variation scale L , the leading test-particle acceleration is

$$\ddot{x}^i = -\partial_i\Phi + O(c^2\epsilon^2/L). \quad (6.11)$$

The error estimate is dimensionful and conditional. Relativistic Recovery develops this limit and the corresponding clock and light relations.

6.7 Observer flow within GR

A normalized timelike field u defines a family of observers. Its derivative separates into expansion, shear, vorticity, and acceleration:

$$\nabla_\mu u_\nu = \frac{1}{3}\theta h_{\mu\nu} + \sigma_{\mu\nu} + \omega_{\mu\nu} - \frac{u_\mu a_\nu}{c^2}, \quad h_{\mu\nu} = g_{\mu\nu} + \frac{u_\mu u_\nu}{c^2}. \quad (6.12)$$

These standard quantities describe volume change, anisotropic deformation rate, local twist, and departure from geodesic motion (Ellis & van Elst, 1999). Their definition presupposes the effective metric and the chosen observers.

The four quantities should not be treated as a one-to-one classification of gravitational effects. Static support may have zero expansion, shear, and vorticity in a curved spacetime. Zero-angular-momentum observers can be hypersurface normal in a frame-dragging geometry (Gourgoulhon, 2010). Geometry develops the relevant counterexamples and the distinction between shear, tidal curvature, and wave strain.

No independent congruence action is added here, so no extra Æther-flow gravitational mode follows from this language. The local GR bulk mode count remains two gravitational configuration degrees of freedom (Arnowitt et al., 1962). This does not prove that the undefined source proposal has the same count or is dynamically consistent.

6.8 The open source derivation

A source action remains schematic. An expression of the form

$$S_{\text{src}}[q] = \int_{\mathcal{A}} \mathcal{L}_{\text{src}}[q] \quad (6.13)$$

would require a specified source configuration q , an integrable density or top-degree form, variations, and appropriate boundary assumptions. A metric or a volume measure cannot be silently imported to complete it. The current symbol Φ_{src} does not provide those ingredients.

A meaningful reconstruction must state which source assumptions give which target result. Reconstructing a causal relation is not automatically a derivation of proper time; constructing a metric is not automatically a derivation of its Einsteinian dynamics or universal matter coupling. The bridge must also specify the regime in which its result applies and any restrictions it imposes on the effective solutions.

6.9 Research outcomes and AI-assisted use

The project combines a physics question with a proposed test of AI-assisted research. The physics question is whether a specified source proposal can recover a declared part of GR without assuming that result. The AI question is whether a research system can make independently checkable progress, including a justified negative finding. Sys4AI is the intended framework for coordinating this work; no completed system evaluation is reported here.

Each task must freeze its assumptions, permitted constructions, target, and validity domain. A fixed-assumption audit keeps those premises unchanged. An open-construction task may add premises, but must record them and state the resulting claim conditionally. A renamed metric or Einstein–Hilbert action remains an assumption when inserted as an input.

A derivation establishes the target under its stated conditions. A counterexample satisfying the premises

but violating the target shows non-entailment. An obstruction excludes recovery for the explicitly defined class and admissible reconstructions. A search finding none of these remains unresolved. The latter is not proof that GR cannot follow, and non-entailment is not proof that no compatible completion exists.

A valid obstruction can be a scientific result against a particular source proposal and a successful demonstration of reasoning. Performance assessment must therefore report the physical outcome separately from the correctness of the AI's argument. The synthesis article gives the proposed evidence and comparison protocol. The present manuscripts are a research case study and reference set, not a validated general test of scientific competence.

6.10 Benchmark policy and conclusion

One may retain the adopted GR model as a reference when a proposed extension fails. This is a research policy, not a logical guarantee that all source hypotheses survive every failure. An unsuccessful extension can still reveal that one of its assumptions is false or incompatible with the intended target.

The benchmark defined in this note is exact GR with a specified matter sector and measurement interpretation. Its equations, observable relations, and local gravitational modes are inherited. The source ontology remains a limited proposal whose physical reality and explanatory connection to that benchmark have not been established. Keeping this distinction explicit preserves both a usable model for calculation and a precise question for further research. The same boundary governs the AI research case study. The desired output is an independently checkable derivation or scoped negative argument, with an unresolved category when neither is obtained. The physical fate of a candidate and the correctness of the research process remain separate judgments.

Overview of the Æther-flow Framework

Exact-GR benchmark and source proposal

Abstract. The Æther-flow program has a physics objective and an artificial-intelligence research objective. The physics task is to determine whether a specified development of its source ontology can recover a declared part of general relativity without assuming that result, or whether an obstruction can be established within an explicit scope. The research-system task is to evaluate independently checkable progress on those questions. The current manuscripts provide a general-relativistic comparison model and a limited source proposal: a smooth four-dimensional arena with an untyped flow placeholder. This overview maps their contributions and sets out a bounded feasibility sequence. It distinguishes fixed-assumption audits from open construction, and successful derivations from counterexamples, scoped impossibility results, and unresolved searches. No source recovery theorem, universal no-go theorem, or completed system evaluation is reported.

7.1 Two questions, two levels of description

The Æther / Æther-flow framework asks whether observed spatial and temporal organization can be understood through a source structure not identified in advance with physical spacetime. The framework also makes a separate effective choice: use general relativity (GR) for the quantitative gravitational description. Keeping these two statements distinct is the organizing principle of the presentation.

The physical intent is an account of GR in which intrinsic ordered motion of a four-dimensional substrate produces the spacetime relations accessible to observers within it. The source is not intended as a wind through pre-existing three-dimensional space. Local spatial experience and S-time express the proposed connection between source organization and measurement. Preserving GR is the target of this construction, not evidence that the construction has already been achieved. The mathematical questions below test that hypothesis and admit a justified result against a specified candidate.

At source level, $\mathcal{A}$ denotes a smooth four-dimensional manifold and Φ_{src} is an untyped placeholder for a possible order or evolution structure. These objects do not yet supply a source metric, causal order, clock, matter model, or reconstruction map. At effective level, (M, g) is a Lorentzian spacetime with ordinary matter laws and operational definitions of measured quantities. There is no identification $\mathcal{A} = M$ or $\Phi_{\text{src}} = u$ between the source placeholder and an observer four-velocity.

The term *exact closure* means that the effective gravitational law is fixed to the corresponding GR law, rather than approximately matched in one limit. It does not mean that the source ontology is complete, that all matter has been specified, or that the physical origin of the Einstein equation has been explained.

7.2 Common effective statement

We use signature $(-+++)$, $x^0 = ct$ in length-valued covariant coordinates, a physical matter action S_m , and a constant Λ . The benchmark action is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad (7.1)$$

with $G > 0$ and specified ordinary matter fields ψ . It yields

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad T_{\mu\nu} = -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}. \quad (7.2)$$

The $\Lambda = 0$ specialization is used for isolated flat-background calculations. No independent source-flow field is varied in (7.1).

Standard vacuum light in geometric optics and ideal timelike clocks obey

$$g_{\mu\nu} k^\mu k^\nu = 0, \quad (7.3)$$

$$d\tau^2 = -\frac{1}{c^2} g_{\mu\nu} dx^\mu dx^\nu. \quad (7.4)$$

These are ordinary GR relations (Carroll, 1997). Observational identity further requires matching matter, constants, admissible solutions, initial and boundary data, and measurement procedures. An added source restriction on solutions would have to be included in the comparison; it cannot be hidden behind equality of the displayed field equations.

A constant Λ is equivalent to the specific vacuum tensor $T_{\mu\nu}^{(\Lambda)} = -\Lambda c^4 g_{\mu\nu} / (8\pi G)$, counted on one side of the equation only. A general conserved dark-energy component is a different matter model and needs its own specification. Cosmological acceleration is an allowed consequence of suitable GR models, not a source-to-GR derivation.

7.3 What each article contributes

The framework is developed across related expositions of one proposal. Their reading order is as follows.

Exact-Closure Note. The compact statement of the effective model and its conditional relationship to GR. It is the shortest route to the central distinction between adopted dynamics and a proposed source explanation.

Foundations. The first technical paper defines the narrow source assumptions. It separates the smooth source arena from spacetime and treats experienced spatial slices and S-time as reconstruction aims rather than derived structures.

Dynamics. The second technical paper fixes the action and stress–energy units, derives the effective equation, and specifies matter coupling, conservation, trajectories, and the weak-field limit. It distinguishes an arbitrary matter template from a fully specified physical model.

Consistency. The third technical paper gives the standard gauge and constraint analysis of the adopted gravitational sector. Its explicit stability calculation concerns weak radiative perturbations about Minkowski vacuum, not every GR background or a source theory that has not yet been defined.

Relativistic Recovery. The fourth technical paper derives representative observable relations: Newtonian acceleration, null propagation, static redshift, proper time, local special relativity, and homogeneous cosmology. Its results are inherited from the corresponding GR models under stated conditions.

Geometry. The fifth technical paper develops observer rest spaces and congruence kinematics. It derives Raychaudhuri evolution with consistent units, distinguishes static support from free fall, gives a zero-vorticity frame-dragging example, and separates shear, curvature, and wave strain.

Synthesis article. The synthesis article gives a self-contained conceptual account of the source proposal and the effective interpretation. It explains the explanatory gap and relates the approach to other gravitational research without claiming a new low-energy field law.

A reader seeking the main conceptual argument can begin with the synthesis, which also specifies the proposed AI evaluation protocol, and use the technical papers for the corresponding calculations. The canonical comprehensive manuscript contains these eight expositions in one document; the standalone versions retain the same scientific content for focused reading.

7.4 Results and shared limits

The effective action has the usual local GR gravitational degrees of freedom. The Arnowitt–Deser–Misner (ADM) count and the Minkowski transverse-traceless analysis yield two radiative gravitational polarizations (Arnowitt et al., 1962). No additional *Æther-flow* vector or scalar gravitational mode has been introduced. This does not count matter modes or prove stability on arbitrary backgrounds.

The geometry paper uses standard observer kinematics: expansion, shear, vorticity, and acceleration (Ellis & van Elst, 1999). These are properties of a metric and chosen observer family. They neither define a source flow nor provide a unique taxonomy of gravity. In particular, congruence vorticity is not synonymous with frame dragging, and zero shear does not imply absence of tidal curvature.

The shared source limit is equally precise. The papers do not derive Lorentzian signature, metric dynamics, ordinary matter, operational clocks, or measured constants from $\mathcal{A}$ and Φ_{src} . They also do not establish that a literal substrate exists. The absence of such results is not a proof that all possible source completions fail; it identifies which argument is still required.

7.5 Relation to other approaches

The effective benchmark belongs to standard GR, not to a preferred-structure modification. Einstein-aether theory adds a dynamical timelike vector, whereas Hořava’s proposal changes the short-distance spacetime scaling structure (Hořava, 2009; Jacobson & Mattingly, 2001). Sharing words such as aether or flow does not make these actions equivalent to (7.1).

The source ambition is closer in subject to approaches that seek an explanation for relativistic dynamics from other assumptions. Examples include thermodynamic reasoning, induced gravity, and consistency-based self-coupling (Deser, 1970; Jacobson, 1995; Visser, 2002). Each has its own premises. Their existence does not supply the missing premises of this ontology, and a successful result in one approach cannot be transferred here by analogy. The detailed comparison is reserved for the synthesis rather than repeated as a large citation cluster in every article.

7.6 Use of the benchmark

The effective GR model can remain a comparison target while source work proceeds. If a proposed extension fails, one may set that extension aside and retain the independent benchmark. This is a research decision, not a proof that source assumptions are immune to failure. Any assumption implicated by a failed calculation

must be reconsidered on its own merits.

7.7 A bounded feasibility sequence

The intended research infrastructure is Sys4AI, an agentic AI framework for coordinating roles, tasks, and verification. The manuscripts specify the research question and comparison physics; they do not report a running evaluation implementation or measured agent performance. The two-track intent is to investigate the source proposal and to assess the reliability of the research process used to investigate it.

The initial feasibility task is narrower than deriving all of GR at once. First, define a versioned source model and one reconstruction target, such as local causal geometry or a clock relation. State every additional assumption and exhibit an admissible source example or the relevant existence argument. Then seek a derivation, a counterexample to the stated implication, or an obstruction within a declared class. Preserve an unresolved outcome when none is established.

The next stage depends on the result. A verified geometric construction may support a separate dynamics or matter-coupling task. A negative result should identify the premise or reconstruction restriction responsible and delimit which alternatives remain open. An unfinished argument should identify its missing step. No stage receives credit for importing the desired target into its assumptions.

Each completed task should yield a self-contained claim, an assumption record, the argument or witness, supporting calculations where needed, and an independent checking record. Resource use and human intervention belong in the AI evaluation record. A valid result is assessed at its actual scope; a partial geometric construction does not become a full GR derivation because later stages are planned.

Two modes organize this sequence. A fixed-assumption audit evaluates the proposal as given. Open construction permits explicit additions and assesses the strengthened proposal. The synthesis article provides the fuller evaluation protocol, including controls, comparable budgets, and the distinction between a base model and the entire research system. This is a proposed procedure, not evidence that these stages have already been completed.

7.8 Conclusion

The framework presents a definite effective gravitational model and a deliberately limited source proposal. Its technical value lies in explicit conventions, calculations, observer choices, and limits. Its open question is whether a specified source theory can reproduce those structures without assuming them in disguised form. The bounded next step is a versioned source specification and a checkable claim. The physics and AI tracks can both learn from a valid obstruction, while an unfinished search remains unfinished. Performance claims await an implemented protocol, recorded runs, and independent assessment.

Æther-flow as an Interpretation of General Relativity

Source assumptions, observer geometry, and the explanatory gap

Abstract. The Æther / Æther-flow proposal seeks a physical account of spatial and temporal organization through intrinsic ordered motion in a proposed four-dimensional substrate, while retaining general relativity as the recovery target. Its present mathematical assumptions are limited to a smooth four-dimensional arena and an unresolved source-flow placeholder. General relativity is adopted separately as the quantitative benchmark, with observer congruences supplying an effective interpretation of flow. This article relates that distinction to a dual research objective: test a source explanation of relativity and evaluate whether an artificial-intelligence research system can make independently checkable progress. It specifies assumption control, recovery targets, scoped negative outcomes, and a proposed evaluation protocol. Fixed-assumption audits are separated from constructions that add new premises. The work establishes the boundaries and evidence requirements of this research case study, not a source derivation, a universal impossibility theorem, or measured model performance.

8.1 The proposal and its central distinction

General relativity (GR) gives a quantitative account of spacetime geometry and its relation to matter. The Æther / Æther-flow framework asks a different, additional question: could the spatial and temporal relations used in that account arise from a source organization not identified at the outset with physical spacetime? The language of Æther and Æther-flow is intended to express this source-level possibility.

A possibility is not yet a model. To explain spacetime from another structure, one must define the source objects, state their laws or constraints, and demonstrate how the operational content of relativity follows. The present source specification has not completed these steps. Its effective model instead takes GR as an explicit starting assumption.

The distinction prevents two opposite mistakes. The first would be to announce that the ontology has derived Einsteinian gravity because its effective equations have been chosen to be Einstein's equations. The second would be to conclude that the source proposal must be false simply because a derivation has not been supplied. Neither conclusion follows from the stated assumptions. The framework provides an adopted comparison model and an unfinished explanatory proposal.

The research goal is to test that explanatory proposal and, separately, the ability of an AI-assisted system to produce reliable results about it. The evidence requirements below allow a valid scoped negative result to count as successful research.

This article brings those parts together. The technical companion papers give the action variation, perturbative mode count, operational limits, and congruence calculations. Their standard GR content should be read as inherited physics, not as a claim that the proposal has independently discovered those results.

8.2 What is assumed about the source

The source arena is denoted by $\mathcal{A}$ and assumed to be a smooth four-dimensional manifold. It is not identified with the effective spacetime manifold M . Its dimension and smooth structure are primitive assumptions,

not products of a source derivation. No particular metric, physical causal order, clock, or matter content is supplied on $\mathcal{A}$.

The symbol Φ_{src} is a placeholder for a possible source-order or evolution structure. Its mathematical type remains unresolved. It has not been specified as a one-parameter flow, a vector field, a stochastic process, or a generator. The usual properties of a mathematical flow therefore cannot be inferred from its name. In particular, no parameter has been identified as physical time.

Intrinsic ordered motion is the working physical hypothesis, not merely an alternative name for observer motion in GR. The proposed substrate is meant to generate the metric rather than move through a previously specified space. Observers belong to the same underlying reality; their accessible spatial and temporal relations would be local manifestations of its organization. This is the intended explanation, not a mechanism already constructed from $\mathcal{A}$ and Φ_{src} .

Experienced space and S-time retain their roles in that hypothesis. A local spatial slice expresses the intended account of three-dimensional measurement, while S-time links temporal experience to ordered change. The present assumptions do not define an embedding of that slice in $\mathcal{A}$, a monotone clock functional, or an explanation of subjective experience. They also do not authorize treating an unspecified source parameter as physical time. Foundations separates these conceptual commitments from the mathematical structures that a reconstruction must supply.

This restraint does not remove the original source question. It makes the proposed explanation evaluable in principle. A future source model must say what carries the order, what counts as a state or process, which transformations are allowed, and how a physical observer can obtain measurements from it. Until those relations exist, referring to $\mathcal{A}$ as underlying reality is a hypothesis about physical interpretation, not an experimentally established fact.

8.3 A definite effective gravitational model

8.3.1 Fields and equations

The effective arena is a time-oriented four-dimensional Lorentzian spacetime (M, g) with signature $(-+++)$ and Levi-Civita connection. In length-valued covariant coordinates, $x^0 = ct$. With a physical matter action S_m , the adopted model is

$$S_{\text{eff}}[g, \psi] = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} (R - 2\Lambda) + S_m[g, \psi], \quad (8.1)$$

where $G > 0$, Λ is constant, and ψ denotes specified ordinary matter. The $\Lambda = 0$ specialization is

$$S_{\text{eff}}|_{\Lambda=0} = \frac{c^3}{16\pi G} \int_M d^4x \sqrt{-g} R + S_m[g, \psi]. \quad (8.2)$$

The stress tensor and bulk metric equation are

$$T_{\mu\nu} = -\frac{2c}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}, \quad G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (8.3)$$

These are the ordinary GR dynamics with the action units stated explicitly. Compactly supported variations suffice for the bulk equation; specific boundary problems need the corresponding boundary terms and data.

No independent Φ_{src} field appears in (8.1). Thus there is no additional low-energy gravitational equation,

coupling parameter, or source-flow mode to compare with the metric sector. The usual local gravitational degree-of-freedom count is retained (Arnowitt et al., 1962). That is a property of the chosen action, not evidence that an undefined microscopic theory has already reduced to two modes.

8.3.2 Operational meaning

For standard Maxwell light in vacuum geometric optics,

$$g_{\mu\nu}k^\mu k^\nu = 0, \quad (8.4)$$

with null geodesic propagation. For an ideal timelike clock,

$$d\tau^2 = -\frac{1}{c^2}g_{\mu\nu}dx^\mu dx^\nu. \quad (8.5)$$

These formulas connect the metric to measurement (Carroll, 1997). They require the ordinary matter laws used to model light and clocks; the phrase minimal coupling is not a proof that an arbitrary matter action has the same propagation properties.

The benchmark agrees exactly with the corresponding GR model only when matter, constants, allowed solutions, initial and boundary data, and measurement procedures match. A hidden source constraint selecting different solutions would change that content. Here no such constraint is specified. The statement of agreement is consequently conditional model identity, not an additional empirical success attributable to the ontology.

The proposed physical explanation aims to preserve this agreement without an additional direct flow force on matter. The intended mechanism would recover the common metric and its coupling to measuring systems. Adopting metric-only coupling and exact GR dynamics already gives the reference model, but does not establish that a deeper source produces it. Nor does metric-only matter coupling alone exclude observable preferred structure in a candidate with different gravitational dynamics or restrictions. Those dependencies belong in the reconstruction and its checks.

8.3.3 Cosmology without a new source law

The cosmological constant in (8.1) may be moved to the matter side as

$$T_{\mu\nu}^{(\Lambda)} = -\frac{\Lambda c^4}{8\pi G}g_{\mu\nu}. \quad (8.6)$$

That specific constant vacuum tensor is equivalent to Λ . A general dark-energy field or fluid is not; its dynamics and perturbations must be supplied separately. The same component must not be counted on both sides of the field equation.

For a Friedmann–Lemaître–Robertson–Walker (FLRW) model with non- Λ density ρ and pressure p ,

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{c^2}\right) + \frac{\Lambda c^2}{3}. \quad (8.7)$$

Positive Λ contributes toward acceleration but need not outweigh matter at every epoch. The supernova and Planck references provide motivation for studying accelerated cosmological benchmarks (Perlmutter et al., 1999; Planck Collaboration, 2020; Riess et al., 1998). They do not identify dark energy with Φ_{src} , derive the value of Λ , or establish a source origin of GR.

8.4 What an observer-based flow description can explain

Once a spacetime and a family of observers have been selected, the word flow has a precise mathematical use. A timelike congruence has four-velocity $u^\mu = dx^\mu/d\tau$, normalized by $u^\mu u_\mu = -c^2$. Its local rest-space projector is

$$h_{\mu\nu} = g_{\mu\nu} + \frac{u_\mu u_\nu}{c^2}. \quad (8.8)$$

The covariant derivative separates as

$$\nabla_\mu u_\nu = \frac{1}{3}\theta h_{\mu\nu} + \sigma_{\mu\nu} + \omega_{\mu\nu} - \frac{u_\mu a_\nu}{c^2}. \quad (8.9)$$

Here θ is the local volume-change rate, σ is the anisotropic deformation rate, ω is vorticity, and a is proper acceleration. This is the standard congruence description (Ellis & van Elst, 1999).

The description is useful because it separates effects that a single image of moving substance would merge. In homogeneous cosmology, the comoving congruence has $\theta = 3H$ and zero shear, vorticity, and acceleration. For static supported observers, all three spatial kinematic rates can vanish while the proper acceleration is $a_\mu = c^2 D_\mu \ln N$. An expanding universe therefore need not swirl, and static gravity need not involve expanding observer volume.

Rotation requires additional care. The zero-angular-momentum observers used in a circular stationary-axisymmetric geometry are hypersurface normal and have zero vorticity, even though their angular velocity relative to an appropriate asymptotic reference can reflect frame dragging (Gourgoulhon, 2010). Conversely, rotating observers in flat spacetime can have nonzero vorticity. The congruence and reference observers must be specified before assigning a physical interpretation to rotational quantities.

Likewise, shear is not curvature itself. In a weak vacuum transverse–traceless (TT) wave, a chosen freely falling congruence has $\sigma_{ij} = \dot{h}_{ij}^{\text{TT}}/2$ at first order, while its tidal acceleration per unit separation involves $\ddot{h}_{ij}^{\text{TT}}/2$. A static congruence can have zero shear in the presence of nonzero tides. Geometry gives these relations, the curvature convention, and the corrected Raychaudhuri identity in full.

The result is an effective interpretive vocabulary, not a source mechanism. All these quantities are calculated after g and u have been supplied. Declaring them to be manifestations of Φ_{src} without a map would rename target objects rather than derive them.

8.5 Explanatory value and empirical limits

A physical interpretation may be useful by clarifying concepts, suggesting a mathematical construction, or connecting otherwise separate phenomena. Those possibilities should be evaluated directly. They do not follow automatically from exact agreement with a theory that was adopted at the outset.

For this proposal, the key explanatory question is whether source assumptions can eventually reduce what must be independently assumed at effective level. If a derivation simply postulates a Lorentzian metric with Einstein–Hilbert dynamics under new names, the measured theory may be sound but the intended explanation has not advanced. If instead independently specified source relations imply an operational causal structure, a metric, or part of its dynamics, the scope and assumptions of that result can be assessed.

Empirical equivalence also limits confirmation. Where the proposed description and ordinary GR assign the same outcomes to every measurement by construction, a successful GR test does not discriminate in favor of this source ontology over GR without it. This does not logically refute the ontology. It means that its additional explanatory claim needs evidence of its own, such as a demonstrable reduction of assumptions or a

discriminating prediction from a completed model.

The sparse source specification cannot currently support a full consistency assessment. Without defined source states and dynamics, one cannot count its propagating modes, derive its conservation laws, or establish its stability. The absence of those definitions is a concrete limitation, not a theorem that no suitable definitions can be found.

8.6 Relationship to other gravitational approaches

8.6.1 Standard GR and preferred-structure theories

The effective benchmark is ordinary GR. Congruence kinematics and Arnowitt–Deser–Misner (ADM) variables are standard ways to analyze that theory, not additional evidence for a source ontology. The word *Æther* must therefore be distinguished from its use in other actions.

Einstein-aether theory introduces a dynamical unit timelike vector and studies its coupled gravitational modes (Jacobson & Mattingly, 2004; Jacobson & Mattingly, 2001). Hořava’s proposal instead uses anisotropic short-distance scaling and a different spacetime-symmetry setup (Hořava, 2009). The present effective action introduces neither structure. A future source completion that produced observable preferred structure would need a new analysis rather than relying on this terminological distinction.

8.6.2 Derivational and analogue approaches

Several established lines of work seek relations between Einsteinian dynamics and other principles. Jacobson’s thermodynamic argument uses horizon entropy and the heat relation for local causal horizons (Jacobson, 1995). Induced-gravity arguments concern how gravitational terms can arise in an effective description of other fields (Visser, 2002). Self-coupling arguments begin with a defined gauge field and use consistency requirements to motivate nonlinear interactions (Deser, 1970). Analogue-gravity systems can reproduce aspects of effective Lorentzian propagation without automatically reproducing the full Einstein dynamics (Barceló et al., 2011).

These are comparisons of explanatory tasks, not derivations of the present proposal. Each begins with definite mathematical and physical assumptions. The untyped source placeholder here does not inherit their conclusions. Related questions about canonical reconstruction and uniqueness likewise require their own hypotheses; general covariance or four-dimensionality alone is not the missing argument.

8.7 A dual physics and AI research objective

The project’s physical aim is to determine whether a specified development of the source ontology can account for relativistic structure without importing the claimed result as a premise. A constructive derivation is one possible outcome. A rigorous obstruction within a declared model class is another. The program does not require a favorable verdict on the ontology to count the investigation as scientifically useful.

The second aim is to study AI-assisted research itself. A system must do more than generate a plausible account of gravity. It must formulate the question, keep assumptions distinct from consequences, check its calculations, seek counterexamples, and report uncertainty when an argument remains incomplete. A negative physical result can demonstrate successful reasoning; an attractive but invalid derivation cannot.

Sys4AI is the intended agentic AI framework for organizing research roles, tasks, tool use, and verification records. These are proposed uses in this project, not reported implementation results. The following protocol defines a research case study that could be implemented and independently assessed. It does not establish the

performance of any particular model, agent arrangement, or software release.

8.8 Task specification and the meaning of success

8.8.1 Freeze the question before assessing the answer

A task must identify a source version, its mathematical objects and laws, the admissible source models, and the permitted reconstruction maps. It must name the effective target and its physical domain. Examples of different targets are a local causal structure, an operational clock, gravitational dynamics in a controlled limit, and exact recovery of a specified classical GR model. A result about one target must not be promoted to another by changing the wording after the search.

The source assumptions need not be complete before AI participates. In a fixed-assumption audit, an agent analyzes the supplied proposal without adding premises. In open construction, it may propose missing definitions or laws, but must label every addition and explain its role. Changes create a new version of the question. The original and strengthened assumption sets must remain distinguishable in the record.

A task must also address existence. An inconsistent source specification does not derive physically meaningful GR through a vacuous implication. An admissible example, a suitable existence argument, or an explicit unresolved existence condition is needed. A theorem may be conditional on such a condition, but its research status must state that dependence.

The protocol does not demand that every task derive all matter or every measured constant. A gravitational result may assume a particular matter sector, and an approximate result may apply only at certain scales. Those restrictions belong in the claim. The question is what explanatory content the source construction adds after its independently supplied inputs are accounted for.

8.8.2 Separate the outcome categories

A *verified derivation* establishes the declared implication under the stated assumptions and regime. A proposed derivation remains provisional until the relevant checks support it. Matching one familiar spacetime or one finite set of observables establishes only that narrower result unless a broader argument is supplied.

A *counterexample to entailment* satisfies the task's source assumptions but violates its required conclusion under the specified reconstruction. It disproves that implication. It does not establish that every other source completion or reconstruction must fail. In particular, premises may permit both GR-compatible and non-GR completions, in which case they do not select GR uniquely.

A *scoped obstruction* establishes that no admissible member of the defined model or reconstruction class can meet the target under the stated conditions. Its quantifiers matter: excluding one construction is not excluding every possible construction. A conclusion about a version of the ontology does not automatically apply to later versions with different premises. A hypothetical model forcing all reconstructed metrics to be flat would exclude recovery of a target class containing curved solutions, but this restriction has not been proved for the present ontology.

An *unresolved search* produces neither a verified positive result nor a valid negative argument. It may yield useful lemmas, calculations, or a precise missing step, which should be reported separately. Exhausting time, tokens, or a list of candidate models is not a proof of impossibility unless the task itself is a justified finite exhaustive decision problem and the search is complete.

Finally, a construction showing that some permitted completion is compatible with GR is an existence result at that scope. It is not a proof that the original assumptions force GR, nor that the completion is unique.

The evaluation must preserve this distinction between compatibility, entailment, and exclusion.

8.9 Proposed evaluation protocol

8.9.1 Evidence beyond restating the manuscripts

The existing papers already document the lack of a source-to-GR reconstruction. An agent that repeats that finding has performed an audit or summary, not demonstrated new research. Credit for research progress requires an additional inspectable result: a precise source specification, a nontrivial implication, a justified reconstruction step, a counterexample, or a previously unrecognized defect in an argument.

Assessment should record the physical outcome separately from the correctness and scope of the research artifact. A correct negative theorem should not be scored as a reasoning failure merely because it rules out the candidate. A partial derivation should receive credit for its proved portion without being called full recovery. Unsupported confidence, concealed assumptions, and invalid counterexamples should be identified directly rather than hidden in an aggregate score.

For each result, retain the claim statement, definitions, assumptions, dependencies, derivation or witness, and checking record. Distinguish calculations performed from checks merely proposed. A missing premise cannot be repaired by a convincing narrative. The final claim must be readable and checkable without access to unrecorded agent exchanges.

8.9.2 Independent checking and physical interpretation

A checking stage should seek errors rather than only confirm the proposed result. It should inspect assumptions, intermediate steps, limiting cases, and possible counterexamples. Where practical, it should use a separate calculation, implementation, or reviewer. Assigning an agent a critic role is a process choice; it is not itself evidence that the result was independently validated.

Symbolic computation can check specified algebraic steps. Numerical calculations can test stated cases with reported precision and error control. Formal proof tools can verify a precisely encoded theorem. None of these checks alone establishes that the encoding or model captures the intended physics. Qualified human assessment is needed for the physical interpretation and scope of any consequential recovery or no-go claim. Unchecked parts remain visible, and agreement among agents is not substituted for a proof.

The verification record should identify who or what checked each component, which inputs were used, and the limits of the check. No claim of external review or human approval follows merely from including that requirement in the protocol.

8.9.3 Controlled comparisons and reproducible records

Before comparing models, fix the task set, source snapshot, permitted information, tool access, resource limits, and assessment criteria. Record the model identifier and available version information, settings, prompts, tool and code outputs, elapsed time, token use, actual costs where available, and human interventions. Record seeds where tools expose them; a seed alone does not guarantee identical reruns. Missing resource measurements must be labeled unavailable rather than estimated as if observed.

The comparison unit must be explicit. A base model with a fixed prompt is different from a system using several agents, retrieval, memory, symbolic tools, and human guidance. A result from the full Sys4AI arrangement demonstrates the behavior of that arrangement under the recorded conditions. It does not isolate the base model's contribution.

Useful comparisons would hold the model fixed while varying the workflow, or hold the workflow and task budget fixed while varying the model. A direct single-agent workflow provides a simple baseline. Removing a component in a controlled comparison can test whether it contributes; such a test is often called an ablation. Human help should either be controlled across conditions or reported as a separate condition. Repeated trials and their variability are preferable to selecting a favorable run.

8.9.4 Controls, exposure, and the limit of generalization

The inherited GR calculations can serve as known-answer control tasks. Carefully specified flawed derivations can test error detection; known counterexamples can test whether the system distinguishes non-entailment from a class-wide obstruction. Such controls should have independently checked answers and criteria established before model outputs are graded. Their usefulness must be tested, not assumed from this proposal.

Evaluation should distinguish access to source physics from access to evaluation answers. The manuscripts may be legitimate task inputs, but repeating an answer supplied in them cannot demonstrate its independent discovery. Record which documents and retrieval sources were available. Keep assessment-only material and held-out checks separate where the task design permits, and report uncertainty about prior model exposure rather than claiming it can always be excluded.

This single ontology is a research case study. Even a strong checked result would support a claim about the demonstrated task, not general scientific competence. Broader evaluation would require additional problems, stable assessment criteria, repeated trials, and examination of failure cases. No pass rate, model ranking, cost comparison, or validated benchmark score is available in the present manuscripts.

8.10 Initial feasibility study and stopping conditions

The first study should select one explicit development of the source proposal and one attainable mathematical question. A causal-geometry or clock-reconstruction target is a possible starting point, not a source assumption adopted here. The study should expose its premises, give a construction or admissible example where possible, and attempt a checkable implication or obstruction.

A positive result justifies only the next task supported by its dependencies. For example, a metric construction may motivate a separate dynamics task; it does not complete that task in advance. A negative result should identify the exact restriction responsible and the alternatives it leaves open. An unresolved result should locate the missing step and preserve the failed attempts needed to avoid repeating them without new information.

Stop a run when its predefined resource limit is reached, when the declared result has been checked to the agreed standard, or when further work would change the question without authorization. New assumptions or a larger target belong in a new task version. The research deliverable is a claim with evidence and scope, not a requirement to reach either a positive or negative verdict at any cost.

8.11 Conclusion

The *Æther-flow* Interpretation of Relativity presently combines a tentative source ontology with an adopted GR model. Observer congruences make its effective flow language precise, but do not convert that language into a derivation of spacetime, matter, or clocks. Its standard relativistic predictions are inherited under explicit matching conditions.

The proposal's unresolved question is therefore clear: can a defined source structure produce some or all of

the effective relational and dynamical content without importing that content as a premise? The adopted GR model supplies an intelligible benchmark for addressing that question. This benchmark neither establishes the physical reality of the proposed source nor shows that all future source constructions must fail.

The dual research objective makes both constructive and negative results relevant, but it does not make either available by declaration. A useful first contribution is a checkable result about a defined source model, accompanied by a reproducible research record. A result against that model can still demonstrate capable reasoning. The project remains an open physical proposal and a proposed AI research case study until such evidence is produced.

References

- Arnowitt, R., Deser, S., & Misner, C. W. (1962). The dynamics of general relativity. In L. Witten (Ed.), *Gravitation: An introduction to current research* (pp. 227–265). Wiley. <https://arxiv.org/abs/gr-qc/0405109>
- Barceló, C., Liberati, S., & Visser, M. (2011). *Analogue gravity* [Updated review, version 3]. <https://arxiv.org/abs/gr-qc/0505065v3>
- Carroll, S. M. (1997). *Lecture notes on general relativity*. arXiv: gr-qc/9712019. <https://arxiv.org/abs/gr-qc/9712019>
- Deser, S. (1970). Self-interaction and gauge invariance [Original article; arXiv reprint first posted in 2004]. *General Relativity and Gravitation*, 1, 9–18. <https://doi.org/10.1007/BF00759198>
- Ellis, G. F. R., & van Elst, H. (1999). Cosmological models (Cargèse lectures 1998). In M. Lachièze-Rey (Ed.), *Theoretical and observational cosmology* (pp. 1–116). Kluwer Academic. <https://arxiv.org/abs/gr-qc/9812046>
- Gourgoulhon, E. (2007). *3+1 formalism and bases of numerical relativity*. <https://arxiv.org/abs/gr-qc/0703035>
- Gourgoulhon, E. (2010). *An introduction to the theory of rotating relativistic stars* [Revised preprint, version 2, 2011]. <https://arxiv.org/abs/1003.5015>
- Hořava, P. (2009). Quantum gravity at a Lifshitz point. *Physical Review D*, 79, Article 084008. <https://doi.org/10.1103/PhysRevD.79.084008>
- Jacobson, T., & Mattingly, D. (2004). Einstein-aether waves. *Physical Review D*, 70, Article 024003. <https://doi.org/10.1103/PhysRevD.70.024003>
- Jacobson, T. (1995). Thermodynamics of spacetime: The Einstein equation of state. *Physical Review Letters*, 75, 1260–1263. <https://doi.org/10.1103/PhysRevLett.75.1260>
- Jacobson, T., & Mattingly, D. (2001). Gravity with a dynamical preferred frame. *Physical Review D*, 64, Article 024028. <https://doi.org/10.1103/PhysRevD.64.024028>
- Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., Deustua, S., Fabbro, S., Goobar, A., Groom, D. E., Hook, I. M., Kim, A. G., Kim, M. Y., Lee, J. C., Nunes, N. J., Pain, R., Pennypacker, C. R., Quimby, R., Lidman, C., . . . Couch, W. J. (1999). Measurements of Ω and Λ from 42 high-redshift supernovae. *The Astrophysical Journal*, 517, 565–586. <https://doi.org/10.1086/307221>
- Planck Collaboration. (2020). Planck 2018 results. VI. cosmological parameters. *Astronomy & Astrophysics*, 641, Article A6. <https://doi.org/10.1051/0004-6361/201833910>
- Raychaudhuri, A. K. (1955). Relativistic cosmology. I. *Physical Review*, 98, 1123–1126. <https://doi.org/10.1103/PhysRev.98.1123>
- Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., Gilliland, R. L., Hogan, C. J., Jha, S., Kirshner, R. P., Leibundgut, B., Phillips, M. M., Reiss, D., Schmidt, B. P., Schommer, R. A., Smith, R. C., Spyromilio, J., Stubbs, C., Suntzeff, N. B., & Tonry, J. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116, 1009–1038. <https://doi.org/10.1086/300499>
- Visser, M. (2002). Sakharov’s induced gravity: A modern perspective. *Modern Physics Letters A*, 17, 977–992. <https://doi.org/10.1142/S0217732302006886>