

The *Æ*ther-flow Interpretation of Relativity

Ontology: Flow Geometry

Project creator: Alexander Samuel Ricciardi

Model-assisted research and drafting tools: GPT-5.4 and GPT-5.5.

Project provenance: this document is part of the current managed ontology package.

This title page records project-origin attribution and model-assisted drafting provenance; it does not change the article's scientific content or claim status.

Abstract

This paper gives the next positive extension of the active $\mathcal{A}$ ether-flow Interpretation of Relativity sequence by formalizing $\mathcal{A}$ ether-flow through congruence-based relativistic geometry. The scope is explicit. The manuscript does not alter the effective law of the theory: $\mathcal{A}$ ether-flow Interpretation of Relativity still adopts Einsteinian gravity with universal matter coupling and a single operational metric. Its purpose is interpretive deepening inside exact closure.

The central move is to provide a target-side interpretive dictionary for $\mathcal{A}$ ether-flow using a future-directed unit timelike congruence u^μ on the exact relativistic spacetime and the standard decomposition of $\nabla_\mu u_\nu$ into expansion, shear, vorticity, and acceleration. The congruence is an admissible, generally non-unique observer field in the adopted GR benchmark. It is not the unresolved source slot Φ_{src} , and no source-to-congruence bridge is claimed. Expansion describes local volume change and cosmological growth of the observer congruence. Acceleration governs static gravitational support and redshift structure. Vorticity captures swirl, rotation, and frame dragging. Shear captures anisotropic distortion, tidal structure, and the strain content associated with gravitational waves. Local special relativity is recovered as the local inertial limit of the same target-side geometry.

The resulting claim is precise. In $\mathcal{A}$ ether-flow Interpretation of Relativity, $\mathcal{A}$ ether-flow is not a second low-energy gravitational field competing with the metric. Congruence kinematics provides a disciplined target-side reading of the historical flow vocabulary, while the canonical source architecture remains a smooth four-dimensional arena carried as primitive debt plus unresolved Φ_{src} . This interpretation neither identifies the source arena with physical spacetime nor pretends that a first-principles bridge has been completed.

Within the flagship exact-closure package, the overview is the front door, the *Exact-Closure Note* is the short anchor, and the present paper is the flow-geometry module that completes the five-paper modular full statement. Its role is to give the $\mathcal{A}$ ether-flow language a disciplined geometric dictionary after the foundations, dynamics, consistency, and relativistic-recovery modules have already fixed the claim boundary. That ordering keeps exact closure first and derivational continuation secondary.

1. Introduction

The active $\mathcal{A}$ ether-flow Interpretation of Relativity sequence fixes three separate points. First, the canonical mainline research architecture uses a smooth four-dimensional source manifold as explicit primitive debt and retains Φ_{src} only as an unresolved abstract source-order or evolution slot. Second, the exact effective gravitational law of the benchmark is general relativity with universal matter coupling. Third, no map between those source and target objects is adopted. The task of this manuscript is therefore to discipline the target-side flow vocabulary without collapsing it into source structure.

That is the task of the present manuscript. The goal is not to replace the metric by a second low-energy object. Nor is the goal to claim a completed microphysical derivation of gravity from substrate variables. The goal is to identify the correct relativistic location of the $\mathcal{A}$ ether-flow idea inside exact closure.

Framework and claim boundary. $\mathcal{A}$ ether / $\mathcal{A}$ ether-flow framework denotes the broader $\mathcal{A}$ ether / $\mathcal{A}$ ether-flow research interpretation. Its source package does not adopt a physical metric, causal cone, clock, slice embedding, matter sector, dynamics, or physical meaning for Φ_{src} . The familiar experiential-space, S-time, and deeper-motion phrases remain interpretive aims. Within that framework, $\mathcal{A}$ ether-flow Interpretation of Relativity denotes the exact-closure benchmark in which

the effective gravitational dynamics are adopted to be exactly Einsteinian with universal matter coupling. In the active sequence, *adoption* means use of that established target-side dynamics without claiming source derivation, while *derivation* is reserved for a first-principles recovery from explicit source variables.

The key discipline is this. A viable flow interpretation of GR cannot identify all gravitational phenomena with one naive image such as a literal vortex. Static spherical gravity exists without vorticity. Cosmological expansion exists without local swirl. Frame dragging does involve rotational structure, but it is not the whole content of gravitation. The correct relativistic move is therefore to interpret $\mathcal{A}$ ether-flow through the full kinematics of a timelike congruence rather than through one over-literal metaphor. The classical language for doing so is the congruence and 1+3 covariant machinery of GR [3, 4].

The sequence role is therefore explicit. After the overview, the short exact-closure anchor, and the foundations, dynamics, consistency, and relativistic-recovery modules, the present manuscript closes the modular full statement by fixing the formal geometric dictionary of the $\mathcal{A}$ ether-flow interpretation. It does not reopen the effective law and it does not move derivational continuation ahead of the exact benchmark already adopted.

2. Flow Geometry in $\mathcal{A}$ ether-flow Interpretation of Relativity

Definition 1 (Flow geometry in $\mathcal{A}$ ether-flow Interpretation of Relativity). *Flow geometry in $\mathcal{A}$ ether-flow Interpretation of Relativity is the pair $(g_{\mu\nu}, u^\mu)$, where $g_{\mu\nu}$ is the exact GR metric adopted by the theory and u^μ is a future-directed unit timelike congruence satisfying*

$$u^\mu u_\mu = -c^2. \quad (1)$$

The congruence provides the observer-based target-side kinematic dictionary by which the historical $\mathcal{A}$ ether-flow vocabulary is read inside the exact relativistic geometry. This definition does not identify u^μ with Φ_{src} or construct either object from the other.

Remark 1. *The congruence is not a second field equation added to GR. It is an admissible, generally non-unique observer field defined on the exact metric spacetime. Different physical situations may call for different natural congruences, but they all live inside the same target metric geometry. No source-level derivation of a preferred choice and no bridge from Φ_{src} is claimed here.*

This point fixes the scientific status of the manuscript. The paper is not proposing a bimetric theory, a preferred-frame matter coupling, or an extra propagating vector sector. It is specifying how the $\mathcal{A}$ ether-flow language may be made mathematically disciplined while the effective law remains exactly Einsteinian.

Proposition 1 (Single-law interpretation). *In $\mathcal{A}$ ether-flow Interpretation of Relativity, the entire flow-geometry interpretation is subordinate to one effective law,*

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad (2)$$

with universal matter coupling to $g_{\mu\nu}$. The congruence u^μ adds interpretation, not a second low-energy dynamics.

2.1. Congruence Choice and Scope

The observer field u^μ is an admissible congruence choice inside the adopted target metric geometry, not a unique field derived from source dynamics. At current scope there is no source-to-congruence map: one chooses the observer family that fits the target-side physical question while remaining inside the same exact relativistic spacetime [3, 4].

Four examples fix the intended use of the dictionary.

1. In static spacetimes, static observer congruences are natural when the goal is to describe gravitational support and redshift.
2. In FLRW cosmology, comoving congruences are natural when the goal is to describe homogeneous expansion.
3. In stationary axisymmetric settings, zero-angular-momentum-observer-like (ZAMO-like) congruences are natural when the goal is to isolate frame dragging and rotational structure.
4. In local inertial or detector problems, freely falling congruences are natural when the goal is to describe local SR recovery, geodesic deviation, and wave strain.

None of these choices is claimed here to be uniquely selected by substrate microphysics. The present manuscript supplies the disciplined dictionary once an admissible observer family has been specified.

3. Observer Rest Space and Local Decomposition

Given a unit timelike congruence u^μ , define the observer rest-space projector by

$$h_{\mu\nu} = g_{\mu\nu} + \frac{1}{c^2} u_\mu u_\nu. \quad (3)$$

Then

$$h_{\mu\nu} u^\nu = 0, \quad h^\mu{}_\alpha h^\alpha{}_\nu = h^\mu{}_\nu, \quad (4)$$

so $h_{\mu\nu}$ defines the spatial geometry measured by observers whose four-velocity is u^μ .

Any vector X^μ decomposes uniquely into temporal and spatial parts relative to the congruence:

$$X^\mu = -\frac{u_\nu X^\nu}{c^2} u^\mu + h^\mu{}_\nu X^\nu. \quad (5)$$

Accordingly, any timelike worldline tangent w^μ may be written as

$$w^\mu = \Gamma (u^\mu + v^\mu), \quad u_\mu v^\mu = 0, \quad (6)$$

where v^μ is the observer-measured spatial velocity and

$$\Gamma = -\frac{u_\mu w^\mu}{c^2} = \frac{1}{\sqrt{1 - v^2/c^2}}, \quad v^2 = h_{\mu\nu} v^\mu v^\nu. \quad (7)$$

Proposition 2 (Local relativistic split). *The pair $(u^\mu, h_{\mu\nu})$ gives the exact observer-based decomposition of relativistic physics in *Aether-flow Interpretation of Relativity*: u^μ fixes the local time direction, $h_{\mu\nu}$ fixes the local rest space, and (7) yields the ordinary Lorentz factor of relative motion.*

This is already enough to see why the $\mathcal{A}$ ether-flow language need not threaten SR. The congruence identifies a local observer family, while the local metric split still gives the ordinary relativistic relation between measured speed and Lorentz factor.

4. Target-Side Kinematic Flow Dictionary

Define the spatially projected derivative

$$\mathcal{D}_\mu \equiv h_\mu{}^\nu \nabla_\nu. \quad (8)$$

The fundamental kinematic quantities of the congruence are then:

$$a_\mu \equiv u^\nu \nabla_\nu u_\mu, \quad (9)$$

$$\theta \equiv \nabla_\mu u^\mu, \quad (10)$$

$$\sigma_{\mu\nu} \equiv h_{(\mu}{}^\alpha h_{\nu)}{}^\beta \nabla_\alpha u_\beta - \frac{1}{3} \theta h_{\mu\nu}, \quad (11)$$

$$\omega_{\mu\nu} \equiv h_{[\mu}{}^\alpha h_{\nu]}{}^\beta \nabla_\alpha u_\beta. \quad (12)$$

These satisfy

$$\sigma_{\mu\nu} u^\nu = 0, \quad \omega_{\mu\nu} u^\nu = 0, \quad \sigma^\mu{}_\mu = 0. \quad (13)$$

With these definitions, the covariant derivative of the congruence decomposes as

$$\nabla_\mu u_\nu = \frac{1}{3} \theta h_{\mu\nu} + \sigma_{\mu\nu} + \omega_{\mu\nu} - \frac{1}{c^2} u_\mu a_\nu. \quad (14)$$

Proposition 3 (Flow-sector map). *In the exact $\mathcal{A}$ ether-flow Interpretation of Relativity line, the congruence kinematics provide the disciplined flow interpretation:*

1. θ describes local volume expansion or contraction of the observer congruence.
2. a_μ describes the proper acceleration structure of non-geodesic observer families and governs static redshift support.
3. $\omega_{\mu\nu}$ describes swirl, twist, and frame-dragging structure.
4. $\sigma_{\mu\nu}$ describes anisotropic distortion, tidal deformation, and wave-induced strain.

This proposition is the key conceptual payoff of the paper. It shows how the flow language becomes mathematically useful once it is distributed across the full congruence kinematics rather than collapsed into one slogan.

4.1. Raychaudhuri Control

The evolution of the expansion is governed by the Raychaudhuri equation [3]

$$u^\mu \nabla_\mu \theta = -\frac{1}{3} \theta^2 - \sigma_{\mu\nu} \sigma^{\mu\nu} + \omega_{\mu\nu} \omega^{\mu\nu} - \nabla_\mu a^\mu - \frac{1}{c^2} R_{\mu\nu} u^\mu u^\nu. \quad (15)$$

This formula is important because it prevents the flow interpretation from becoming vague. Focusing, defocusing, local support against free fall, and the effect of matter on congruence evolution are all carried by definite geometric invariants.

4.2. Frobenius Criterion

The vorticity sector is equally controlled. The congruence is hypersurface orthogonal if and only if

$$\omega_{\mu\nu} = 0. \quad (16)$$

Thus swirl is not a metaphorical add-on. It has a sharp geometric meaning: failure of the observer flow lines to be orthogonal to a family of spatial hypersurfaces.

5. Expansion as Ordered Volume Change

The ontology of observed expansion is sharpened immediately by the expansion scalar θ . For any infinitesimal rest-space volume element V carried by the congruence,

$$\frac{1}{V} \frac{dV}{d\tau} = \theta. \quad (17)$$

This is the exact relativistic meaning of local ordered growth or contraction of the observer flow.

In a homogeneous and isotropic cosmological geometry with comoving congruence u^μ , one has

$$a_\mu = 0, \quad \sigma_{\mu\nu} = 0, \quad \omega_{\mu\nu} = 0, \quad \theta = 3H, \quad (18)$$

where H is the Hubble parameter. The observed expansion of cosmology is therefore represented, in the congruence language, by nonzero θ rather than by motion into an external container.

This is the comoving-FLRW example from the congruence-choice discussion above.

Remark 2. *This clarifies only the target-side dictionary. Congruence expansion inside the exact relativistic geometry is not evidence that Φ_{src} has an adopted evolution law or that observed expansion has been derived from source structure.*

5.1. Dark Energy and Congruence Defocusing

The same congruence language accommodates late-time accelerated expansion without leaving exact closure. For the comoving FLRW congruence, the reduction of Raychaudhuri's equation is

$$u^\mu \nabla_\mu \theta + \frac{1}{3} \theta^2 = -4\pi G \left(\rho + \frac{3p}{c^2} \right) + \Lambda c^2, \quad (19)$$

or, equivalently,

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2} \right) + \frac{\Lambda c^2}{3}. \quad (20)$$

Thus positive Λ , or more generally any GR-compatible component satisfying $\rho + 3p/c^2 < 0$, counteracts focusing and drives accelerated growth of the comoving observer congruence [5, 6, 7].

This is the correct exact-closure location of dark energy in the target-side Æther-flow dictionary. Dark energy is not a literal fluid wind pushing galaxies through a pre-existing space. It is the

standard GR cosmological term or stress tensor that changes the evolution of the same congruence expansion $\theta = 3H$. No inference from this target-side acceleration to source Φ_{src} , a source law, or a metric derivation is licensed.

6. Static Gravity and Redshift as Acceleration Structure

The next step is equally important for physical interpretation. Static gravity is not fundamentally a vorticity effect. It is encoded, for the natural static observer congruence, by acceleration structure together with tidal curvature.

Consider a static line element

$$ds^2 = -N^2(\mathbf{x}) c^2 dt^2 + \gamma_{ij}(\mathbf{x}) dx^i dx^j, \quad (21)$$

and choose the static observer congruence

$$u^\mu = N^{-1} \delta^\mu_t. \quad (22)$$

For this congruence,

$$\theta = 0, \quad \sigma_{\mu\nu} = 0, \quad \omega_{\mu\nu} = 0, \quad (23)$$

while the acceleration is

$$a_\mu = \mathcal{D}_\mu \ln N. \quad (24)$$

Thus the static observers are generally not geodesic even though the spacetime may be stationary and non-rotating. The proper acceleration needed to hold station in the gravitational field is encoded directly in the gradient of the lapse.

This is the static-observer example from the congruence-choice discussion above.

In the weak-field limit,

$$N(\mathbf{x}) = 1 + \frac{\Phi(\mathbf{x})}{c^2} + \mathcal{O}(c^{-4}), \quad (25)$$

so

$$a_i = \frac{1}{c^2} \partial_i \Phi + \mathcal{O}(c^{-4}), \quad (26)$$

or equivalently

$$c^2 a_i = \partial_i \Phi + \mathcal{O}(c^{-2}). \quad (27)$$

This is the relativistic statement that the proper support acceleration of the static observer family reproduces the Newtonian gravitational potential gradient.

The same lapse factor controls gravitational redshift. For stationary observers with $dx^i = 0$,

$$d\tau = N(\mathbf{x}) dt, \quad (28)$$

so for light emitted at A and received at B ,

$$\frac{\nu_B}{\nu_A} = \frac{N(\mathbf{x}_A)}{N(\mathbf{x}_B)}. \quad (29)$$

In the weak-field regime this becomes

$$\frac{\nu_B - \nu_A}{\nu_A} = \frac{\Phi_A - \Phi_B}{c^2} + \mathcal{O}(c^{-4}). \quad (30)$$

Proposition 4 (Static-gravity interpretation). *For the natural static observer congruence, the flow-geometry interpretation of gravity is carried primarily by congruence acceleration a_μ and curvature-induced tidal structure, not by vorticity.*

This proposition is essential. It shows why the language of a universal vortex cannot serve as the whole theory. Static gravitational fields are physically real even when $\omega_{\mu\nu} = 0$.

7. Swirl, Rotation, and Frame Dragging

The vortex intuition does have a legitimate relativistic home, but only in the rotational sector. When $\omega_{\mu\nu} \neq 0$, the congruence has local twist or swirl. This is the sector in which rotational gravity and frame dragging belong.

In stationary axisymmetric settings, the natural illustrative choice is often a zero-angular-momentum-observer-like (ZAMO-like) congruence. More generally, in a stationary but non-static geometry with nontrivial time-space metric components g_{0i} , the congruence of observers held at fixed spatial coordinates generally has nonzero vorticity. In the weak-field regime, $\omega_{\mu\nu}$ is controlled by the antisymmetric spatial derivatives of those time-space components. Thus the rotational part of the metric carries a genuine swirl content in the congruence description.

Proposition 5 (Rotational sector). *Within the $\mathcal{A}$ ether-flow interpretation of $\mathcal{A}$ ether-flow Interpretation of Relativity, vorticity $\omega_{\mu\nu}$ is the correct mathematical carrier of swirl, rotational dragging, and local failure of hypersurface orthogonality.*

This is the disciplined way to keep the user's physical intuition without distorting the mathematics. Swirl is real, but it is not the whole of gravity. It is the rotational sector of the flow geometry.

The same point may be stated operationally. In the presence of frame dragging, gyroscope axes, local inertial directions, and orbit precession respond to the rotational structure of spacetime. In the congruence picture, that response is read through the vorticity sector rather than through the static acceleration sector.

8. Shear, Tidal Gravity, and Wave Strain

The tracefree symmetric part of the flow gradient is the shear tensor $\sigma_{\mu\nu}$. It governs anisotropic distortion of an observer bundle and therefore gives the correct home for tidal stretching and squeezing.

The curvature content measured by the observer field is encoded in the electric part of the Weyl tensor,

$$E_{\mu\nu} \equiv \frac{1}{c^2} C_{\mu\alpha\nu\beta} u^\alpha u^\beta, \quad (31)$$

which is spatial, symmetric, and tracefree in vacuum. For nearby freely falling observers separated by a spatial vector ξ^μ , the vacuum geodesic-deviation law reduces to

$$\frac{D^2 \xi^\mu}{D\tau^2} = -E^\mu{}_\nu \xi^\nu. \quad (32)$$

Thus the tidal content of gravity is registered directly as differential stretching and compression in the observer rest space.

Proposition 6 (Tidal sector). *In the $\mathcal{A}$ ether-flow interpretation of $\mathcal{A}$ ether-flow Interpretation of Relativity, shear $\sigma_{\mu\nu}$ and the tidal tensor $E_{\mu\nu}$ together provide the correct mathematical reading of anisotropic gravitational distortion.*

This is again important for conceptual clarity. What people loosely call the bending of space and time is not one undifferentiated effect. In the flow-geometry description, part of it appears as static acceleration, part as rotational vorticity, and part as tidal shear driven by curvature.

The gravitational-wave sector fits naturally into the same picture. In transverse-traceless form around local Minkowski space,

$$ds^2 = -c^2 dt^2 + \left(\delta_{ij} + h_{ij}^{\text{TT}} \right) dx^i dx^j, \quad \partial_i h_{ij}^{\text{TT}} = 0, \quad h_{ii}^{\text{TT}} = 0. \quad (33)$$

The wave does not create a second low-energy propagation law. It is already part of the exact GR metric sector. For freely falling detector congruences, the fourth example from the scope discussion above, it appears as oscillatory tidal deformation and therefore as time-dependent shear and geodesic deviation in the observer rest space.

9. Local Special Relativity as the Local Inertial Limit

The local relation to SR can now be stated sharply. Freely falling congruences provide the cleanest local inertial realization of the dictionary. At every event p , one may choose local inertial coordinates such that

$$g_{\mu\nu}(p) = \eta_{\mu\nu}, \quad \partial_\alpha g_{\mu\nu}(p) = 0. \quad (34)$$

If the observer congruence is chosen so that

$$u^\mu(p) = (c, 0, 0, 0) \quad (35)$$

in those coordinates, then the rest-space projector reduces at p to the ordinary Euclidean spatial projector and the relative-motion decomposition (6) becomes the usual SR split.

Corollary 1 (Local SR recovery). *For any event p and any unit timelike observer $u^\mu(p)$, the flow geometry of $\mathcal{A}$ ether-flow Interpretation of Relativity reduces locally to standard special-relativistic kinematics with Lorentz factor*

$$\Gamma = \frac{1}{\sqrt{1 - v^2/c^2}}. \quad (36)$$

This is the precise sense in which the $\mathcal{A}$ ether / $\mathcal{A}$ ether-flow interpretation does not deny SR. The theory does not add a measurable preferred-frame correction to local inertial physics in the exact-closure theory. Instead, it provides a deeper reading of why embedded observers always recover local Minkowski order inside the exact relativistic geometry.

10. Interpretive Dictionary and Explicit Nonclaims

The flow-geometry manuscript fixes the interpretive/congruence dictionary for the active $\mathcal{A}$ ether-flow Interpretation of Relativity line. Its positive contribution is the following.

1. It gives the historical $\mathcal{A}$ ether-flow vocabulary a congruence-based target-side interpretation without identifying the congruence with Φ_{src} , a second low-energy gravitational field, or a source dynamics.
2. It provides the exact observer rest-space split $(u^\mu, h_{\mu\nu})$ appropriate to that interpretation.
3. It assigns distinct physical roles to expansion, acceleration, vorticity, and shear.
4. It shows that static gravity, redshift, swirl, frame dragging, tidal distortion, gravitational-wave strain, and local SR can all be read within one geometric dictionary.
5. It clarifies that vortex language is at most a statement about the vorticity sector and cannot be the whole account of gravity.

The manuscript also has explicit nonclaims that protect the benchmark package discipline.

1. It does not alter the one operative metric, the universal matter coupling rule, or the exact GR observer-level dynamics already fixed elsewhere in the active sequence.
2. It does not introduce an independent low-energy preferred-frame field, a second universal causal cone, or a new radiative gravitational sector.
3. It does not claim a first-principles source derivation of the congruence dictionary or of Einsteinian gravity from explicit source variables.
4. It does not type Φ_{src} , identify the source arena with physical spacetime, or supply a source-to-congruence bridge.
5. It does not replace the dynamics or relativistic-recovery modules; it presupposes them and sharpens the ontology only at the level of interpretation.

Accordingly, the present contribution is interpretive sharpening through formal geometry. It is a disciplined dictionary for reading the ontology inside exact closure, not a new low-energy dynamics.

11. Conclusion

The clean way to make the $\mathcal{A}$ ether and $\mathcal{A}$ ether-flow interpretation useful inside $\mathcal{A}$ ether-flow Interpretation of Relativity is not to oppose GR or SR or to identify source and target objects by fiat. It is to read the exact relativistic geometry through a disciplined target-side dictionary while keeping the source bridge open. The congruence-based formalism provides that dictionary.

In this target-side language, the historical flow idea is distributed across structured relativistic quantities. Expansion θ captures ordered growth and cosmological volume change. Acceleration a_μ captures static gravitational support and redshift structure. Vorticity $\omega_{\mu\nu}$ captures swirl, rotation, and frame dragging. Shear $\sigma_{\mu\nu}$ captures anisotropic tidal distortion and wave strain. Local special relativity appears as the local inertial limit of the same geometry. None of these target-side quantities supplies the missing semantics or dynamics of Φ_{src} .

That is the appropriate status of the manuscript. It does not claim that GR has been derived from source microphysics or that the selected source arena is physical spacetime. It claims only that, once exact closure is adopted, the $\mathcal{A}$ ether / $\mathcal{A}$ ether-flow interpretation has a disciplined target-side flow dictionary. A first-principles connection from $(\mathcal{A}, \Phi_{\text{src}})$ to that measured relativistic world remains open.

References

- Einstein, A. (1916). The foundation of the general theory of relativity. *Annalen der Physik*, 49, 769–822.
- Arnowitt, R., Deser, S., & Misner, C. W. (1962). The dynamics of general relativity. In L. Witten (Ed.), *Gravitation: An introduction to current research* (pp. 227–265). Wiley. Reprinted in *General Relativity and Gravitation*, 40, 1997–2027.
- Raychaudhuri, A. (1955). Relativistic cosmology. I. *Physical Review*, 98, 1123–1126.
- Ellis, G. F. R., & van Elst, H. (1999). Cosmological models (Cargèse lectures 1998). In M. Lachièze-Rey (Ed.), *Theoretical and observational cosmology* (NATO Science Series C: Mathematical and Physical Sciences, Vol. 541, pp. 1–116). Kluwer. [arXiv:gr-qc/9812046](https://arxiv.org/abs/gr-qc/9812046)
- Riess, A. G., et al. [Supernova Search Team]. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astronomical Journal*, 116, 1009–1038. [arXiv:astro-ph/9805201](https://arxiv.org/abs/astro-ph/9805201)
- Perlmutter, S., et al. [Supernova Cosmology Project]. (1999). Measurements of Ω and Λ from 42 high-redshift supernovae. *Astrophysical Journal*, 517, 565–586. [arXiv:astro-ph/9812133](https://arxiv.org/abs/astro-ph/9812133)
- Aghanim, N., et al. [Planck Collaboration]. (2020). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641, A6. [arXiv:1807.06209](https://arxiv.org/abs/1807.06209)
- Jacobson, T., & Mattingly, D. (2001). Gravity with a dynamical preferred frame. *Physical Review D*, 64, 024028.
- Jacobson, T., & Mattingly, D. (2004). Einstein-aether waves. *Physical Review D*, 70, 024003.
- Eling, C., & Jacobson, T. (2004). Static post-Newtonian equivalence of general relativity and gravity with a dynamical preferred frame. *Physical Review D*, 69, 064005.
- Jacobson, T. (2010). Extended Hořava gravity and Einstein-aether theory. *Physical Review D*, 81, 101502(R).
- Jacobson, T. (2014). Undoing the twist: The Hořava limit of Einstein-aether theory. *Physical Review D*, 89, 081501(R).
- Jacobson, T., & Speranza, A. J. (2015). Variations on an aethereal theme. *Physical Review D*, 92, 044030.
- Hořava, P. (2009). Quantum gravity at a Lifshitz point. *Physical Review D*, 79, 084008.
- Jacobson, T. (1995). Thermodynamics of spacetime: The Einstein equation of state. *Physical Review Letters*, 75, 1260–1263.
- Visser, M. (2002). Sakharov’s induced gravity: A modern perspective. *Modern Physics Letters A*, 17, 977–992. [arXiv:gr-qc/0204062](https://arxiv.org/abs/gr-qc/0204062)
- Hojman, S. A., Kuchař, K., & Teitelboim, C. (1976). Geometrodynamics regained. *Annals of Physics*, 96, 88–135.
- Deser, S. (1970). Self-interaction and gauge invariance. *General Relativity and Gravitation*, 1, 9–18. Reprinted with 2004 note at [arXiv:gr-qc/0411023](https://arxiv.org/abs/gr-qc/0411023)
- Lovelock, D. (1971). The Einstein tensor and its generalizations. *Journal of Mathematical Physics*, 12, 498–501.
- Barcelo, C., Liberati, S., & Visser, M. (2011). Analogue gravity. *Living Reviews in Relativity*, 14, 3.
- Sakharov, A. D. (1968). Vacuum quantum fluctuations in curved space and the theory of gravitation. *Soviet Physics Doklady*, 12, 1040–1041. Translated from *Doklady Akademii Nauk SSSR*, 177, 70–71.