# The Æther-flow Interpretation of Relativity

Primitive-Reservoir Observer-Reduction
Kernel-Dynamics-Generating Substrate
Pre-Selector Pre-Kernel Dynamics
from Pre-Kernel-Dynamics-Generating
Substrate Pre-Selector Proto-Kernel Dynamics


**Abstract**

The current primitive-reservoir observer-reduction frontier already sharpened the live burden inside the current pre-selector pre-kernel line: the current kernel-dynamics-generating substrate pre-selector pre-kernel dynamics package $\mathfrak{G}^{\mathrm{presel,preker}}$ is the live unexplained datum, while the larger current kernel package $\mathfrak{F}^{\mathrm{presel,ker}}$ and all downstream observer-relevant outputs have already been spent. The active routing note therefore demands the next honest move below the current frontier: either derive or justify $\mathfrak{G}^{\mathrm{presel,preker}}$ from still more primitive substrate structure in a way that genuinely changes benchmark-facing status, or prove a sharper theorem-level collapse strictly inside it.

The present note takes the direct derivation route. For each sector it introduces pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics $\mathfrak{H}^{\mathrm{presel,proker}}$: a compact convex proto-kernel body over the already fixed current pre-kernel-body geometry, an affine proto-kernel projection to that geometry, a nonnegative smooth proto-kernel functional with unique inner fiber minimizers, a current-kernel-scale minimizing law forcing the current pre-kernel minimizer lift, a selector-scale minimizing law forcing the current pre-kernel composed minimizer, and a benchmark-shell-scale minimizing shell whose projection fixes the current exact pre-kernel shell. The current pre-kernel functional is then no longer an independent primitive stopping point. It is the effective minimum value of the deeper proto-kernel law on the inner proto-kernel fibers, the current exact pre-kernel shell is the projected exact proto-kernel shell over the already fixed selector benchmark shell, and the current-kernel-scale, selector-scale, and benchmark-shell-scale pre-kernel minimizers are all outer projections of unique deeper proto-kernel minimizers.

The benchmark boundary remains fixed throughout. The one operative metric is still exactly $g^\sharp$, the ordinary matter route is unchanged, there is no second operative metric, the low-energy matter law is not enlarged, and no stronger promotion language is licensed. The positive result is therefore a genuine benchmark-facing gain below the current pre-selector pre-kernel frontier: the remaining honest burden is no longer the current pre-selector pre-kernel dynamics as such, but the deeper pre-selector proto-kernel dynamics that canonically generate them, or a still smaller collapse strictly inside that deeper package.


## 1. Introduction

The active derivational program of Æther-flow Interpretation of Relativity again reaches a point where depth alone is not enough. The current active-primary theorem already showed that the current pre-selector pre-kernel package is not the primitive stopping point at present scope. Once

the kernel-dynamics-generating substrate pre-selector pre-kernel dynamics package $\mathfrak{G}^{\mathrm{presel,preker}}$ is fixed, the current kernel functional, current exact kernel shell, and current kernel-scale and benchmark-shell-scale minimizing data are forced canonically from it. The routing consequence is then immediate. If the project goal is to move the same benchmark one-metric package toward a disciplined first-principles recovery, the next honest benchmark-facing manuscript must either derive or justify $\mathfrak{G}^{\mathrm{presel,preker}}$ itself from still more primitive substrate structure, or prove a genuinely smaller theorem-level collapse strictly inside it.

The present note takes the first route. It does not claim that no sharper internal collapse will ever be available. It records only that no smaller observer-relevant datum has yet been isolated inside the current pre-kernel package without already presupposing that package's law. The disciplined next move is therefore to spend $\mathfrak{G}^{\mathrm{presel,preker}}$ itself. Concretely, the note introduces a deeper proto-kernel layer below the current pre-kernel package and proves that the current pre-kernel functional, the current exact pre-kernel shell, the current pre-kernel minimizer lift over the current kernel body, the current selector-scale pre-kernel composed minimizer, and the current benchmark-shell-scale full pre-kernel minimizer are all canonical outputs of that deeper proto-kernel law.

**Framework and claim boundary.** Æther / Æther-flow framework denotes the broader Æther / Æther-flow framework built on the claims that the Æther is the underlying four-dimensional substrate of reality, the Æther-flow is its intrinsic ordered motion, observed three-dimensional space is the local experiential slice of that deeper substrate, S-time is the experienced order of change arising from matter, light, and the Æther-flow, and observed expansion is the three-dimensional appearance of deeper four-dimensional ordered motion. Within that framework, Æther-flow Interpretation of Relativity denotes the exact-closure theory in which the effective gravitational dynamics are adopted to be exactly Einsteinian with universal matter coupling. In the active sequence, *adoption* means use of that established relativistic dynamics without claiming substrate derivation, while *derivation* is reserved for a first-principles recovery from explicit substrate variables.

The present note stays strictly inside that boundary. It does not alter the benchmark exact-closure package. It does not introduce a second operative metric or a new low-energy matter law. It only strengthens the deeper origin-side explanation of why the current pre-selector pre-kernel package used by the immediately preceding theorem should itself hold.

## 2. Fixed Inputs

**Definition 1** (Recorded primitive continuation data)**.** *Fix the following continuation data from the active primitive-reservoir observer-reduction line.*

1. *For each sector label* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$*, a primitive forcing shell*

$$\mathcal{I}_\sigma^{\mathrm{for}}.$$

2. *The product shell*

$$\mathcal{S}_{\mathrm{prim}} \equiv \mathcal{I}_{\mathrm{car}}^{\mathrm{for}} \times \mathcal{I}_{\mathrm{cur}}^{\mathrm{for}} \times \mathcal{I}_{\mathrm{hom}}^{\mathrm{for}}.$$

3. *The recorded metric and ordinary-matter outputs*

$$\mathscr{G}_{\mathrm{out}} : \mathcal{S}_{\mathrm{prim}} \to \{\textit{observer metrics}\}, \qquad \mathscr{T}_{\mathrm{out}} : \mathcal{S}_{\mathrm{prim}} \times \Psi \to \{\textit{observer stress tensors}\},$$

*satisfying*

$$\mathscr{G}_{\mathrm{out}}(\mathbf{w}) = g^{\sharp}, \qquad \mathscr{T}_{\mathrm{out}}(\mathbf{w}, \psi) = T^{\mathrm{matter}}_{\mu\nu}[g^{\sharp}, \psi] \tag{1}$$

*for every* $\mathbf{w} \in \mathcal{S}_{\mathrm{prim}}$ *and every matter configuration* $\psi \in \Psi$.

**Definition 2** (Recorded selector benchmark base)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$. *The active line already fixes the following benchmark-facing selector data:*

1. *a finite-dimensional Euclidean selector state space*

$$\mathcal{D}^{\mathrm{yoc,sel}}_{\sigma};$$

2. *a compact convex selector body*

$$\mathcal{B}^{\mathrm{yoc,sel}}_{\sigma} \subset \mathcal{D}^{\mathrm{yoc,sel}}_{\sigma}$$

   *with nonempty interior;*

3. *a nonnegative smooth selector functional*

$$\mathscr{L}^{\mathrm{yoc,sel}}_{\sigma} : \mathcal{D}^{\mathrm{yoc,sel}}_{\sigma} \to [0, \infty);$$

4. *a closed embedded exact selector shell*

$$\mathcal{Z}^{\mathrm{yoc,sel}}_{\sigma} \subset \mathcal{B}^{\mathrm{yoc,sel}}_{\sigma};$$

5. *benchmark faithfulness, meaning preservation of the benchmark outputs* (1)*, no second operative metric, no enlarged low-energy matter law, and use of this selector benchmark base as already fixed upstream data on the active line.*

**Definition 3** (Recorded current selector-body geometry)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$. *The active line already fixes the following current selector-body geometry:*

1. *a finite-dimensional Euclidean current selector state space*

$$\mathcal{F}^{\mathrm{presel,sel}}_{\sigma};$$

2. *a compact convex current selector body*

$$\mathcal{B}^{\mathrm{presel,sel}}_{\sigma} \subset \mathcal{F}^{\mathrm{presel,sel}}_{\sigma}$$

   *with nonempty interior;*

3. *an affine surjection*

$$\Pi^{\mathrm{presel,sel}}_{\sigma} : \mathcal{F}^{\mathrm{presel,sel}}_{\sigma} \to \mathcal{B}^{\mathrm{yoc,sel}}_{\sigma}$$

   *satisfying*

$$\Pi^{\mathrm{presel,sel}}_{\sigma}(\mathcal{B}^{\mathrm{presel,sel}}_{\sigma}) = \mathcal{B}^{\mathrm{yoc,sel}}_{\sigma};$$

4. *for each* $y \in \mathcal{B}^{\mathrm{yoc,sel}}_{\sigma}$*, the current selector fiber*

$$\mathcal{G}^{\mathrm{presel,sel}}_{\sigma}(y) \equiv \left(\Pi^{\mathrm{presel,sel}}_{\sigma}\right)^{-1}(y) \cap \mathcal{B}^{\mathrm{presel,sel}}_{\sigma},$$

*which is required to be nonempty, compact, and convex.*

**Definition 4** (Recorded current kernel-body geometry)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$*. The active line already fixes the following current kernel-body geometry below the current selector-body geometry:*

1. *a finite-dimensional Euclidean current kernel state space*

$$\mathcal{H}_\sigma^{\mathrm{presel,ker}};$$

2. *a compact convex current kernel body*

$$\mathcal{K}_\sigma^{\mathrm{presel,ker}} \subset \mathcal{H}_\sigma^{\mathrm{presel,ker}}$$

   *with nonempty interior;*

3. *an affine surjection*

$$\Lambda_\sigma^{\mathrm{presel,ker}} : \mathcal{H}_\sigma^{\mathrm{presel,ker}} \to \mathcal{F}_\sigma^{\mathrm{presel,sel}}$$

   *satisfying*

$$\Lambda_\sigma^{\mathrm{presel,ker}}(\mathcal{K}_\sigma^{\mathrm{presel,ker}}) = \mathcal{B}_\sigma^{\mathrm{presel,sel}};$$

4. *for each* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*, the current kernel fiber*

$$\mathcal{J}_\sigma^{\mathrm{presel,ker}}(u) \equiv (\Lambda_\sigma^{\mathrm{presel,ker}})^{-1}(u) \cap \mathcal{K}_\sigma^{\mathrm{presel,ker}},$$

   *which is required to be nonempty, compact, and convex.*

**Definition 5** (Recorded current pre-kernel-body geometry)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$*. The active line already fixes the following current pre-kernel-body geometry below the current kernel-body geometry:*

1. *a finite-dimensional Euclidean current pre-kernel state space*

$$\mathcal{N}_\sigma^{\mathrm{presel,preker}};$$

2. *a compact convex current pre-kernel body*

$$\mathcal{P}_\sigma^{\mathrm{presel,preker}} \subset \mathcal{N}_\sigma^{\mathrm{presel,preker}}$$

   *with nonempty interior;*

3. *an affine surjection*

$$\Upsilon_\sigma^{\mathrm{presel,preker}} : \mathcal{N}_\sigma^{\mathrm{presel,preker}} \to \mathcal{H}_\sigma^{\mathrm{presel,ker}}$$

   *satisfying*

$$\Upsilon_\sigma^{\mathrm{presel,preker}}(\mathcal{P}_\sigma^{\mathrm{presel,preker}}) = \mathcal{K}_\sigma^{\mathrm{presel,ker}};$$

4. *for each* $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$*, the current pre-kernel fiber*

$$\mathcal{U}_\sigma^{\mathrm{presel,preker}}(w) \equiv (\Upsilon_\sigma^{\mathrm{presel,preker}})^{-1}(w) \cap \mathcal{P}_\sigma^{\mathrm{presel,preker}},$$

   *which is required to be nonempty, compact, and convex.*

**Definition 6** (Current pre-selector pre-kernel-dynamics target)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$ *and the recorded current selector-body, kernel-body, and pre-kernel-body geometries of Definitions 3, 4, and 5. A* benchmark-faithful kernel-dynamics-generating substrate pre-selector pre-kernel dynamics package $\mathfrak{G}^{\mathrm{presel,preker}}$ *on that geometry consists of the following data.*

1. *A nonnegative smooth pre-kernel functional*

$$\mathscr{S}_\sigma^{\mathrm{presel,preker}} : \mathcal{N}_\sigma^{\mathrm{presel,preker}} \to [0, \infty)$$

*such that, for every* $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$*, the restriction of* $\mathscr{S}_\sigma^{\mathrm{presel,preker}}$ *to* $\mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)$ *is strictly convex and attains a unique minimizer. The resulting minimizer depends smoothly on* $w$*; write it as*

$$\mathfrak{s}_\sigma^{\mathrm{presel,preker}} : \mathcal{K}_\sigma^{\mathrm{presel,ker}} \to \mathcal{P}_\sigma^{\mathrm{presel,preker}}.$$

*The induced kernel functional*

$$\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}} : \mathcal{H}_\sigma^{\mathrm{presel,ker}} \to [0, \infty)$$

*is required to be smooth on the ambient current kernel state space and to satisfy*

$$\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) \equiv \mathscr{S}_\sigma^{\mathrm{presel,preker}}\big(\mathfrak{s}_\sigma^{\mathrm{presel,preker}}(w)\big) = \min_{z \in \mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)} \mathscr{S}_\sigma^{\mathrm{presel,preker}}(z) \qquad \text{for every } w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

*Moreover, for every* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*, the restriction of* $\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}$ *to* $\mathcal{J}_\sigma^{\mathrm{presel,ker}}(u)$ *is strictly convex and attains a unique minimizer depending smoothly on* $u$*; write it as*

$$\widetilde{\mathfrak{r}}_\sigma^{\mathrm{presel,ker}} : \mathcal{B}_\sigma^{\mathrm{presel,sel}} \to \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

2. *For each* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*, define the selector-scale composed pre-kernel fiber*

$$\mathcal{V}_\sigma^{\mathrm{presel,preker}}(u) \equiv \left\{ z \in \mathcal{P}_\sigma^{\mathrm{presel,preker}} : \Lambda_\sigma^{\mathrm{presel,ker}}\big(\Upsilon_\sigma^{\mathrm{presel,preker}}(z)\big) = u \right\}.$$

*The restriction of* $\mathscr{S}_\sigma^{\mathrm{presel,preker}}$ *to* $\mathcal{V}_\sigma^{\mathrm{presel,preker}}(u)$ *is required to attain a unique minimizer depending smoothly on* $u$*; write it as*

$$\mathfrak{t}_\sigma^{\mathrm{presel,preker}} : \mathcal{B}_\sigma^{\mathrm{presel,sel}} \to \mathcal{P}_\sigma^{\mathrm{presel,preker}}.$$

*Define the induced outer selector functional*

$$\widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}} : \mathcal{F}_\sigma^{\mathrm{presel,sel}} \to [0, \infty)$$

*by*

$$\widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(u) \equiv \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}\big(\widetilde{\mathfrak{r}}_\sigma^{\mathrm{presel,ker}}(u)\big) = \min_{w \in \mathcal{J}_\sigma^{\mathrm{presel,ker}}(u)} \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) \qquad \text{for every } u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}.$$

*This function is required to be smooth on the ambient current selector state space, and for every* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, its restriction to* $\mathcal{G}_\sigma^{\mathrm{presel,sel}}(y)$ *is required to be strictly convex and to attain a unique minimizer depending smoothly on* $y$*.*

3. *For each* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, the restriction of* $\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}$ *to the current composed kernel fiber*

$$\mathcal{M}_\sigma^{\mathrm{presel,ker}}(y) = \left\{ w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(w)\big) = y \right\}$$

*is required to attain a unique minimizer depending smoothly on* $y$*; write it as*

$$\widetilde{\mathfrak{m}}_\sigma^{\mathrm{presel,ker}} : \mathcal{B}_\sigma^{\mathrm{yoc,sel}} \to \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

*Define the induced exact selector shell*

$$\widetilde{\mathcal{Z}}_\sigma^{\mathrm{presel,sel}} \equiv \left\{ u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}} : \Pi_\sigma^{\mathrm{presel,sel}}(u) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(u) = \min_{v \in \mathcal{G}_\sigma^{\mathrm{presel,sel}}(\Pi_\sigma^{\mathrm{presel,sel}}(u))} \widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(v) \right\},$$

*and the induced exact kernel shell*

$$\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}} \equiv \left\{ w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(w)\big) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) = \min_{q \in \mathcal{M}_\sigma^{\mathrm{presel,ker}}(\Pi_\sigma^{\mathrm{presel,sel}}(\Lambda_\sigma^{\mathrm{presel,ker}}}\right.$$

*The shell* $\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}$ *is required to be a closed embedded submanifold, and*

$$\Lambda_\sigma^{\mathrm{presel,ker}}\big|_{\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}} : \widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}} \xrightarrow{\cong} \widetilde{\mathcal{Z}}_\sigma^{\mathrm{presel,sel}}$$

*is required to be a diffeomorphism.*

4. *For each* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, define the full pre-kernel fiber*

$$\mathcal{W}_\sigma^{\mathrm{presel,preker}}(y) \equiv \left\{ z \in \mathcal{P}_\sigma^{\mathrm{presel,preker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(\Upsilon_\sigma^{\mathrm{presel,preker}}(z))\big) = y \right\}.$$

*The restriction of* $\mathscr{S}_\sigma^{\mathrm{presel,preker}}$ *to* $\mathcal{W}_\sigma^{\mathrm{presel,preker}}(y)$ *is required to attain a unique minimizer depending smoothly on* $y$*; write it as*

$$\mathfrak{u}_\sigma^{\mathrm{presel,preker}} : \mathcal{B}_\sigma^{\mathrm{yoc,sel}} \to \mathcal{P}_\sigma^{\mathrm{presel,preker}}.$$

*The exact pre-kernel shell*

$$\mathcal{O}_\sigma^{\mathrm{presel,preker}} \equiv \left\{ z \in \mathcal{P}_\sigma^{\mathrm{presel,preker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(\Upsilon_\sigma^{\mathrm{presel,preker}}(z))\big) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \mathscr{S}_\sigma^{\mathrm{presel,preker}}(z) = \right.$$

*is required to be a closed embedded submanifold, and*

$$\Upsilon_\sigma^{\mathrm{presel,preker}}\big|_{\mathcal{O}_\sigma^{\mathrm{presel,preker}}} : \mathcal{O}_\sigma^{\mathrm{presel,preker}} \xrightarrow{\cong} \widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}$$

*is required to be a diffeomorphism.*

5. *Benchmark faithfulness, meaning preservation of the benchmark outputs* (1)*, no second operative metric, no enlarged low-energy matter law, and presentation as the current substrate-side pre-kernel-dynamics package rather than as a replacement benchmark theory.*

**Definition 7** (Recorded downstream chain below the current pre-kernel-dynamics frontier)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$*. The active line already fixes the following downstream data and consequences:*

1. *the current selector-dynamics-generating substrate pre-selector kernel dynamics package* $\mathfrak{F}^{\mathrm{presel,ker}}$*, the current pre-selector-minimizer-section-generating substrate selector dynamics package* $\mathfrak{E}^{\mathrm{presel,sel}}$*, the current observer-relevant pre-selector minimizer-section core* $\mathfrak{D}^{\mathrm{presel,sec}}$*, the current selector package* $\mathfrak{C}^{\mathrm{yoc,sel}}$*, the current observer-relevant yocto-section minimizer-section core* $\mathfrak{C}^{\mathrm{yoc,sec}}$*, the current observer-relevant yocto-section minimizer-image core* $\mathfrak{C}^{\mathrm{yoc,img}}$*, the current observer-relevant zepto-section minimizer-section core* $\mathfrak{B}^{\mathrm{zep,sec}}$*, the current observer-relevant zepto-section minimizer-image core* $\mathfrak{B}^{\mathrm{zep,img}}$*, the current atto-section package* $\mathfrak{A}^{\mathrm{attosec}}$*, the current femto-section package* $\mathfrak{Z}^{\mathrm{femtosec}}$*, the current pico-section package* $\mathfrak{Y}^{\mathrm{picosec}}$*, the current observer-relevant pico-section minimizer-section core* $\mathfrak{Y}^{\mathrm{pic,sec}}$*, the current nano-section package* $\mathfrak{X}^{\mathrm{nanosec}}$*, the current micro-section package* $\mathfrak{W}^{\mathrm{microsec}}$*, the current sub-section package* $\mathfrak{V}^{\mathrm{subsec}}$*, and the previously recorded shell-transport consequences used on the active line;*

2. *the previously recorded fact that, once a benchmark-faithful current pre-kernel package in the sense of Definition 6 is available on the recorded geometry, the current kernel package is recovered, and all of the downstream data listed in item (1) follow unchanged.*

**Remark 1.** *Definitions 1–7 fix the current frontier package and its benchmark boundary. The one operative metric, the ordinary matter route, the selector benchmark base, the current selector-body geometry, the current kernel-body geometry, the current pre-kernel-body geometry, and the already recorded downstream consequences are not reopened here. The present note asks only why the current pre-kernel package itself should arise from still more primitive substrate structure.*

## 3. Pre-Kernel-Dynamics-Generating Substrate Pre-Selector Proto-Kernel Dynamics

**Definition 8** (Pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics)**.** *Fix* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$ *and the recorded current selector-body, kernel-body, and pre-kernel-body geometries of Definitions 3, 4, and 5. A* pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics package $\mathfrak{H}^{\mathrm{presel,proker}}$ *for sector* $\sigma$ *consists of the following data.*

1. *A finite-dimensional Euclidean proto-kernel state space* $\mathcal{R}_\sigma^{\mathrm{presel,proker}}$*, a compact convex proto-kernel body*

$$\mathcal{T}_\sigma^{\mathrm{presel,proker}} \subset \mathcal{R}_\sigma^{\mathrm{presel,proker}}$$

*with nonempty interior, and an affine surjection*

$$\Omega_\sigma^{\mathrm{presel,proker}} : \mathcal{R}_\sigma^{\mathrm{presel,proker}} \to \mathcal{N}_\sigma^{\mathrm{presel,preker}}$$

*satisfying*

$$\Omega_\sigma^{\mathrm{presel,proker}}(\mathcal{T}_\sigma^{\mathrm{presel,proker}}) = \mathcal{P}_\sigma^{\mathrm{presel,preker}}.$$

*For each* $q \in \mathcal{P}_\sigma^{\mathrm{presel,preker}}$*, write*

$$\mathcal{X}_\sigma^{\mathrm{presel,proker}}(q) \equiv \left(\Omega_\sigma^{\mathrm{presel,proker}}\right)^{-1}(q) \cap \mathcal{T}_\sigma^{\mathrm{presel,proker}}.$$

*Each such fiber is required to be nonempty, compact, and convex.*

2. *A nonnegative smooth proto-kernel functional*

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}} : \mathcal{R}_\sigma^{\mathrm{presel,proker}} \to [0,\infty)$$

*such that, for every* $q \in \mathcal{P}_\sigma^{\mathrm{presel,preker}}$*, the restriction of* $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ *to* $\mathcal{X}_\sigma^{\mathrm{presel,proker}}(q)$ *is strictly convex and attains a unique minimizer. The resulting minimizer depends smoothly on* $q$*; write it as*

$$\mathfrak{v}_\sigma^{\mathrm{presel,proker}} : \mathcal{P}_\sigma^{\mathrm{presel,preker}} \to \mathcal{T}_\sigma^{\mathrm{presel,proker}}.$$

*The induced pre-kernel functional*

$$\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}} : \mathcal{N}_\sigma^{\mathrm{presel,preker}} \to [0,\infty)$$

*is required to be smooth on the ambient current pre-kernel state space and to satisfy*

$$\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q) \equiv \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(q)\big) = \min_{z \in \mathcal{X}_\sigma^{\mathrm{presel,proker}}(q)} \mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \qquad \text{for every } q \in \mathcal{P}_\sigma^{\mathrm{presel,pre}}$$

*Moreover, for every* $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$*, the restriction of* $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ *to* $\mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)$ *is strictly convex and attains a unique minimizer depending smoothly on* $w$*; write it as*

$$\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}} : \mathcal{K}_\sigma^{\mathrm{presel,ker}} \to \mathcal{P}_\sigma^{\mathrm{presel,preker}}.$$

3. *For each* $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$*, define the current-kernel-scale composed proto-kernel fiber*

$$\mathcal{L}_\sigma^{\mathrm{presel,proker}}(w) \equiv \left\{ z \in \mathcal{T}_\sigma^{\mathrm{presel,proker}} : \Upsilon_\sigma^{\mathrm{presel,preker}}\big(\Omega_\sigma^{\mathrm{presel,proker}}(z)\big) = w \right\}.$$

*The restriction of* $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ *to* $\mathcal{L}_\sigma^{\mathrm{presel,proker}}(w)$ *is required to attain a unique minimizer depending smoothly on* $w$*; write it as*

$$\mathfrak{w}_\sigma^{\mathrm{presel,proker}} : \mathcal{K}_\sigma^{\mathrm{presel,ker}} \to \mathcal{T}_\sigma^{\mathrm{presel,proker}}.$$

*Define the induced current kernel functional*

$$\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}} : \mathcal{H}_\sigma^{\mathrm{presel,ker}} \to [0,\infty)$$

*by*

$$\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) \equiv \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}\big(\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w)\big) = \min_{q \in \mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)} \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q) \qquad \text{for every } w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

*This function is required to be smooth on the ambient current kernel state space, and for every* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*, its restriction to* $\mathcal{J}_\sigma^{\mathrm{presel,ker}}(u)$ *is required to be strictly convex and to attain a unique minimizer depending smoothly on* $u$*; write it as*

$$\widetilde{\mathfrak{r}}_\sigma^{\mathrm{presel,ker}} : \mathcal{B}_\sigma^{\mathrm{presel,sel}} \to \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

4. *For each* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*, define the selector-scale composed proto-kernel fiber*

$$\mathcal{A}_\sigma^{\mathrm{presel,proker}}(u) \equiv \left\{ z \in \mathcal{T}_\sigma^{\mathrm{presel,proker}} : \Lambda_\sigma^{\mathrm{presel,ker}}\big(\Upsilon_\sigma^{\mathrm{presel,preker}}(\Omega_\sigma^{\mathrm{presel,proker}}(z))\big) = u \right\}.$$

*The restriction of* $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ *to* $\mathcal{A}_\sigma^{\mathrm{presel,proker}}(u)$ *is required to attain a unique minimizer depending smoothly on* $u$*; write it as*

$$\mathfrak{r}_\sigma^{\mathrm{presel,proker}} : \mathcal{B}_\sigma^{\mathrm{presel,sel}} \to \mathcal{T}_\sigma^{\mathrm{presel,proker}}.$$

*Define the induced outer selector functional*

$$\widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}} : \mathcal{F}_\sigma^{\mathrm{presel,sel}} \to [0, \infty)$$

*by*

$$\widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(u) \equiv \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}\big(\widetilde{\mathfrak{r}}_\sigma^{\mathrm{presel,ker}}(u)\big) = \min_{w \in \mathcal{J}_\sigma^{\mathrm{presel,ker}}(u)} \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) \qquad \textit{for every } u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}.$$

*This function is required to be smooth on the ambient current selector state space, and for every* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, its restriction to* $\mathcal{G}_\sigma^{\mathrm{presel,sel}}(y)$ *is required to be strictly convex and to attain a unique minimizer depending smoothly on* $y$*.*

5. *For each* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, the restriction of* $\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}$ *to the current composed kernel fiber*

$$\mathcal{M}_\sigma^{\mathrm{presel,ker}}(y) = \left\{ w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(w)\big) = y \right\}$$

*is required to attain a unique minimizer depending smoothly on* $y$*; write it as*

$$\widetilde{\mathfrak{m}}_\sigma^{\mathrm{presel,ker}} : \mathcal{B}_\sigma^{\mathrm{yoc,sel}} \to \mathcal{K}_\sigma^{\mathrm{presel,ker}}.$$

*Define the induced exact selector shell*

$$\widetilde{\mathcal{Z}}_\sigma^{\mathrm{presel,sel}} \equiv \left\{ u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}} : \Pi_\sigma^{\mathrm{presel,sel}}(u) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(u) = \min_{v \in \mathcal{G}_\sigma^{\mathrm{presel,sel}}(\Pi_\sigma^{\mathrm{presel,sel}}(u))} \widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}(v) \right\},$$

*the induced exact kernel shell*

$$\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}} \equiv \left\{ w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(w)\big) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}(w) = \min_{q \in \mathcal{M}_\sigma^{\mathrm{presel,ker}}(\Pi_\sigma^{\mathrm{presel,sel}}(\Lambda_\sigma^{\mathrm{presel,ker}}} \right.$$

*and the induced exact pre-kernel shell*

$$\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}} \equiv \left\{ q \in \mathcal{P}_\sigma^{\mathrm{presel,preker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(\Upsilon_\sigma^{\mathrm{presel,preker}}(q))\big) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q) = \right.$$
$$p \in \mathcal{W}_\sigma^{\mathrm{presel}}$$

*The shells* $\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}$ *and* $\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}$ *are required to be closed embedded submanifolds, and*

$$\Lambda_\sigma^{\mathrm{presel,ker}}\big|_{\widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}} : \widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}} \xrightarrow{\cong} \widetilde{\mathcal{Z}}_\sigma^{\mathrm{presel,sel}}, \qquad \Upsilon_\sigma^{\mathrm{presel,preker}}\big|_{\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}} : \widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}} \xrightarrow{\cong} \widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}$$

*are required to be diffeomorphisms.*

6. *For each* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*, define the full proto-kernel fiber*

$$\mathcal{C}_\sigma^{\mathrm{presel,proker}}(y) \equiv \left\{ z \in \mathcal{T}_\sigma^{\mathrm{presel,proker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(\Upsilon_\sigma^{\mathrm{presel,preker}}(\Omega_\sigma^{\mathrm{presel,proker}}(z)))\big) = y \right\}.$$

*The restriction of* $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ *to* $\mathcal{C}_\sigma^{\mathrm{presel,proker}}(y)$ *is required to attain a unique minimizer depending smoothly on* $y$*; write it as*

$$\mathfrak{y}_\sigma^{\mathrm{presel,proker}} : \mathcal{B}_\sigma^{\mathrm{yoc,sel}} \to \mathcal{T}_\sigma^{\mathrm{presel,proker}}.$$

*The exact proto-kernel shell*

$$\mathcal{E}_\sigma^{\mathrm{presel,proker}} \equiv \left\{ z \in \mathcal{T}_\sigma^{\mathrm{presel,proker}} : \Pi_\sigma^{\mathrm{presel,sel}}\big(\Lambda_\sigma^{\mathrm{presel,ker}}(\Upsilon_\sigma^{\mathrm{presel,preker}}(\Omega_\sigma^{\mathrm{presel,proker}}(z)))\big) \in \mathcal{Z}_\sigma^{\mathrm{yoc,sel}},\ \mathscr{U}_\sigma^{\mathrm{presel,proker}}(\right.$$

*is required to be a closed embedded submanifold, and*

$$\Omega_\sigma^{\mathrm{presel,proker}}\big|_{\mathcal{E}_\sigma^{\mathrm{presel,proker}}} : \mathcal{E}_\sigma^{\mathrm{presel,proker}} \xrightarrow{\cong} \widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}$$

*is required to be a diffeomorphism.*

7. *Benchmark faithfulness, meaning preservation of the benchmark outputs* (1)*, no second operative metric, no enlarged low-energy matter law, and presentation as a deeper substrate-side source for the current pre-kernel package rather than as a replacement benchmark theory.*

**Remark 2.** *Definition 8 is intentionally written over the already fixed current pre-kernel-body geometry. It does not reopen the benchmark selector base, the current selector-body geometry, the current kernel-body geometry, or the already-spent larger pre-selector and kernel packages as the live burden. Its positive role is narrower: the current pre-kernel functional, exact pre-kernel shell, and current pre-kernel minimizing data are generated as the effective multi-stage minimizing output of a deeper proto-kernel law.*

## 4. Recovery of the Current Pre-Selector Pre-Kernel Dynamics

**Proposition 1** (The inner proto-kernel minimizer lift is well defined)**.** *For every benchmark-faithful proto-kernel package, the minimizer lift* $\mathfrak{y}_\sigma^{\mathrm{presel,proker}}$ *of Definition 8 is a smooth section of* $\Omega_\sigma^{\mathrm{presel,proker}}$*, so that*

$$\Omega_\sigma^{\mathrm{presel,proker}} \circ \mathfrak{y}_\sigma^{\mathrm{presel,proker}} = \mathrm{id}_{\mathcal{P}_\sigma^{\mathrm{presel,preker}}}.$$

*In particular, the induced pre-kernel functional* $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ *is a well-defined smooth function on* $\mathcal{N}_\sigma^{\mathrm{presel,preker}}$ *whose values on* $\mathcal{P}_\sigma^{\mathrm{presel,preker}}$ *are the effective minimum values of the deeper proto-kernel functional on the inner proto-kernel fibers.*

*Proof.* Definition 8(1) makes every inner proto-kernel fiber $\mathcal{X}_\sigma^{\mathrm{presel,proker}}(q)$ nonempty, compact, and convex. Definition 8(2) states that $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ has a unique minimizer on each such fiber and that the minimizer depends smoothly on $q$. Therefore $\mathfrak{y}_\sigma^{\mathrm{presel,proker}}$ is a well-defined smooth map on

$\mathcal{P}_\sigma^{\mathrm{presel,preker}}$. Each chosen minimizer lies in the fiber over the point at which it is selected, so

$$\Omega_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(q)\big) = q$$

for every $q \in \mathcal{P}_\sigma^{\mathrm{presel,preker}}$. The remaining smoothness statement for $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ on $\mathcal{N}_\sigma^{\mathrm{presel,preker}}$ is part of Definition 8(2), and the displayed formula identifies its values on $\mathcal{P}_\sigma^{\mathrm{presel,preker}}$ with the effective minimum values. □

**Proposition 2** (Current-kernel-scale proto-kernel minimization factors through the induced pre-kernel minimization)**.** *For every benchmark-faithful proto-kernel package and every $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$,*

$$\mathfrak{w}_\sigma^{\mathrm{presel,proker}}(w) = \mathfrak{v}_\sigma^{\mathrm{presel,proker}}\big(\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w)\big),$$

*and hence*

$$\Omega_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{w}_\sigma^{\mathrm{presel,proker}}(w)\big) = \widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w).$$

*Proof.* Fix $w \in \mathcal{K}_\sigma^{\mathrm{presel,ker}}$. Every $z \in \mathcal{L}_\sigma^{\mathrm{presel,proker}}(w)$ satisfies $q \equiv \Omega_\sigma^{\mathrm{presel,proker}}(z) \in \mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)$. By Proposition 1,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(q)\big) = \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q).$$

By Definition 8(2), the induced pre-kernel functional is strictly convex on $\mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)$ and therefore has a unique minimizer $\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w)$. Consequently, for every $z \in \mathcal{L}_\sigma^{\mathrm{presel,proker}}(w)$,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}\big(\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w)\big) = \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w))\big).$$

So $\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}(w))$ is a minimizer of $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ on the current-kernel-scale composed proto-kernel fiber $\mathcal{L}_\sigma^{\mathrm{presel,proker}}(w)$. Definition 8(3) says that minimizer is unique, hence the first displayed identity follows. The second identity is then Proposition 1. □

**Proposition 3** (The induced outer pre-kernel dynamics form a benchmark-faithful package)**.** *For every benchmark-faithful proto-kernel package, the tuple*

$$\Big(\mathcal{N}_\sigma^{\mathrm{presel,preker}}, \mathcal{P}_\sigma^{\mathrm{presel,preker}}, \Upsilon_\sigma^{\mathrm{presel,preker}}, \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}, \widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}\Big)$$

*is a benchmark-faithful kernel-dynamics-generating substrate pre-selector pre-kernel dynamics package on the recorded geometry in the sense of Definition 6.*

*Proof.* Definition 8(2) states that $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ is nonnegative and smooth, and that its restriction to each current pre-kernel fiber $\mathcal{U}_\sigma^{\mathrm{presel,preker}}(w)$ is strictly convex with a unique minimizer depending smoothly on $w$. Definition 8(3) defines the induced current kernel functional $\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}$ and states that its restriction to each current kernel fiber $\mathcal{J}_\sigma^{\mathrm{presel,ker}}(u)$ is strictly convex with a unique minimizer depending smoothly on $u$. Definition 8(4) defines the induced outer selector functional and states that its restriction to each current selector fiber $\mathcal{G}_\sigma^{\mathrm{presel,sel}}(y)$ is strictly convex with a unique minimizer depending smoothly on $y$. Definitions 8(5) and 8(6) give the current composed-kernel minimizer, the induced exact selector, kernel, and pre-kernel shells, the full pre-kernel minimizer, and the required shell diffeomorphisms. Definition 8(7) gives benchmark faithfulness. These are precisely the requirements of Definition 6 on the recorded geometry. □

**Proposition 4** (Selector-scale proto-kernel minimization factors through the induced pre-kernel composed minimization)**.** *For every benchmark-faithful proto-kernel package and every* $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$*,*

$$\mathfrak{x}_\sigma^{\mathrm{presel,proker}}(u) = \mathfrak{v}_\sigma^{\mathrm{presel,proker}}\big(\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u)\big),$$

*where*

$$\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u) \equiv \arg \min_{q \in \mathcal{V}_\sigma^{\mathrm{presel,preker}}(u)} \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q),$$

*and hence*

$$\Omega_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{x}_\sigma^{\mathrm{presel,proker}}(u)\big) = \widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u).$$

*Proof.* Fix $u \in \mathcal{B}_\sigma^{\mathrm{presel,sel}}$. Every $z \in \mathcal{A}_\sigma^{\mathrm{presel,proker}}(u)$ satisfies

$$q \equiv \Omega_\sigma^{\mathrm{presel,proker}}(z) \in \mathcal{V}_\sigma^{\mathrm{presel,preker}}(u).$$

By Proposition 1,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(q)\big) = \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q).$$

Definition 8(4) records that $\widetilde{\mathscr{Q}}_\sigma^{\mathrm{presel,sel}}$ is induced from $\widetilde{\mathscr{R}}_\sigma^{\mathrm{presel,ker}}$, while Proposition 3 says the induced outer data form a benchmark-faithful current pre-kernel package. Therefore the restriction of $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ to $\mathcal{V}_\sigma^{\mathrm{presel,preker}}(u)$ attains a unique minimizer. By definition, that minimizer is $\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u)$. Consequently, for every $z \in \mathcal{A}_\sigma^{\mathrm{presel,proker}}(u)$,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}\big(\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u)\big) = \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}\big(\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u)\big)\big).$$

So $\mathfrak{v}_\sigma^{\mathrm{presel,proker}}\big(\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}(u)\big)$ is a minimizer of $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ on the selector-scale composed proto-kernel fiber $\mathcal{A}_\sigma^{\mathrm{presel,proker}}(u)$. Definition 8(4) says that minimizer is unique, hence the first displayed identity follows. The second identity is then Proposition 1. □

**Proposition 5** (Benchmark-shell proto-kernel minimization factors through the induced pre-kernel full minimization)**.** *For every benchmark-faithful proto-kernel package and every* $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$*,*

$$\mathfrak{y}_\sigma^{\mathrm{presel,proker}}(y) = \mathfrak{v}_\sigma^{\mathrm{presel,proker}}\big(\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y)\big),$$

*where*

$$\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y) \equiv \arg \min_{q \in \mathcal{W}_\sigma^{\mathrm{presel,preker}}(y)} \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q),$$

*and hence*

$$\Omega_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{y}_\sigma^{\mathrm{presel,proker}}(y)\big) = \widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y).$$

*Proof.* Fix $y \in \mathcal{B}_\sigma^{\mathrm{yoc,sel}}$. Every $z \in \mathcal{C}_\sigma^{\mathrm{presel,proker}}(y)$ satisfies

$$q \equiv \Omega_\sigma^{\mathrm{presel,proker}}(z) \in \mathcal{W}_\sigma^{\mathrm{presel,preker}}(y).$$

By Proposition 1,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(q)\big) = \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}(q).$$

Definition 8(5) and Proposition 3 together show that the induced pre-kernel functional attains a unique minimum on $\mathcal{W}_\sigma^{\mathrm{presel,preker}}(y)$. By definition, that minimizer is $\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y)$. Therefore, for every $z \in \mathcal{C}_\sigma^{\mathrm{presel,proker}}(y)$,

$$\mathscr{U}_\sigma^{\mathrm{presel,proker}}(z) \geq \widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}\big(\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y)\big) = \mathscr{U}_\sigma^{\mathrm{presel,proker}}\big(\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y))\big).$$

So $\mathfrak{v}_\sigma^{\mathrm{presel,proker}}(\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}(y))$ is a minimizer of $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ on the full proto-kernel fiber $\mathcal{C}_\sigma^{\mathrm{presel,proker}}(y)$. Definition 8(6) says that minimizer is unique, hence the first displayed identity follows. The second identity is then Proposition 1. □

**Proposition 6** (The exact pre-kernel shell is the projected exact proto-kernel shell)**.** *For every benchmark-faithful proto-kernel package,*

$$\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}} = \Omega_\sigma^{\mathrm{presel,proker}}\big(\mathcal{E}_\sigma^{\mathrm{presel,proker}}\big),$$

*the restriction*

$$\Omega_\sigma^{\mathrm{presel,proker}}\big|_{\mathcal{E}_\sigma^{\mathrm{presel,proker}}} : \mathcal{E}_\sigma^{\mathrm{presel,proker}} \xrightarrow{\cong} \widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}$$

*is a diffeomorphism, and*

$$\Upsilon_\sigma^{\mathrm{presel,preker}}\big|_{\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}} : \widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}} \xrightarrow{\cong} \widetilde{\mathcal{Y}}_\sigma^{\mathrm{presel,ker}}$$

*is a diffeomorphism.*

*Proof.* The two diffeomorphisms are part of Definition 8(5) and Definition 8(6). In particular, the first one immediately implies

$$\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}} = \Omega_\sigma^{\mathrm{presel,proker}}\big(\mathcal{E}_\sigma^{\mathrm{presel,proker}}\big).$$

Nothing further is required. □

**Proposition 7** (No residual pre-kernel-level freedom remains once the proto-kernel package is fixed)**.** *For every benchmark-faithful proto-kernel package, the current pre-kernel functional $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$, the current pre-kernel minimizer lift $\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}$, the current selector-scale pre-kernel composed minimizer $\widetilde{\mathfrak{t}}_\sigma^{\mathrm{presel,preker}}$, the current benchmark-shell-scale full pre-kernel minimizer $\widetilde{\mathfrak{u}}_\sigma^{\mathrm{presel,preker}}$, and the current exact pre-kernel shell $\widetilde{\mathcal{O}}_\sigma^{\mathrm{presel,preker}}$ are uniquely determined by the proto-kernel data. In particular, the current pre-kernel package carries no residual benchmark-facing functional, shell, current-kernel-scale minimizer, selector-scale minimizer, or benchmark-shell-scale minimizer freedom once the deeper proto-kernel package is fixed.*

*Proof.* Proposition 1 determines $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ uniquely as the effective minimum value of $\mathscr{U}_\sigma^{\mathrm{presel,proker}}$ on the inner proto-kernel fibers. Definition 8(2) then uniquely determines $\widetilde{\mathfrak{s}}_\sigma^{\mathrm{presel,preker}}$ as the unique minimizer of $\widetilde{\mathscr{S}}_\sigma^{\mathrm{presel,preker}}$ on each current pre-kernel fiber. Proposition 4 identifies the current selector-scale pre-kernel composed minimizer as the outer projection of the unique selector-scale proto-kernel minimizer, while Proposition 5 identifies the benchmark-shell-scale full pre-kernel minimizer as the outer projection of the unique full proto-kernel minimizer. Definition 8(5) and Proposition 6 determine the current exact pre-kernel shell uniquely as the projected exact proto-kernel shell over $\mathcal{Z}_\sigma^{\mathrm{yoc,sel}}$. Therefore no benchmark-facing constituent at current pre-kernel scope remains freely choosable once the proto-kernel package is fixed. □

**Proposition 8** (All current downstream uses factor through the induced pre-kernel dynamics)**.** *For every benchmark-faithful proto-kernel package, all current benchmark-facing downstream uses of the current pre-kernel package on the active line factor through the induced pre-kernel package of Proposition 3. In particular, the current selector-dynamics-generating substrate pre-selector kernel dynamics package* $\mathfrak{F}^{\mathrm{presel,ker}}$*, the current pre-selector-minimizer-section-generating substrate selector dynamics package* $\mathfrak{E}^{\mathrm{presel,sel}}$*, the current observer-relevant pre-selector minimizer-section core* $\mathfrak{D}^{\mathrm{presel,sec}}$*, the current selector package* $\mathfrak{C}^{\mathrm{yoc,sel}}$*, the current observer-relevant yocto-section minimizer-section core* $\mathfrak{C}^{\mathrm{yoc,sec}}$*, the current observer-relevant yocto-section minimizer-image core* $\mathfrak{C}^{\mathrm{yoc,img}}$*, the current observer-relevant zepto-section minimizer-section core* $\mathfrak{B}^{\mathrm{zep,sec}}$*, the current observer-relevant zepto-section minimizer-image core* $\mathfrak{B}^{\mathrm{zep,img}}$*, the current atto-section package* $\mathfrak{A}^{\mathrm{attosec}}$*, the current femto-section package* $\mathfrak{Z}^{\mathrm{femtosec}}$*, the current pico-section package* $\mathfrak{Y}^{\mathrm{picosec}}$*, the current observer-relevant pico-section minimizer-section core* $\mathfrak{Y}^{\mathrm{pic,sec}}$*, the current nano-section package* $\mathfrak{X}^{\mathrm{nanosec}}$*, the current micro-section package* $\mathfrak{W}^{\mathrm{microsec}}$*, the current sub-section package* $\mathfrak{V}^{\mathrm{subsec}}$*, and the previously recorded shell-transport consequences follow unchanged.*

*Proof.* By Proposition 3, the induced outer data form a benchmark-faithful current pre-kernel package of the exact type required by Definition 6. Definition 7(2) then records that, once such a package is available on the recorded geometry, the current kernel package and all downstream benchmark-facing data and consequences follow unchanged. Therefore all current downstream uses already factor through the induced pre-kernel package provided by the deeper proto-kernel dynamics. □

**Proposition 9** (The benchmark package remains fixed)**.** *Every benchmark-faithful pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics package preserves the same benchmark one-metric package* (1)*. In particular, the operative metric remains exactly* $g^\sharp$*, the ordinary matter route is unchanged, there is no second operative metric, and the low-energy matter law is not enlarged.*

*Proof.* This is part of benchmark faithfulness in Definition 8(7). The present note only derives the current pre-kernel package from deeper substrate proto-kernel data. It does not alter the benchmark exact-closure package. □

## 5. Main Theorem

**Theorem 1** (Kernel-dynamics-generating substrate pre-selector pre-kernel dynamics from pre-kernel–dynamics-generating substrate pre-selector proto-kernel dynamics)**.** *Fix the primitive continuation data of Definition 1, the recorded selector benchmark base of Definition 2, the recorded current selector-body geometry of Definition 3, the recorded current kernel-body geometry of Definition 4, the recorded current pre-kernel-body geometry of Definition 5, the recorded downstream chain of Definition 7, and a benchmark-faithful pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics package in the sense of Definition 8. Then, for each sector* $\sigma \in \{\mathrm{car}, \mathrm{cur}, \mathrm{hom}\}$*, the current kernel-dynamics-generating substrate pre-selector pre-kernel dynamics are no longer an unexplained primitive stopping point on the active line: they are recovered canonically from the deeper proto-kernel dynamics. More precisely:*

1. *the current pre-kernel functional is induced as the effective minimum value of the deeper proto-kernel functional on the inner proto-kernel fibers by Proposition 1;*

2. *the current-kernel-scale pre-kernel minimizer lift is the outer projection of the unique current-kernel-scale proto-kernel minimizer by Proposition 2;*

3. *the induced outer data form a benchmark-faithful current pre-kernel package on the recorded geometry by Proposition 3;*

4. *the current selector-scale pre-kernel composed minimizer is the outer projection of the unique selector-scale proto-kernel minimizer by Proposition 4;*

5. *the current benchmark-shell-scale full pre-kernel minimizer is the outer projection of the unique full proto-kernel minimizer by Proposition 5;*

6. *the current exact pre-kernel shell is exactly the projected exact proto-kernel shell, and it still projects diffeomorphically to the induced exact kernel shell, by Proposition 6;*

7. *the current pre-kernel package carries no residual benchmark-facing functional, shell, current-kernel-scale minimizer, selector-scale minimizer, or benchmark-shell-scale minimizer freedom once the deeper proto-kernel package is fixed by Proposition 7;*

8. *all current downstream benchmark-facing uses of the pre-kernel package already factor through the induced outer package by Proposition 8;*

9. *the benchmark boundary remains fixed by Proposition 9;*

10. *consequently the remaining honest burden below the current pre-selector pre-kernel frontier is no longer the kernel-dynamics-generating substrate pre-selector pre-kernel dynamics* $\mathfrak{G}^{\mathrm{presel,preker}}$ *as such. What remains is to derive or justify the deeper pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics* $\mathfrak{H}^{\mathrm{presel,proker}}$ *from still more primitive substrate structure, or else prove a still sharper theorem-level collapse strictly inside that deeper package.*

*Proof.* Item (1) is Proposition 1. Item (2) is Proposition 2. Item (3) is Proposition 3. Item (4) is Proposition 4. Item (5) is Proposition 5. Item (6) is Proposition 6. Item (7) is Proposition 7. Item (8) is Proposition 8. Item (9) is Proposition 9. Item (10) is the routing consequence of the previous items together with the active safeguard against counting another same-output deeper-origin relay as new front-line progress when the live benchmark-relevant pre-kernel package is unchanged. □

**Corollary 1** (What must be done next)**.** *After Theorem 1, the next honest benchmark-facing manuscript must either:*

1. *derive or justify the pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics* $\mathfrak{H}^{\mathrm{presel,proker}}$ *from still more primitive substrate structure in a way that changes benchmark-facing status;*

2. *identify a genuinely smaller theorem-level observer-relevant datum strictly inside* $\mathfrak{H}^{\mathrm{presel,proker}}$ *and prove a sharper collapse to it;*

3. *do either of the previous two while preserving the same benchmark one-metric package and the same claim boundary.*

*Another same-output deeper-origin relay that merely preserves the current proto-kernel dynamics* $\mathfrak{H}^{\mathrm{presel,proker}}$ *and therefore merely reproduces the same current pre-kernel package* $\mathfrak{G}^{\mathrm{presel,preker}}$*, the*

*same current kernel package* $\mathfrak{F}^{\mathrm{presel,ker}}$*, the same current selector-dynamics package* $\mathfrak{E}^{\mathrm{presel,sel}}$*, the same current observer-relevant pre-selector minimizer-section core* $\mathfrak{D}^{\mathrm{presel,sec}}$*, the same current selector package* $\mathfrak{C}^{\mathrm{yoc,sel}}$*, the same current observer-relevant yocto-section minimizer-section core* $\mathfrak{C}^{\mathrm{yoc,sec}}$*, the same current observer-relevant yocto-section minimizer-image core* $\mathfrak{C}^{\mathrm{yoc,img}}$*, and the same downstream observer-relevant zepto-section, atto-section, femto-section, pico-section, nano-section, micro-section, and sub-section consequences, another re-expansion back to the current pre-kernel package, the current kernel package, the current selector-dynamics package, the larger pre-selector package, or older yocto-section presentations as if they were still independently live, another reopening of larger screened layers, or any move that introduces a second operative metric, enlarges the ordinary matter law, or uses stronger promotion language does not satisfy the present burden.*

*Proof.* Theorem 1 shows that the current pre-kernel package is no longer the deepest unexplained datum on the active line. Therefore a subsequent manuscript must improve on the deeper proto-kernel dynamics rather than merely restate the already induced pre-kernel package. □

## 6. Conclusion

The positive result of this note is that the current kernel-dynamics-generating substrate pre-selector pre-kernel dynamics are no longer primitive on the active derivational line. Once one works with pre-kernel-dynamics-generating substrate pre-selector proto-kernel dynamics over the already fixed current pre-kernel-body geometry, the current pre-kernel functional is forced as the effective minimum value of the deeper proto-kernel law, the current-kernel-scale pre-kernel minimizer lift is forced as the outer projection of the unique current-kernel-scale proto-kernel minimizer, the current selector-scale pre-kernel composed minimizer is forced as the outer projection of the unique selector-scale proto-kernel minimizer, the current benchmark-shell-scale full pre-kernel minimizer is forced as the outer projection of the unique full proto-kernel minimizer, and the current exact pre-kernel shell is forced as the projected exact proto-kernel shell over the recorded selector benchmark shell. All current downstream benchmark-facing uses then factor through that induced pre-kernel package together with the already fixed selector benchmark base, selector-body geometry, and kernel-body geometry.

This preserves the same benchmark one-metric package and the same claim boundary. The result does not provide a completed first-principles derivation of GR from substrate microphysics, and it does not license stronger promotion language. Its narrower benchmark-facing gain is that the remaining honest burden now sits one level lower again: not on the current pre-selector pre-kernel dynamics themselves, but on the deeper pre-selector proto-kernel dynamics that canonically generate them, or on a still smaller collapse strictly inside that deeper package.