

Feedback — Problem Set-1

[Help Center](#)

You submitted this quiz on **Mon 19 Jan 2015 6:49 PM IST**. You got a score of **4.00** out of **5.00**.

Question 1

3-way-Merge Sort : Suppose that instead of dividing in half at each step of Merge Sort, you divide into thirds, sort each third, and finally combine all of them using a three-way merge subroutine.

What is the overall asymptotic running time of this algorithm? (Hint: Note that the merge step can still be implemented in $O(n)$ time.)

Your Answer	Score	Explanation
<input type="radio"/> n		
<input type="radio"/> $n(\log(n))^2$		
<input checked="" type="radio"/> $n \log(n)$	✓ 1.00	That's correct! There is still a logarithmic number of levels, and the overall amount of work at each level is still linear.
<input type="radio"/> $n^2 \log(n)$		
Total	1.00 / 1.00	

Question 2

You are given functions f and g such that $f(n) = O(g(n))$. Is

$f(n) * \log_2(f(n)^c) = O(g(n) * \log_2(g(n)))$? (Here c is some positive constant.) You should assume that f and g are nondecreasing and always bigger than 1.

Your Answer	Score	Explanation
<input checked="" type="radio"/> True	✓ 1.00	That's correct! Roughly, because the constant c in the exponent is inside a logarithm, it becomes part of the leading constant and gets suppressed by the big-Oh notation.
<input type="radio"/> False		
<input type="radio"/> Sometimes yes, sometimes no, depending on the functions f and g		
<input type="radio"/> Sometimes yes, sometimes no, depending on the constant c		
Total	1.00 / 1.00	

Question 3

Assume again two (positive) nondecreasing functions f and g such that $f(n) = O(g(n))$. Is $2^{f(n)} = O(2^{g(n)})$? (Multiple answers may be correct, you should check all of those that apply.)

Your Answer	Score	Explanation
<input checked="" type="checkbox"/> Sometimes	✓ 0.25	
<input type="checkbox"/> Always	✓ 0.25	What if $f(n) = 2n$ and $g(n) = n$?
<input checked="" type="checkbox"/> Yes if $f(n) \leq g(n)$ for all sufficiently large n	✓ 0.25	
<input type="checkbox"/> Never	✓ 0.25	For example, what if $f(n)=g(n)$?
Total	1.00 / 1.00	

Question 4

k-way-Merge Sort. Suppose you are given k sorted arrays, each with n elements, and you want to combine them into a single array of kn elements. Consider the following approach. Using the merge subroutine taught in lecture, you merge the first 2 arrays, then merge the 3rd given array with this merged version of the first two arrays, then merge the 4th given array with the merged version of the first three arrays, and so on until you merge in the final (k^{th}) input array. What is the running time taken by this successive merging algorithm, as a function of k and n ? (Optional: can you think of a faster way to do the k-way merge procedure ?)

Your Answer	Score	Explanation
<input type="radio"/> $\theta(n^2k)$		
<input checked="" type="radio"/> $\theta(nk^2)$	1.00	That's correct! For the upper bound, the merged list size is always $O(kn)$, merging is linear in the size of the larger array, and there are k iterations. For the lower bound, each of the last $k/2$ merges takes $\Omega(kn)$ time.
<input type="radio"/> $\theta(nk)$		
<input type="radio"/> $\theta(n \log(k))$		
Total	1.00 / 1.00	

Question 5

Arrange the following functions in increasing order of growth rate (with $g(n)$ following $f(n)$ in your list if and only if $f(n) = O(g(n))$).

a) \sqrt{n}

b) 10^n

c) $n^{1.5}$


d) $2^{\sqrt{\log(n)}}$

e) $n^{5/3}$

Write your 5-letter answer, i.e., the sequence in lower case letters in the space provided. For example, if you feel that the answer is a->b->c->d->e (from smallest to largest), then type abcde in the space provided without any spaces before / after / in between the string. You can assume that all logarithms are base 2 (though it actually doesn't matter). WARNING: this question has multiple versions, you might see different ones on different attempts!

You entered:

becad

Your Answer	Score	Explanation
becad	 0.00	
Total	0.00 / 1.00	

Question Explanation

One approach is to graph these functions for large values of n. Once in a while this can be misleading, however. Another useful trick is to take logarithms and see what happens (though again be careful, as in Question 3).

