

Colombia

Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening

Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is among the top 10 cancers in men.
- Incidence rate: Around 11 per 100,000 men per year.
- Total new cases (2022): Approximately 8,938 men.
- Daily diagnoses (2022): About 24 men per day.
- Deaths (2022): Roughly 2,591 men.
- 5-year survival rate: Estimated below 40–45%.
- Most affected age group: Typically men aged 60–75.
- Screening participation: No formal screening; diagnosis usually occurs at symptomatic stages.

Colombia



Infrastructure



Strengths

- Urban cancer centers in Bogotá (Instituto Nacional de Cancerología) and Medellín (Clínica Antioquia) provide advanced endoscopy, surgery, imaging, and pathology, including molecular diagnostics.
- National cancer registry and regional oncology networks support quality monitoring and patient referrals.

Weakness

- Rural and underserved regions (e.g. Chocó, Arauca) lack endoscopic units, pathology services, and oncology specialists.
- Referral systems from primary to tertiary care are inefficient, leading to diagnostic and treatment delays.

Opportunity

- Expansion of regional cancer infrastructure through public funding or partnership with academic institutions.
- Implementation of telepathology and mobile endoscopy units to serve geographically remote populations.

Threats


- Urban facility overcrowding, particularly in Bogotá and Medellín, causes long wait times and diagnostic bottlenecks.
- Healthcare provider migration from public to private sector and abroad drains talent from public institutions.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Colombia

Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Public insurance systems (SISBEN/SGSSS) cover standard gastric cancer treatments including surgery and chemotherapy.
- Participation in international GC clinical trials provides patients with access to biomarker-driven therapies (e.g., HER2-targeted agents, immunotherapy in MSI-H or PD-L1-positive cases).

Weakness

- Funding and clinical research capacity for gastric cancer remain limited; few nationally led biomarker-specific trials.
- Low public awareness of gastric cancer risk factors (high-salt diet, H. pylori, smoking) results in late presentation.

Opportunity

- Collaborate with academic and NGO partners to increase funding into nationwide research and awareness campaigns.
- Leverage media and community health worker networks for culturally adapted education about prevention and early symptoms.

Threats

- Budget competition with more high-profile cancers (breast, prostate, cervical) can result in underinvestment in GC initiatives.
- Socioeconomic barriers may limit uptake of awareness campaigns among marginalized groups.



5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.



4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.



3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.



2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.




1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Colombia

Survival Rates, Early Detection and Palliative Care



Strengths

- In regional cancer centers offering early diagnosis and multidisciplinary care, 5-year survival for localized GC exceeds 40%.
- Palliative care integration is increasing in major hospitals, improving symptom control and quality of life for advanced GC patients.

Weakness


- Nationwide, most GC patients present at advanced stage III–IV; overall 5-year survival remains below 30%.
- Palliative care is poorly available in non-urban areas, with limited hospices or home-care programs.

Opportunity

- Deploy early detection programs using symptom screening and risk stratification in primary care settings.
- Train rural and district-level providers in palliative best practices and pain management

Threats

- Cultural stigma around cancer may discourage early symptom presentation.
- Fragmented follow-up systems delay post-surgical care or palliative referrals.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.



3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Colombia



Utilization of Biomarkers



Strengths

- HER2 testing is available and guiding trastuzumab use in public and private tertiary centers for advanced GC.
- MSI-H/dMMR and PD-L1 (CPS ≥ 1) profiling is performed at academic institutions to select for immunotherapy.

Opportunity




- CLDN18.2 and FGFR2b testing are rarely available outside trial settings—even where clinical utility is emerging.
- Variability in lab accreditation and biomarker reporting contributes to inconsistent test quality across regions.


Weakness

- CLDN18.2 and FGFR2b testing are rarely available outside trial settings—even where clinical utility is emerging.
- Variability in lab accreditation and biomarker reporting contributes to inconsistent test quality across regions.

Threats



- High costs and low insurance coverage for less common biomarkers hinder equitable access.
- Delay in reimbursement policy for biomarker assays leads to underutilization by clinicians.

-  Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
-  Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
-  Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.



Colombia

Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	✗	◯	✗	✗
Feasibility of Integration	✗	✗	◯	✗	✗
Adoption of International Guidelines	✗	✗	◯	✗	✗
Engagement with Updates	✗	✗	✗	◯	✗
ESMO Guidelines Implementation	✗	✗	◯	✗	✗

Colombia

Reimbursement



Strengths

- Public schemes cover essential treatments, including surgery, chemotherapy, and trastuzumab for HER2+ GC.
- Clinical trial participation provides alternative routes for access to advanced therapies not yet reimbursed officially.

Weakness

- Newer agents (e.g., immune checkpoint inhibitors in PD-L1 positive patients, CLDN18.2 or FGFR2b therapies) are not well-reimbursed.
- Diagnostic testing for biomarkers such as MSI, PD-L1, CLDN18.2 is rarely covered and often requires out-of-pocket payment.

Opportunity

- Advocate for inclusion of critical biomarker tests and immunotherapy agents in public reimbursement frameworks.
- Pilot outcome-based payment or managed access schemes to accelerate coverage for effective therapies.

Threats

- Budget constraints and rising pharmaceutical costs limit the expansion of public coverage.
- Fragmentation among health insurers and regional systems creates inequitable access pathways.























































































A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.



A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.




No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States		
United Kingdom		
Canada		
Australia		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Japan		
South Korea		
China		
India		
Singapore		
Thailand		
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Colombia



Gastric Cancer Screening



Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities