



Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening

Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is not among the top 5, but remains significant in men.
- Incidence rate: Approximately 14 per 100,000 men per year.
- Total new cases (2022): Around 5,000–5,500 men.
- Daily diagnoses: Approximately 14–15 men per day.
- Deaths (2022): Around 3,500 men.
- 5-year survival rate: Estimated 30–40%.
- Most affected age group: Primarily men aged 70 and above.
- Screening participation: No national screening; detection is mostly symptom-driven

Poland



Infrastructure



Strengths

- Poland has a network of public oncology centers under the National Oncology Network (Narodowa Sieć Onkologiczna) providing multidisciplinary care.
- Modern cancer centers like the Maria Skłodowska-Curie National Research Institute of Oncology in Warsaw offer advanced diagnostics and treatment.

Weakness


- Regional inequalities persist, with rural and eastern provinces facing shortages of oncology specialists and endoscopy units.
- Waiting times for diagnostic procedures such as endoscopy can delay timely diagnosis and treatment.

Opportunity

- Ongoing investments through EU structural funds and national health programs support modernization of cancer care infrastructure.
- Potential for mobile diagnostic clinics and expanded regional hubs to decentralize access.

Threats


- Workforce shortages in oncology and pathology can hamper infrastructure utilization.
- Aging hospital facilities in smaller towns limit the expansion of high-tech cancer services.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Poland

Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Gastric cancer treatment is publicly funded under the National Health Fund (NFZ), ensuring broad financial coverage.
- Active participation in clinical trials, especially for HER2-positive and immunotherapy gastric cancer subtypes.

Weakness

- Limited national public awareness campaigns specifically targeting gastric cancer symptoms and risk factors.
- Research funding often prioritizes cancers with higher media coverage (e.g., breast, prostate), reducing resources for gastric cancer.


















































































































Opportunity

- Expansion of public education under the Polish National Oncology Strategy could include gastric cancer prevention and risk factors like H. pylori.
- Cross-border collaboration with European cancer consortia can boost funding and innovation.

Threats


- Health policy instability or shifting government priorities may affect sustained support for awareness and research.
- Low public engagement in screening campaigns could limit campaign effectiveness.

5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Poland

Survival Rates, Early Detection and Palliative Care



Strengths

- National cancer registry and survival data allow ongoing monitoring and targeted interventions.
- Palliative care is integrated into cancer services, with legal provisions supporting end-of-life care.

Weakness

- Many gastric cancers are diagnosed at advanced stages due to late symptom recognition.
- Referral and diagnostic delays persist, especially outside major urban centers.

Opportunity

- Early detection programs through general practitioners and screening in high-risk groups could significantly improve stage at diagnosis.
- Expanded community palliative care models can improve access in underserved regions.

Threats

- Social stigma or fatalistic attitudes toward cancer still delay early medical help-seeking.
- Insufficient trained personnel in palliative care in certain voivodeships.


5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.
4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.
3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.
2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.
1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Poland



Utilization of Biomarkers



Strengths

- HER2 testing is routinely performed in metastatic gastric cancer patients and trastuzumab is reimbursed.
- Some reference laboratories perform PD-L1 and MSI testing for immunotherapy eligibility.

Weakness




- Limited routine testing for emerging biomarkers like CLDN18.2 or FGFR2b in non-trial settings.
- Regional disparities in molecular diagnostics infrastructure and turnaround times.

Opportunity


- EU-funded precision oncology projects can support expansion of biomarker testing capabilities.
- Centralization of testing to specialized hubs could ensure standardization and quality control.

Threats

- Reimbursement uncertainty for advanced biomarker panels may deter broader adoption.
- Delays in incorporating emerging biomarker-based therapies into national protocols.

-  Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
-  Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
-  Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.

Poland



Poland



Clinical Guidelines

Strengths

- National guidelines for gastric cancer align closely with ESMO and NCCN standards.
- Multidisciplinary teams (MDTs) are mandated in cancer centers for treatment planning.

Weakness



- Guideline adherence can vary, particularly in less specialized regional hospitals.
- Slow dissemination of updates to frontline practitioners, especially in smaller centers.

Opportunity

- Digital dissemination and e-learning platforms can improve clinician access to updated protocols.
- Greater integration of European guideline frameworks and real-world data to support policy adjustments.

Threats

- Resistance to practice change or lack of continuing education may reduce uptake of new guidelines.
- Delays in national adoption of emerging evidence-based treatments due to bureaucratic processes.




	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	○	✗	✗	✗
Feasibility of Integration	✗	○	✗	✗	✗
Adoption of International Guidelines	✗	○	✗	✗	✗
Engagement with Updates	✗	✗	✗	○	✗
ESMO Guidelines Implementation	✗	○	✗	✗	✗

Poland



Reimbursement



Poland



Colorectal Cancer Screening

Strengths

- Opportunistic gastroscopy is widely available for symptomatic or at-risk patients.
- Pilot programs and research initiatives have evaluated *H. pylori* prevalence and eradication strategies.

Weakness

- No population-wide gastric cancer screening program despite relatively high incidence in some regions.
- Lack of integration between primary care and gastroenterology hampers proactive case finding.

Opportunity

- National efforts on cancer prevention (e.g., tobacco/alcohol control, *H. pylori* testing) can be aligned with early gastric cancer detection.
- Research into risk stratification models may support targeted screening.

Threats

- Public reluctance toward invasive procedures (endoscopy) limits screening participation.
- Competing priorities in national screening programs (e.g., colorectal and breast cancer) may reduce attention to gastric cancer.

Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities