



Colorectal Cancer Factsheet: Insights & Key Developments

Key Insights on Colorectal Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Colorectal Cancer Screening

Colorectal cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping colorectal cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Prostate cancer ranks among the top cancers in Rwandan men, though overall cancer burden is low.
- Incidence rate: Estimated at low single digits to low teens per 100,000 men per year (registry-based estimate).
- Total new cases (2022): Approximately approximately 500–800 men based on registry coverage.
- Daily diagnoses (2022): Fewer than 3 men per day.
- Deaths (2022): Estimated around 300–350 men.
- 5-year survival rate: Likely below 50%, due to late-stage diagnosis and limited treatment infrastructure.
- Most affected age group: Mostly men aged 65 and older.
- Screening participation: Virtually no PSA screening or awareness programs; diagnosis often occurs at advanced stages, even in referral centers.

Rwanda



Infrastructure



Strengths

- Rwanda has made notable progress in strengthening its public healthcare infrastructure, especially after its national strategic health investments.
- Referral hospitals like King Faisal Hospital and Rwanda Military Hospital provide oncology and urology services.

Weakness

- Limited access to cancer diagnostic and treatment facilities outside Kigali; rural areas lack imaging and pathology services.
- Shortage of specialists, including oncologists, pathologists, and trained urologists

Opportunity

- Strengthen decentralized cancer units and mobile diagnostic units for rural populations.
- Invest in public-private partnerships to build regional diagnostic labs.

Threats

- Heavy reliance on external aid and donor support for advanced diagnostics and infrastructure.
- Delays in equipment procurement due to regulatory or import challenges




Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Rwanda



Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Rwanda’s universal healthcare coverage under the Community-Based Health Insurance (Mutuelle de Santé) helps reduce cost barriers.
- Some awareness campaigns on non-communicable diseases have included male reproductive health topics.

Weakness


- Very limited availability of radiotherapy, chemotherapy, and surgical oncology for prostate cancer.
- Prostate cancer is underfunded in national research priorities and receives less attention than infectious diseases.

Opportunity

- Include prostate cancer more explicitly in national NCD plans and awareness drives.
- Build local research capacity through collaborations with African and global cancer networks.

Threats

- Competing health priorities like HIV, malaria, and maternal care may crowd out cancer care investments.
- Sociocultural taboos around male health can reduce campaign effectiveness.




5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●
Malaysia	●	●	●

Rwanda

Survival Rates, Early Detection and Palliative Care



Strengths

- Rwanda has improved palliative care access, integrated into primary health facilities in some districts.
- Community health workers (CHWs) are trained to identify symptoms and refer patients early.

Weakness

- No nationwide early detection program, leading to late-stage presentations of prostate cancer.
- Survival data are not systematically collected; cancer registries are still under development.

Opportunity

- Train CHWs and primary care providers to include prostate symptoms in routine male screenings.
- Invest in data collection and cancer registry development to monitor outcomes.

Threats

- Late diagnosis and limited treatment capacity increase morbidity and mortality.
- Stigma around prostate issues, especially among older men, reduces early reporting.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.






























































































































3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Rwanda



Utilization of Biomarkers



Strengths

- PSA testing is available at tertiary hospitals in Kigali and some private clinics.
- National efforts are underway to strengthen laboratory services for better diagnostics.

Weakness

- Advanced biomarkers (BRCA1/2, PTEN, TMPRSS2-ERG) are largely unavailable and rarely used in clinical decision-making.
- Most clinicians lack training on interpreting genomic or molecular biomarker results.

Opportunity


- Introduce low-cost PSA screening in primary health centers targeting men above 50.
- Collaborate with regional genomics labs (e.g., in Kenya or South Africa) to offer reference-level testing.

Threats

- Financial and logistical constraints hinder scaling up biomarker testing.
- Biomarker-based decisions cannot be implemented effectively without linked treatment options.



Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.




Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.



Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.

Rwanda




Rwanda



Clinical Guidelines

Strengths

- Rwanda follows WHO and ESMO recommendations in developing cancer treatment policies.
- Ongoing national plans aim to standardize cancer care pathways.



Weakness


- There are no specific national prostate cancer guidelines, leading to inconsistencies in care.
- Clinical practice heavily depends on individual provider knowledge or facility protocols.

Opportunity

- Develop context-appropriate prostate cancer guidelines, incorporating PSA and treatment pathways.
- Use digital platforms to disseminate guidelines and provide ongoing training.

Threats

- Lack of coordination across providers and weak referral systems can fragment patient care.
- Outdated protocols can persist if guideline updates are not institutionalized.







	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	✗	✗	✗	○
Feasibility of Integration	✗	✗	✗	✗	○
Adoption of International Guidelines	✗	✗	✗	✗	○
Engagement with Updates	✗	✗	✗	✗	○
ESMO Guidelines Implementation	✗	✗	✗	✗	○





















































































Rwanda



Reimbursement




-  A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
-  A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
-  No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States		
United Kingdom		
Canada		
Australia		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Japan		
South Korea		
China		
India		
Singapore		
Thailand		
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Rwanda



Colorectal Cancer Screening



Country	Colorectal Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Colorectal Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities