

# Colorectal Cancer Factsheet: Insights & Key Developments

Key Insights on Colorectal Cancer  
Care and Infrastructure

## Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Colorectal Cancer Screening

Colorectal cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping colorectal cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Among the top 3 cancers in Italian men.
- Incidence rate: Approximately 49 per 100,000 men per year.
- Total new cases (2022): Around 21,000 men.
- Daily diagnoses (2022): About 58 men per day.
- Deaths (2022): Approximately 9,800 men.
- 5-year survival rate: Estimated 65–70%, due to early detection.
- Most affected age group: Men aged 60–79.
- Screening participation: National screening program (biennial FIT for 50–69); participation is moderate to high depending on region.



# Italy

## Infrastructure



### Strengths

- Italy has a well-established national healthcare system (Servizio Sanitario Nazionale, SSN), offering equitable cancer services.
- Specialized cancer centers (IRCCS) and university hospitals provide comprehensive colorectal cancer treatment and diagnostics.

### Weakness



- Geographic disparities: Northern regions (e.g., Lombardy, Emilia-Romagna) have more advanced facilities than Southern regions (e.g., Calabria, Sicily).
- Waiting times for colonoscopy and surgery can be long, especially in under-resourced areas.

### Opportunity

- Continued investment in digital health infrastructure and centralized cancer registries to improve resource allocation.
- EU funding can support upgrades in oncology services in southern Italy.

### Threats


- Regionalized health administration sometimes leads to inconsistent care quality across the country.
- Aging infrastructure in older hospitals may not meet modern oncology standards.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		


# Italy

## Treatment Access, Research Funding and Awareness Campaigns



### Strengths

- Universal health coverage ensures access to standard colorectal cancer treatment, including surgery, chemotherapy, and radiotherapy.
- Italy is active in EU-funded oncology research, with several ongoing trials in precision medicine.



### Weakness











- Limited access to the newest targeted therapies in smaller cities or peripheral hospitals.
- Public campaigns on colorectal cancer awareness lag behind breast and prostate cancer initiatives.

























































































































### Opportunity

- Enhancing collaboration with European Cancer Mission projects can improve research capacity.
- Expanding media and digital awareness campaigns focused on colorectal cancer prevention and early signs

### Threats

- Budget limitations may hinder sustained research funding or rapid adoption of innovative treatments.
- Risk of unequal access to cutting-edge therapies based on hospital or region.


-   5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
-   4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
-   3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
-   2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
-   1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			



# Italy

## Survival Rates, Early Detection and Palliative Care



### Strengths

- Italy has one of the highest 5-year survival rates for colorectal cancer in Europe (above 60%), thanks to early detection and standardized treatment.
- Strong palliative care framework integrated into oncology units and supported by national policy.

### Weakness

- Late-stage diagnosis still occurs, especially in populations not reached by screening programs.
- Psychological and social support in palliative care is underdeveloped in non-urban regions.

### Opportunity

- Expanding early detection through mobile units and general practitioner engagement.
- Continued training of palliative care specialists and integration with community health services.

### Threats

- Rising aging population may increase demand for end-stage cancer care.
- Regional differences in palliative care access could exacerbate inequalities.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.






























































































































3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

# Italy

Utilization of Biomarkers



## Strengths

- Biomarker testing for KRAS, NRAS, and MSI is standard in metastatic colorectal cancer cases, guiding use of targeted therapies.
- Testing for BRAF V600E and PIK3CA is available in major centers and supported by pathology guidelines.

## Opportunity

- National guidelines promoting mandatory biomarker testing for all advanced colorectal cancer patients.
- Wider use of liquid biopsy and cost-effective genomic technologies can expand precision oncology.


## Weakness

- Access to comprehensive genomic profiling (NGS panels) may be limited to major cancer centers.
- Reimbursement for all biomarker tests is not uniformly enforced across regions.

## Threats

- Variability in biomarker testing infrastructure between regions may lead to unequal care.
- Over-reliance on local lab standards without external quality controls may risk diagnostic accuracy.

- Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.







# Italy



## Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	○	✗	✗	✗	✗
Feasibility of Integration	○	✗	✗	✗	✗
Adoption of International Guidelines	○	✗	✗	✗	✗
Engagement with Updates	✗	○	✗	✗	✗
ESMO Guidelines Implementation	○	✗	✗	✗	✗

# Indonesia

## Reimbursement



### Strengths

- SSN covers standard colorectal cancer diagnostics and treatment, including surgery, chemotherapy, and radiotherapy.
- Reimbursement for KRAS/NRAS biomarker testing is generally provided in metastatic settings.

### Weakness

- Some advanced targeted therapies and companion diagnostics may require case-by-case approval or delay.
- Regional health authority decisions can result in inconsistent reimbursement policies.

### Opportunity

- National reimbursement packages for next-generation sequencing and emerging therapies (e.g., immunotherapies).
- Use of value-based assessments to streamline coverage of high-cost drugs.

### Threats

- Economic constraints may impact reimbursement for high-cost drugs or delay listing of newer agents.
- Reimbursement rules may discourage hospitals from performing multiple biomarker tests.


- A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
- A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
- No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States	●	●
United Kingdom	●	●
Canada	●	●
Australia	●	●
Germany	●	●
France	●	●
Netherlands	●	●
Sweden	●	●
Italy	●	●
Spain	●	●
Poland	●	●
Japan	●	●
South Korea	●	●
China	●	●
India	●	●
Singapore	●	●
Thailand	●	●
South Africa	●	●
Kenya	●	●
Nigeria	●	●
Egypt	●	●
Morocco	●	●
Algeria	●	●
Ethiopia	●	●
Mexico	●	●
Brazil	●	●
Argentina	●	●
Chile	●	●
Colombia	●	●
New Zealand	●	●
Greece	●	●
Rwanda	●	●
Uganda	●	●
Serbia	●	●
Saudi Arabia	●	●
UAE	●	●
Syria	●	●
Indonesia	●	●
Vietnam	●	●
Philippines	●	●
Russia	●	●
Malaysia	●	●



# Indonesia

## Colorectal Cancer Screening



Country	Colorectal Cancer Screening
United States	Annual LDCT (50–80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55–74 years)
Canada	LDCT for high-risk individuals (55–74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50–74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Colorectal Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities