

Nigeria

Colorectal Cancer Factsheet: Insights & Key Developments

Key Insights on Colorectal Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Colorectal Cancer Screening

Colorectal cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping colorectal cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Colorectal cancer is increasing, now among top 5 digestive cancers in men.
- Incidence rate: Around 5 per 100,000 men per year.
- Total new cases (2022): Approximately 2,300 men.
- Daily diagnoses (2022): About 6 men per day.
- Deaths (2022): Around 2,000 men.
- 5-year survival rate: Likely under 30%, due to late diagnosis and limited treatment.
- Most affected age group: Mostly men aged 60 and older.
- Screening participation: No national program; screening is virtually absent.

Nigeria




Infrastructure



Strengths

- Major cancer centers like the **Lagos University Teaching Hospital** and **National Hospital Abuja** provide CRC care with surgery and limited oncology services.
- Growing adoption of digital health platforms for patient monitoring and referrals in urban centers.



Weakness


- Severe infrastructure deficits in rural areas; many regions lack CT, MRI, or endoscopy equipment.
- Frequent power outages, understaffing, and lack of multidisciplinary tumor boards impair care quality.












Opportunity

- Public-private infrastructure partnerships can upgrade diagnostic facilities and regional oncology hubs.
- Investment in mobile endoscopy and telepathology services for remote outreach.

Threats


- Continued reliance on external donors for major upgrades poses sustainability risks.
- Brain drain of trained oncologists and surgeons to higher-income countries.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Nigeria

Treatment Access, Research Funding and Awareness Campaigns



Strengths

- NGOs like Project PINK BLUE and Breast Without Spot engage in awareness campaigns for various cancers, including CRC.
- Pilot partnerships with global agencies support training of oncology nurses and pathologists.

Weakness

- CRC treatment costs are largely out-of-pocket; public hospitals often face drug and equipment shortages.
- Extremely limited CRC-specific research funding and clinical trials within Nigeria.

Opportunity

- Increased domestic philanthropy and diaspora investments could fund awareness and access programs.
- Regional collaboration (e.g., West African Health Organization) may help secure pooled research grants.

Threats

- Rising CRC incidence, especially among younger adults, without corresponding national investment.
- Misinformation and cancer stigma hinder early care-seeking behavior.


5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●
Malaysia	●	●	●

Nigeria



Survival Rates, Early Detection and Palliative Care



Strengths

- Urban centers like Ibadan and Abuja have established histopathology labs supporting earlier-stage diagnosis.
- Traditional and community health systems are increasingly engaged for early symptom referral.

Weakness

- Late-stage presentation is the norm; **>70%** of CRC cases are diagnosed at advanced stages.
- Very few structured palliative care services outside capital cities, with morphine availability being inconsistent.

Opportunity

- Education of primary care providers on CRC red-flag symptoms can boost early referral rates.
- Incorporation of home-based palliative models to reach underserved populations.

Threats

- Delayed diagnosis due to long waiting times, diagnostic bottlenecks, and lack of cancer awareness.
- Poverty and poor transportation limit follow-up after initial consultations.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.





















































































































3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Nigeria



Utilization of Biomarkers



Strengths

- Some tertiary centers have begun offering **KRAS and BRAF mutation** testing via research collaborations.
- Pathology labs in Lagos, Ibadan, and Port Harcourt are slowly incorporating IHC testing for MSI/dMMR.

Weakness

- Biomarker testing remains rare, expensive, and mostly research-driven; not standard in clinical pathways.
- No national mandate or reimbursement for molecular profiling in CRC.

Opportunity


- Regional lab networks or centralized genomic testing could reduce biomarker testing costs.
- Partnering with universities and private labs for biomarker-driven drug trials.

Threats

- High costs of test kits and reagents due to currency instability and import dependency.
- Risk of outdated treatment decisions due to unavailable molecular data.

- Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.


Nigeria



Nigeria



Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	✗	✗	○	✗
Feasibility of Integration	✗	✗	✗	○	✗
Adoption of International Guidelines	✗	✗	✗	○	✗
Engagement with Updates	✗	✗	✗	○	✗
ESMO Guidelines Implementation	✗	✗	✗	✗	○

Nigeria



Reimbursement



Strengths

- CRC surgery and select chemotherapy are included under **NHIS (National Health Insurance Scheme)** in certain pilot states.
- Faith-based and NGO hospitals sometimes subsidize cancer treatment.

Weakness

- Limited NHIS penetration (~10% of the population); most cancer care remains paid out-of-pocket.
- No reimbursement for diagnostics like colonoscopy, CT scan, or biomarker tests.

Opportunity

- Advocacy for cancer-specific funding within **NHIS expansion reforms** can broaden reimbursement scope.
- Emerging microinsurance and community-based health schemes could cover diagnostic costs.

Threats

- Unstable insurance funding mechanisms and delays in provider reimbursements reduce effectiveness.
- Economic constraints may further delay inclusion of new treatments or tests.


- A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
- A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
- No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States	●	●
United Kingdom	●	●
Canada	●	●
Australia	●	●
Germany	●	●
France	●	●
Netherlands	●	●
Sweden	●	●
Italy	●	●
Spain	●	●
Poland	●	●
Japan	●	●
South Korea	●	●
China	●	●
India	●	●
Singapore	●	●
Thailand	●	●
South Africa	●	●
Kenya	●	●
Nigeria	●	●
Egypt	●	●
Morocco	●	●
Algeria	●	●
Ethiopia	●	●
Mexico	●	●
Brazil	●	●
Argentina	●	●
Chile	●	●
Colombia	●	●
New Zealand	●	●
Greece	●	●
Rwanda	●	●
Uganda	●	●
Serbia	●	●
Saudi Arabia	●	●
UAE	●	●
Syria	●	●
Indonesia	●	●
Vietnam	●	●
Philippines	●	●
Russia	●	●
Malaysia	●	●

Nigeria



Colorectal Cancer Screening



Country	Colorectal Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Colorectal Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities