

United Kingdom

Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening


Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.

This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is not in the top 10, but still relevant in older men.
- Incidence rate: Around 7 per 100,000 men per year.
- Total new cases (2022): Around 3,500 men.
- Daily diagnoses: Roughly 10 men per day.
- Deaths (2022): About 2,500 men.
- 5-year survival rate: Estimated 35–45%.
- Most affected age group: Mostly 70 years and above.
- Screening participation: No routine screening; endoscopy performed only when symptoms arise.


United Kingdom

Infrastructure



Strengths

- Robust National Health Service (NHS) infrastructure with comprehensive cancer centres (e.g., Royal Marsden, Christie).
- Widespread access to modern diagnostic equipment (endoscopy, CT, PET-CT, MRI)



Weakness


- Long wait times for diagnostic endoscopy in some regions due to NHS backlog.
- Disparities in infrastructure between urban and rural areas or across devolved nations (England vs. Wales/Scotland/Northern Ireland).






















































Opportunity

- Investment in AI-assisted endoscopy and digital pathology to improve early gastric cancer diagnosis.
- Capital projects to upgrade cancer units under NHS Long-Term Plan.

Threats



- Workforce shortages (pathologists, radiologists) could bottleneck diagnostic capacity.
- Post-Brexit economic constraints may slow infrastructure expansion.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

United Kingdom

Treatment Access, Research Funding and Awareness Campaigns




5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	<div></div>	<div></div>	<div></div>
Kenya	<div></div>	<div></div>	<div></div>
Nigeria	<div></div>	<div></div>	<div></div>
Egypt	<div></div>	<div></div>	<div></div>
Morocco	<div></div>	<div></div>	<div></div>
Algeria	<div></div>	<div></div>	<div></div>
Ethiopia	<div></div>	<div></div>	<div></div>
India	<div></div>	<div></div>	<div></div>
Japan	<div></div>	<div></div>	<div></div>
South Korea	<div></div>	<div></div>	<div></div>
China	<div></div>	<div></div>	<div></div>
Thailand	<div></div>	<div></div>	<div></div>
Singapore	<div></div>	<div></div>	<div></div>
United Kingdom	<div></div>	<div></div>	<div></div>
Germany	<div></div>	<div></div>	<div></div>
France	<div></div>	<div></div>	<div></div>
Netherlands	<div></div>	<div></div>	<div></div>
Sweden	<div></div>	<div></div>	<div></div>
Italy	<div></div>	<div></div>	<div></div>
Spain	<div></div>	<div></div>	<div></div>
Poland	<div></div>	<div></div>	<div></div>
Mexico	<div></div>	<div></div>	<div></div>
Brazil	<div></div>	<div></div>	<div></div>
Argentina	<div></div>	<div></div>	<div></div>
Chile	<div></div>	<div></div>	<div></div>
Colombia	<div></div>	<div></div>	<div></div>
United States	<div></div>	<div></div>	<div></div>
Canada	<div></div>	<div></div>	<div></div>
Australia	<div></div>	<div></div>	<div></div>
New Zealand	<div></div>	<div></div>	<div></div>
Greece	<div></div>	<div></div>	<div></div>
Rwanda	<div></div>	<div></div>	<div></div>
Uganda	<div></div>	<div></div>	<div></div>
Serbia	<div></div>	<div></div>	<div></div>
Saudi Arabia	<div></div>	<div></div>	<div></div>
UAE	<div></div>	<div></div>	<div></div>
Syria	<div></div>	<div></div>	<div></div>
Indonesia	<div></div>	<div></div>	<div></div>
Vietnam	<div></div>	<div></div>	<div></div>
Philippines	<div></div>	<div></div>	<div></div>
Russia	<div></div>	<div></div>	<div></div>
Malaysia	<div></div>	<div></div>	<div></div>

United Kingdom

Survival Rates, Early Detection and Palliative Care



Strengths

- Structured cancer pathways (2-week referral) enable relatively timely diagnosis for suspected cancer.
- Excellent access to NHS-funded palliative care and hospice services.

Weakness



- Gastric cancer 5-year survival remains poor (~20%) due to late detection.
- Limited use of screening tools in asymptomatic populations.

Opportunity

- AI-assisted early detection in Barrett’s and gastric mucosal changes via advanced endoscopy.
- Expand primary care education to identify early alarm symptoms.

Threats


- Ageing population could increase caseload and worsen late-stage detection patterns.
- Stigma and denial may delay healthcare-seeking in some population segments.



Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

United Kingdom

Utilization of Biomarkers



Strengths

- HER2 testing is routinely done for advanced gastric cancer; trastuzumab is standard in eligible cases.
- UK centres participate in trials assessing PD-L1, MSI-H and emerging biomarkers (e.g., FIGHT, CheckMate trials).

Weakness


- Lack of routine testing for CLDN18.2 and FGFR2b limits stratification for newer therapies.
- Biomarker testing access may vary by Trust and region.




Opportunity


- NHS Genomic Medicine Service (GMS) provides a pathway for wider biomarker integration.
- Use of Real-World Evidence (RWE) from NHS datasets to evaluate biomarker utility.

Threats

- Reimbursement hurdles for novel targeted therapies like zolbetuximab (CLDN18.2).
- Rapid biomarker evolution could outpace NHS diagnostic capacity.




-  Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
-  Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
-  Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.



United Kingdom


Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	○	×	×	×	×
Feasibility of Integration	○	×	×	×	×
Adoption of International Guidelines	○	×	×	×	×
Engagement with Updates	×	○	×	×	×
ESMO Guidelines Implementation	○	×	×	×	×


United Kingdom

Reimbursement



United Kingdom

Colorectal Cancer Screening



Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Russia	No formal national LDCT program; regional pilot screening programs in large cities