

Prostate Cancer Factsheet: Insights & Key Developments

Key Insights on Prostate Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Prostate Cancer Screening

Prostate cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.

This factsheet provides a comprehensive overview of key pillars shaping Prostate cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- **Incidence share:** Most common cancer in Chilean men, accounting for roughly 20% of all male cancer diagnoses
- **Incidence rate:** Approximately 24.4 per 100,000 men per year
- **Total new cases (2022/2023):** Around 8,157 men diagnosed annually
- **Daily diagnoses:** Roughly 22 new cases per day
- **Deaths:** About 2,029 deaths per year
- **5-year survival rate:** Estimated around 57% (based on incidence vs mortality trends)
- **Most affected age group:** Primarily men aged 60 and above, with increasing rates over time
- **Screening participation:** No national organized screening program; opportunistic PSA testing common, with some regional outreach

Chile



Infrastructure

Strengths

- High-quality care centers like **Instituto Nacional del Cáncer (INC)** and major university hospitals (e.g., **Hospital Clínico Universidad de Chile, PUC, FACH**) offer **comprehensive prostate cancer care**.
- Radiotherapy and urological surgery available in all regions with **cancer-designated public hospitals**.
- GES (AUGE) plan legally guarantees access to prostate cancer diagnosis and treatment

Opportunity



- Expand **satellite oncology centers** and train local personnel in underserved regions.
- Invest in tele-urology and support early detection pathways in municipal clinics.

Weakness

- Significant **urban-rural divide** in access to specialized care—**northern and southern regions** (e.g., Arica, Aysén) have fewer trained specialists.
- **Shortage of radiotherapy machines** leads to treatment delays in certain areas.
- Limited access to **robotic-assisted surgery**, concentrated in Santiago.

Threats

- **Health worker strikes or public system overload** may disrupt treatment continuity.
- Public-private gaps risk **fragmented patient journeys**.



- 5. Advanced nationwide infrastructure, widespread availability in public and private sectors, integration with clinical practice.
- 4. Strong infrastructure in major hospitals and cancer centers, some regional disparities.
- 3. Moderate infrastructure, primarily in private settings or research institutions.
- 2. Limited infrastructure, available only in select centers or for high-cost private testing.
- 1. Minimal or no infrastructure, testing mostly unavailable or sent abroad.

Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Chile



Treatment Access, Research Funding and Awareness Campaigns

Strengths

- Prostate cancer is part of GES (Explicit Guarantees in Health): state guarantees diagnosis within 45 days and treatment within 60 days.
- Active urology research groups and participation in Latin American oncology trials.
- Annual public awareness campaigns supported by the Ministry of Health and groups like Fundación Cáncer Vida.

Opportunity



- Promote clinical trial decentralization to regional hospitals.
- Strengthen Men’s Health and preventive cancer education through PHC campaigns.

Weakness












































































































- Delays in implementation of GES deadlines due to high patient load and capacity limits.
- Access to next-generation hormone therapies (enzalutamide, abiraterone) is still limited in public hospitals.
- Research output is concentrated in a few urban academic hubs.

Threats

- Economic disparities affect care continuity for patients transitioning between public and private systems.
- Over-reliance on urban infrastructure for high-cost drug access.



- 5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
- 4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
- 3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
- 2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
- 1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Chile



Survival Rates, Early Detection and Palliative Care

Strengths

- Early-stage patients treated under GES show **5-year survival >90%**.
- Chile has a strong palliative care model, integrated into both **public and home-based services**.
- Coverage of **pain management and hospice care** included under public benefits.

Opportunity



- Embed **early symptom alerts and PSA checks** in PHC visits.
- Strengthen community-based palliative education for caregivers.

Weakness

- Still, ~35–40% of cases are diagnosed at an advanced stage, especially in men without routine PSA testing.
- Geriatric and end-of-life care models not fully scaled in **rural health posts**.

Threats

- Ageing population may increase demand for palliative oncology beyond current capacity.
- **Cultural stigma among older men** still limits help-seeking behavior.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.




























































































































3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.


Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Chile

Utilization of Biomarkers





- Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.



Chile



Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	○	✗	✗	✗
Feasibility of Integration	✗	○	✗	✗	✗
Adoption of International Guidelines	✗	○	✗	✗	✗
Engagement with Updates	✗	✗	○	✗	✗
ESMO Guidelines Implementation	✗	○	✗	✗	✗

Chile



Reimbursement



Strengths

- GES/AUGE ensures free coverage of prostate cancer diagnosis and treatment timelines for **all public-insured citizens (FONASA)**.
- Private insurers (ISAPRES) also cover prostate cancer under **explicit legal guarantees**.

Weakness

- Gaps in coverage for **advanced therapies, especially for castration-resistant prostate cancer (CRPC)**.
- Approval delays and **bureaucracy hinder access to newer agents** in public system.

Opportunity

- Add **next-generation hormone therapy** to the explicit benefits list (GES).
- Pilot **value-based reimbursement** programs for innovative treatments.

Threats

- Economic fluctuations or policy shifts could **jeopardize continuity of full coverage**.
- High drug prices strain public sector's ability to provide equitable access.


- A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
- A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
- No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States	●	●
United Kingdom	●	●
Canada	●	●
Australia	●	●
Germany	●	●
France	●	●
Netherlands	●	●
Sweden	●	●
Italy	●	●
Spain	●	●
Poland	●	●
Japan	●	●
South Korea	●	●
China	●	●
India	●	●
Singapore	●	●
Thailand	●	●
South Africa	●	●
Kenya	●	●
Nigeria	●	●
Egypt	●	●
Morocco	●	●
Algeria	●	●
Ethiopia	●	●
Mexico	●	●
Brazil	●	●
Argentina	●	●
Chile	●	●
Colombia	●	●
New Zealand	●	●
Greece	●	●
Rwanda	●	●
Uganda	●	●
Serbia	●	●
Saudi Arabia	●	●
UAE	●	●
Syria	●	●
Indonesia	●	●
Vietnam	●	●
Philippines	●	●
Russia	●	●
Malaysia	●	●

Chile



Prostate Cancer Screening



Country	Prostate Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Prostate Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities