

Netherlands

Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening

Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is not among top male cancers, but incidence is higher in older adults.
- Incidence rate: Around 7–8 per 100,000 men per year.
- Total new cases (2022): Around 2,000 men.
- Daily diagnoses: About 5–6 men per day.
- Deaths (2022): Approximately 1,300 men.
- 5-year survival rate: Estimated 30–40%.
- Most affected age group: Mostly men aged 70+.
- Screening participation: No national screening; endoscopy is performed only when clinically indicated.

Netherlands



Infrastructure



Strengths

- The Netherlands has a highly organized healthcare system with advanced cancer centers like the Netherlands Cancer Institute (NKI) and University Medical Centers (e.g., UMC Utrecht, Erasmus MC).
- National electronic health records and integrated diagnostic pathways support streamlined referrals, diagnostics, and treatment planning.

Weakness


- Specialized gastric cancer care is centralized to a limited number of high-volume centers, which may delay treatment initiation for patients in peripheral regions.
- Limited endoscopy capacity in some areas causes delays in early diagnosis, especially post-COVID

Opportunity

- Expand telepathology and remote tumor boards to enhance access for patients in non-academic hospitals.
- Increase training in gastrointestinal oncology and surgical specialization to reduce dependency on central hubs.

Threats


- Increasing pressure on hospital capacity due to an aging population and staff shortages, particularly in surgical and pathology departments.
- Delayed upgrades in regional hospitals due to budget caps and public-private funding constraints.



Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Netherlands

Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Strong government and EU-level funding for cancer research (e.g., Dutch Cancer Society, Horizon Europe projects).
- Universal healthcare coverage ensures equitable access to chemotherapy, surgery, and radiotherapy.

Weakness


- Public awareness of gastric cancer remains low, especially in comparison to breast or colorectal cancer.
- Access to clinical trials is often limited to major academic hospitals, reducing opportunity for patients in community settings.

Opportunity

















































































































- Boost funding for targeted awareness campaigns on H. pylori and gastric cancer risk factors.
- Leverage Dutch research strengths in precision medicine and AI-driven diagnostics to improve gastric cancer outcomes.

Threats

- Rising costs of oncology care and new drugs may strain reimbursement systems.
- Patient advocacy is weaker in gastric cancer compared to other cancer types, potentially affecting fundraising and policy influence.




- 5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
- 4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
- 3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
- 2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
- 1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Netherlands

Survival Rates, Early Detection and Palliative Care



Strengths

- 5-year relative survival rates for localized gastric cancer can reach over 60% when detected early and treated in high-volume centers.
- Strong palliative care infrastructure, including home-based and hospice care integrated into the national health system.

Weakness


- Most gastric cancers in the Netherlands are diagnosed at an advanced stage, often stage III or IV.
- Early detection programs for gastric cancer are non-existent, and H. pylori testing is not routine.

Opportunity

- Introduce risk-based screening pilots for high-risk populations (e.g., migrants from high-incidence regions).
- Expand non-invasive testing pathways, such as stool-based markers or serological risk panels.

Threats

- Continued absence of national gastric cancer screening may prevent significant survival improvement.
- Delayed symptom recognition among both GPs and patients, due to vague early signs and low awareness.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.



3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			
Malaysia			

Netherlands

Utilization of Biomarkers



Strengths

- HER2 testing is routine in advanced gastric cancer, and trastuzumab is reimbursed for HER2+ patients.
- PD-L1 testing (CPS≥1) and MSI testing are increasingly implemented to guide immunotherapy use.

Opportunity

- Expand biomarker panels through national cancer genomics initiatives like the Hartwig Medical Foundation.
- Integrate comprehensive molecular profiling into routine advanced-stage gastric cancer workup.


Weakness

- Limited adoption of newer biomarkers like CLDN18.2 and FGFR2b in standard clinical workflows.
- Variability in testing access across smaller or non-academic hospitals, especially for less common biomarkers.

Threats



- Reimbursement delays for newer biomarker tests and therapies may limit their uptake.
- Risk of disparity in access between urban and rural hospitals or academic vs. community centers.

- Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.



Netherlands


Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	○	×	×	×	×
Feasibility of Integration	○	×	×	×	×
Adoption of International Guidelines	○	×	×	×	×
Engagement with Updates	×	×	○	×	×
ESMO Guidelines Implementation	○	×	×	×	×

Netherlands

Reimbursement




- A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
- A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
- No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States	●	●
United Kingdom	●	●
Canada	●	●
Australia	●	●
Germany	●	●
France	●	●
Netherlands	●	●
Sweden	●	●
Italy	●	●
Spain	●	●
Poland	●	●
Japan	●	●
South Korea	●	●
China	●	●
India	●	●
Singapore	●	●
Thailand	●	●
South Africa	●	●
Kenya	●	●
Nigeria	●	●
Egypt	●	●
Morocco	●	●
Algeria	●	●
Ethiopia	●	●
Mexico	●	●
Brazil	●	●
Argentina	●	●
Chile	●	●
Colombia	●	●
New Zealand	●	●
Greece	●	●
Rwanda	●	●
Uganda	●	●
Serbia	●	●
Saudi Arabia	●	●
UAE	●	●
Syria	●	●
Indonesia	●	●
Vietnam	●	●
Philippines	●	●
Russia	●	●
Malaysia	●	●

Netherlands

Gastric Cancer Screening



Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities