

United States

Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening


Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.

This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is relatively rare, not in the top male cancers.
- Incidence rate: Around 6 per 100,000 men per year.
- Total new cases (2022): Around 13,000–14,000 men.
- Daily diagnoses: About 35–38 men per day.
- Deaths (2022): Around 9,500 men.
- 5-year survival rate: Estimated 32% overall; higher for early-stage cases.
- Most affected age group: Mostly men aged 70+.
- Screening participation: No population-based screening; diagnosis occurs after symptom onset.

United States

Infrastructure



Strengths

- Extensive network of NCI-designated cancer centers (e.g., MD Anderson, Memorial Sloan Kettering).
- Advanced diagnostic capabilities with widespread access to endoscopy, imaging, and pathology services.

Weakness


- Urban-rural healthcare disparities affect access to timely diagnostic and treatment services.
- Fragmented healthcare system leads to variability in care quality across regions and insurance types.

Opportunity

- Expansion of telehealth and mobile health units for underserved areas.
- Infrastructure modernization through federal cancer moonshot and NIH-backed initiatives.

Threats

- Rising healthcare costs may limit future infrastructure investments.
- Unequal access to technology (AI diagnostics, molecular imaging) across institutions




Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

United States



Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Leading country in gastric cancer clinical trials and translational research.
- NIH and NCI provide strong financial backing for cancer research, including GI oncology.

Weakness


- Low public awareness of gastric cancer risk factors and early symptoms.
- Access to advanced treatments heavily influenced by insurance coverage.

Opportunity

- Launch targeted awareness campaigns for high-risk groups (e.g., Asian American and Hispanic populations).
- Increase funding for rare cancer subtypes and improve clinical trial diversity.

Threats

- Underinsurance or lack of coverage in certain populations hinders treatment continuity.
- Economic inequality may limit patient participation in clinical trials or access to high-cost therapies.




- 5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
- 4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
- 3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
- 2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
- 1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●
Malaysia	●	●	●

United States



Survival Rates, Early Detection and Palliative Care



Strengths

- Access to high-quality palliative and hospice care services nationwide.
- Multidisciplinary care models in comprehensive cancer centers improve patient outcomes.

Weakness

- Overall 5-year survival rate for gastric cancer remains low (~32%), mainly due to late-stage diagnoses.
- Lack of routine early detection strategies outside high-risk cohorts.

Opportunity

- AI-assisted endoscopy and liquid biopsy research to detect cancer earlier.
- Enhance symptom-based screening in primary care and urgent care settings.

Threats

- Disparities in early detection and survival outcomes by race, ethnicity, and socioeconomic status.
- Delays in diagnosis due to healthcare fragmentation or insurance denials.




- 5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.
- 4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.
- 3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.
- 2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.
- 1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●
Malaysia	●	●	●

United States

Utilization of Biomarkers



Strengths

- HER2, MSI-H, PD-L1 testing is widely adopted in major institutions for gastric cancer management.
- FDA-approved drugs available for HER2+, MSI-H, and PD-L1+ cases (e.g., pembrolizumab, trastuzumab).

Weakness




- CLDN18.2 and FGFR2b testing not yet standard; may be limited to clinical trials or major centers.
- Limited biomarker testing in community hospitals due to cost or infrastructure gaps.

Opportunity


- CLDN18.2 and FGFR2b testing not yet standard; may be limited to clinical trials or major centers.
- Limited biomarker testing in community hospitals due to cost or infrastructure gaps.

Threats

- High costs of molecular testing and therapies may limit equitable adoption.
- Insurance may not cover newer biomarker tests or off-label drug use.



-  Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
-  Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
-  Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.

United States



United States


Clinical Guidelines




















































































	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	○	×	×	×	×
Feasibility of Integration	○	×	×	×	×
Adoption of International Guidelines	○	×	×	×	×
Engagement with Updates	×	○	×	×	×
ESMO Guidelines Implementation	○	×	×	×	×

United States

Reimbursement




-  A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
-  A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
-  No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States		
United Kingdom		
Canada		
Australia		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Japan		
South Korea		
China		
India		
Singapore		
Thailand		
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

United States

Colorectal Cancer Screening



Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities