

# Germany

## Breast Cancer Factsheet: Insights & Key Developments

Key Insights on Breast Cancer Care  
and Infrastructure

### Core Pillars:

1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Breast Cancer Screening

Breast cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping breast cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Breast cancer incidence: Approximately 74,512 new cases annually, accounting for a significant proportion of all female cancers.
- Incidence rate: 117.5 per 100,000 women per year.
- Total cases (2022): Approximately 74,512 reported cases.
- Lifetime risk in Germany: Approximately 1 in 8 women.
- Daily diagnoses (2022): Approximately 204 women per day.
- Breast cancer deaths (2023): Approximately 18,527 deaths.
- 5-year survival rate: 88%.
- 10-year survival rate: 83%.
- Most affected age group: Women aged 50 to 69 years.
- Mammogram detection: Approximately 6 out of 1,000 women screened are diagnosed with breast cancer.
- Screening participation (ages 50-69): Approximately 51% of invited women participate in the German breast cancer screening program.



# Germany

## Infrastructure



### Strengths

- Universal access to HER2 therapies through public coverage
- Strong clinical trial infrastructure and public engagement initiatives

### Weakness



- Advanced therapies and clinical trials more accessible in urban centers
- Variability in implementation of awareness campaigns across regions










































### Opportunity

- Expand clinical trials and new therapies to community hospitals
- Strengthen targeted campaigns for underrepresented populations

### Threats

- Access delays for innovative treatments in lower-volume hospitals
- Resource concentration in large institutions could exacerbate disparities




Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		

# Germany



## Treatment Access, Research Funding and Awareness Campaigns



### Strengths

- Five-year breast cancer survival rate exceeds 88%
- National screening and palliative care systems are well-established

### Weakness


- Screening participation remains around 50%, below EU targets
- End-of-life and psycho-oncological services uneven in rural areas

### Opportunity

- Boost outreach to improve screening participation and early detection
- Expand access to outpatient and community-based palliative care

### Threats

- Regional disparities may affect quality-of-life outcomes
- Demographic shifts could increase pressure on supportive services




5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●




# Germany

## Survival Rates, Early Detection and Palliative Care



### Strengths

- 5-year breast cancer survival exceeds 90%—among the world’s highest.
- Palliative care services are nationally integrated and expanding.



### Weakness

- Screening participation remains ~55%, below the WHO 70% target.
- Rural regions have fewer palliative care specialists and services.

### Opportunity

- Increase outreach and screening participation through personalized invitations.
- Train more rural GPs and nurses in early detection and palliative support.

### Threats

- Aging population may increase demand for end-of-life care.
- Lower participation in vulnerable populations may limit national gains.



5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.



4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.


























































































































3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.



2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.




1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa			
Kenya			
Nigeria			
Egypt			
Morocco			
Algeria			
Ethiopia			
India			
Japan			
South Korea			
China			
Thailand			
Singapore			
United Kingdom			
Germany			
France			
Netherlands			
Sweden			
Italy			
Spain			
Poland			
Mexico			
Brazil			
Argentina			
Chile			
Colombia			
United States			
Canada			
Australia			
New Zealand			
Greece			
Rwanda			
Uganda			
Serbia			
Saudi Arabia			
UAE			
Syria			
Indonesia			
Vietnam			
Philippines			
Russia			

# Germany

## Utilization of Biomarkers




### Strengths

- HER2, ER, PR, and BRCA testing widely available and reimbursed
- NGS and expanded panels increasingly used in major centers

### Weakness

- Regional disparities in advanced biomarker testing and genetic counseling
- Gaps in integration of newer technologies in all care settings



### Opportunity


- Standardize access through national genomic platforms
- Strengthen education on precision medicine for clinicians

### Threats

- Uneven biomarker use could delay access to targeted therapies
- Resource limitations in non-academic hospitals may hinder uptake

- 5. 80% Biomarker testing is widely available and routinely performed as part of standard clinical practice. Strong integration into treatment decisions, with national coverage and reimbursement ensuring accessibility.
- 4. 61-80%. Biomarker testing is commonly used, but access may be limited in certain regions or patient groups. Some disparities exist in coverage or affordability, but it is still a crucial part of cancer diagnostics
- 3. 41-60% Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- 2. 20-40% Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- 1. <20% Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.


### Germany





# Germany

## Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	○	×	×	×	×
Feasibility of Integration	×	○	×	×	×
Adoption of International Guidelines	○	×	×	×	×
Engagement with Updates	○	×	×	×	×
ESMO Guidelines Implementation	○	×	×	×	×

# Germany

## Reimbursement



### Strengths

- Robust, structured reimbursement under SHI for diagnostics and therapies
- HER2-targeted treatments fully covered, including in early stages

### Weakness

- Bureaucratic complexity in approval of novel therapies
- Differences in reimbursement navigation between institutions

### Opportunity

- Streamline fast-track access to innovative therapies
- Enhance transparency in pricing and patient access pathways

### Threats

- Financial sustainability may be challenged by rising innovation costs
- Delays in access due to administrative bottlenecks



Yes - A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.



Partial - A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.



No - No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement	No-cost Access
South Africa	○	✗
Kenya	✗	✗
Nigeria	✗	✗
Egypt	○	◐
Morocco	○	✗
Algeria	○	✗
Ethiopia	✗	✗
India	◐	✗
Japan	○	○
South Korea	○	○
China	○	◐
Thailand	○	◐
Singapore	○	○
United Kingdom	○	○
Germany	○	○
France	○	○
Netherlands	○	○
Sweden	○	○
Italy	○	○
Spain	○	○
Poland	○	◐
Mexico	○	✗
Brazil	○	✗
Argentina	○	✗
Chile	○	◐
Colombia	○	✗
United States	○	○
Canada	○	○
Australia	○	○
New Zealand	○	○
Greece	○	◐
Rwanda	✗	✗
Uganda	✗	✗
Serbia	○	◐
Saudi Arabia	○	○
UAE	○	○
Syria	○	○
Indonesia	○	◐
Vietnam	✗	✗
Philippines	✗	✗
Russia	○	◐




# Germany

## Breast Cancer Screening

### Strengths

- National screening program reimbursed for women aged 50–75
- High detection rates in screened populations



### Weakness

- Participation rate ~50%, below EU benchmark
- Screening less accessible to high-risk younger populations

### Opportunity

- Expand mobile screening units and local outreach strategies
- Adopt risk-adapted screening models for improved targeting

### Threats

- Low uptake among vulnerable groups may delay diagnosis
- Cultural and regional barriers to participation persist

Country	Breast Cancer Screening
United States	Biennial mammograms (50-74 years)
United Kingdom	Triennial mammograms (50-71 years)
Canada	Mammograms every 2-3 years (50-74 years)
Australia	Biennial mammograms (50-74 years)
Germany	Mammograms every 2 years (50-69 years)
France	Biennial mammograms (50-74 years)
Netherlands	Mammograms every 2 years (50-75 years)
Sweden	Mammograms every 18-24 months (40-74 years)
Italy	Mammograms every 2 years (50-69 years)
Spain	Mammograms every 2 years (50-69 years)
Poland	Mammograms every 2 years (50-69 years)
Japan	Mammograms every 2 years (40+ years)
South Korea	Biennial mammograms (40+ years)
China	Regional mammogram programs (40-69 years)
India	Opportunistic screening
Singapore	Biennial mammograms (50-69 years)
Saudi Arabia	Opportunistic screening; regional programs for women aged 40+
UAE	Opportunistic screening; encouraged every 2 years for 40-69 years
Syria	No national program; limited local initiatives due to conflict

Country	Breast Cancer Screening
Thailand	Biennial mammograms (50-69 years)
South Africa	Opportunistic screening
Kenya	No national program
Nigeria	No national program
Egypt	National awareness campaigns
Morocco	National program for 45-69 years
Algeria	Planned national program (50-69 years)
Ethiopia	No national program
Mexico	Biennial mammograms (40-69 years)
Brazil	Biennial mammograms (50-69 years)
Argentina	Biennial mammograms (50-69 years)
Chile	Mammograms every 3 years (50-69 years)
Colombia	Biennial mammograms (50-69 years)
New Zealand	Biennial mammograms (45-69 years)
Greece	Biennial mammograms (50-69 years)
Rwanda	No national program
Uganda	No national program
Serbia	Biennial mammograms (50-69 years)
Indonesia	Opportunistic screening; no national mammography program
Vietnam	Regional mammography programs; pilot programs in urban areas (age 45-69)
Philippines	Opportunistic screening; mammography recommended every 2 years for women 50+
Russia	National program for biennial mammograms (50-69 years)