

Gastric Cancer Factsheet: Insights & Key Developments

Key Insights on Gastric Cancer
Care and Infrastructure

Core Pillars:


1. Infrastructure
2. Treatment Access, Research Funding and Awareness Campaigns
3. Survival Rates, Early Detection and Palliative Care
4. Utilization of Biomarkers
5. Clinical Guidelines
6. Reimbursement
7. Gastric Cancer Screening

Gastric cancer remains one of the most prevalent cancers worldwide, affecting millions of individuals each year. Despite advancements in diagnostics, treatment, and awareness, disparities in access to care, molecular testing, and specialized centers persist.


This factsheet provides a comprehensive overview of key pillars shaping Gastric cancer care, including specialized infrastructure, treatment accessibility, research funding, early detection, and palliative care.

- Incidence share: Gastric cancer is not among the top digestive cancers in Kenyan men.
- Incidence rate: Approximately 4 per 100,000 men per year.
- Total new cases (2022): Around 600–700 men.
- Daily diagnoses: Roughly 2 men per day.
- Deaths (2022): About 500–600 men.
- 5-year survival rate: Likely < 40%, with many late-stage presentations.
- Most affected age group: Men aged 60 and older.
- Screening participation: None; diagnosis usually occurs at advanced stages.

Kenya



Infrastructure



Strengths

- Existence of referral hospitals such as Kenyatta National Hospital and Moi Teaching & Referral Hospital offering oncology services.
- Public-private partnerships (e.g., Aga Khan University Hospital) provide access to diagnostics and surgery in major cities.

Opportunity



- Expansion of County Cancer Centres as part of Universal Health Coverage (UHC) agenda.
- Telemedicine and mobile diagnostic vans for early detection in rural counties.

Weakness

- Severe shortages of oncologists (only ~35 medical oncologists in the country).
- Rural infrastructure gaps: Most cancer services centralized in Nairobi, Kisumu, Eldoret—limiting reach to 75%+ of the population.

Threats

- Equipment breakdowns (e.g., radiotherapy machines), electricity outages, and lack of maintenance limit service delivery.
- Growing cancer burden may overwhelm under-resourced infrastructure.




Country	Specialized Centers	Genetic & Molecular Testing Infrastructure
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
India		
Japan		
South Korea		
China		
Thailand		
Singapore		
United Kingdom		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
United States		
Canada		
Australia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Kenya



Treatment Access, Research Funding and Awareness Campaigns



Strengths

- Recent National Cancer Control Strategy (2023–2027) emphasizes access, affordability, and awareness.
- Civil society and NGOs (e.g., Faraja Cancer Support, Hope for Cancer Kids) lead local awareness and patient support

Weakness

- Most Kenyans pay out-of-pocket; NHIF covers some treatments but excludes high-cost drugs and biomarker testing.
- Minimal local gastric cancer research or funding from national institutions.

Opportunity

- Increased global funding (e.g., WHO, IAEA) for oncology scale-up can support early detection and training.
- Public education on risk factors like H. pylori, tobacco, diet could reduce incidence.

Threats

- Cultural stigma and late health-seeking behavior hinder awareness efforts.
- Economic instability reduces donor and government cancer funding.


5. Strong healthcare infrastructure with comprehensive treatment access, high research funding, and nationwide awareness campaigns. Patients have access to advanced therapies, clinical trials, and widespread early detection programs.
4. Well-developed system with good treatment availability, strong research funding, and effective but regionally focused awareness campaigns. Some disparities may exist in rural areas or between public and private sectors.
3. Moderate development, with specialized treatments available in major hospitals, research funding concentrated on specific cancers, and occasional but limited awareness efforts. Healthcare access may be restricted by cost or geography.
2. Limited system where cancer treatment is available only in select urban centers, research funding is minimal or sporadic, and awareness campaigns are rare or underfunded. Patients often face long wait times or financial barriers.
1. Poor infrastructure with severe barriers to treatment, little to no research funding, and lack of structured awareness campaigns. Cancer care is largely inaccessible, with many patients relying on out-of-pocket expenses or external aid.

Country	Treatment Access	Research Funding	Awareness Campaigns
South Africa	<div></div>	<div></div>	<div></div>
Kenya	<div></div>	<div></div>	<div></div>
Nigeria	<div></div>	<div></div>	<div></div>
Egypt	<div></div>	<div></div>	<div></div>
Morocco	<div></div>	<div></div>	<div></div>
Algeria	<div></div>	<div></div>	<div></div>
Ethiopia	<div></div>	<div></div>	<div></div>
India	<div></div>	<div></div>	<div></div>
Japan	<div></div>	<div></div>	<div></div>
South Korea	<div></div>	<div></div>	<div></div>
China	<div></div>	<div></div>	<div></div>
Thailand	<div></div>	<div></div>	<div></div>
Singapore	<div></div>	<div></div>	<div></div>
United Kingdom	<div></div>	<div></div>	<div></div>
Germany	<div></div>	<div></div>	<div></div>
France	<div></div>	<div></div>	<div></div>
Netherlands	<div></div>	<div></div>	<div></div>
Sweden	<div></div>	<div></div>	<div></div>
Italy	<div></div>	<div></div>	<div></div>
Spain	<div></div>	<div></div>	<div></div>
Poland	<div></div>	<div></div>	<div></div>
Mexico	<div></div>	<div></div>	<div></div>
Brazil	<div></div>	<div></div>	<div></div>
Argentina	<div></div>	<div></div>	<div></div>
Chile	<div></div>	<div></div>	<div></div>
Colombia	<div></div>	<div></div>	<div></div>
United States	<div></div>	<div></div>	<div></div>
Canada	<div></div>	<div></div>	<div></div>
Australia	<div></div>	<div></div>	<div></div>
New Zealand	<div></div>	<div></div>	<div></div>
Greece	<div></div>	<div></div>	<div></div>
Rwanda	<div></div>	<div></div>	<div></div>
Uganda	<div></div>	<div></div>	<div></div>
Serbia	<div></div>	<div></div>	<div></div>
Saudi Arabia	<div></div>	<div></div>	<div></div>
UAE	<div></div>	<div></div>	<div></div>
Syria	<div></div>	<div></div>	<div></div>
Indonesia	<div></div>	<div></div>	<div></div>
Vietnam	<div></div>	<div></div>	<div></div>
Philippines	<div></div>	<div></div>	<div></div>
Russia	<div></div>	<div></div>	<div></div>
Malaysia	<div></div>	<div></div>	<div></div>

Kenya



Survival Rates, Early Detection and Palliative Care



Strengths

- Growing focus on palliative care, supported by Kenya Hospices and Palliative Care Association (KEHPCA).
- Increasing training of palliative nurses and counselors through national programs.

Opportunity


- Scale-up of early H. pylori screening and treatment in high-incidence counties (e.g., Rift Valley, Central).
- Train community health workers in basic symptom screening and referral.

Weakness

- Survival rate for gastric cancer is <15%, largely due to late diagnosis.
- No national screening program, and most cases are detected at Stage III or IV.

Threats


- Poor diagnostic access, especially pathology turnaround times (2–4 weeks), delays staging and treatment.
- Low availability of pain medications and poor palliation in remote regions.




- 5. High survival rates, strong early detection programs, and well-established palliative care services. Patients have access to timely diagnosis, advanced treatments, and comprehensive end-of-life care.
- 4. Good survival rates, effective early detection efforts, and accessible but regionally limited palliative care. Some disparities may exist in rural areas or for specific cancer types.
- 3. Moderate survival rates, early detection available but not widespread, and palliative care services mainly in urban centers. Some patients experience delays in diagnosis or limited end-of-life care.
- 2. Low survival rates, early detection efforts are inconsistent or underfunded, and palliative care is minimal or only available in select hospitals. Cancer patients face significant access barriers.
- 1. Very low survival rates, poor early detection infrastructure, and almost no palliative care services. Many patients are diagnosed late and lack proper support for pain management and end-of-life care.

Country	Survival Rates	Early Detection	Palliative Care
South Africa	●	●	●
Kenya	●	●	●
Nigeria	●	●	●
Egypt	●	●	●
Morocco	●	●	●
Algeria	●	●	●
Ethiopia	●	●	●
India	●	●	●
Japan	●	●	●
South Korea	●	●	●
China	●	●	●
Thailand	●	●	●
Singapore	●	●	●
United Kingdom	●	●	●
Germany	●	●	●
France	●	●	●
Netherlands	●	●	●
Sweden	●	●	●
Italy	●	●	●
Spain	●	●	●
Poland	●	●	●
Mexico	●	●	●
Brazil	●	●	●
Argentina	●	●	●
Chile	●	●	●
Colombia	●	●	●
United States	●	●	●
Canada	●	●	●
Australia	●	●	●
New Zealand	●	●	●
Greece	●	●	●
Rwanda	●	●	●
Uganda	●	●	●
Serbia	●	●	●
Saudi Arabia	●	●	●
UAE	●	●	●
Syria	●	●	●
Indonesia	●	●	●
Vietnam	●	●	●
Philippines	●	●	●
Russia	●	●	●
Malaysia	●	●	●

Kenya



Utilization of Biomarkers



Strengths

- HER2 testing occasionally available in private labs (e.g., Lancet Labs, Pathologists Lancet Kenya).
- Global clinical trials (especially HER2+) increasingly enrolling from sub-Saharan Africa, including Kenya.

Weakness

- Biomarker testing is unaffordable for most patients and not available in public hospitals.
- Lack of national laboratory capacity for advanced genomic profiling (e.g., NGS).

Opportunity


- Collaboration with international biopharma and NGOs to improve access to testing and targeted therapy.
- Inclusion of HER2 and PD-L1 testing in NHIF-covered diagnostic packages.

Threats


- Delayed diagnosis and inability to perform timely biomarker testing makes targeted therapy ineffective.
- Dependence on imported reagents and supply chain delays.

- Moderate utilization, often restricted to major hospitals or private healthcare settings. Some patients may not receive biomarker testing due to cost or limited availability in public healthcare systems.
- Biomarker testing is available but underutilized, with significant barriers such as high costs, lack of awareness, or limited infrastructure. Many patients may not receive recommended biomarker assessments.
- Biomarker testing is rarely performed, often due to lack of infrastructure, awareness, or financial barriers. Patients typically do not receive targeted therapies based on biomarker status.



Kenya



Kenya



Clinical Guidelines



	Very High	High	Medium	Low	Very Low
Clinical Guideline Implementation	✗	✗	✗	○	✗
Feasibility of Integration	✗	✗	✗	○	✗
Adoption of International Guidelines	✗	✗	✗	○	✗
Engagement with Updates	✗	✗	✗	✗	○
ESMO Guidelines Implementation	✗	✗	✗	✗	○

Kenya



Reimbursement



Strengths

- NHIF offers basic oncology care, including chemotherapy, surgery, and radiotherapy at designated centers.
- Recent NHIF reforms are increasing outpatient cancer care packages.

Weakness




- NHIF does not reimburse biomarker testing or targeted therapies (e.g., trastuzumab, nivolumab).
- Private insurance is unaffordable for most Kenyans; many patients go without full treatment.





















































































Opportunity

- NHIF strategic purchasing from private labs and treatment centers can broaden access.
- Pilot outcomes-based reimbursement models for select targeted therapies.

Threats

- Rising cancer treatment costs may lead to benefit package cuts.
- Informal sector populations often not enrolled in NHIF, limiting their cancer coverage

-  A structured reimbursement system exists, ensuring biomarker testing is covered through national healthcare systems, insurance, or public-private partnerships. Patients face no direct financial burden.
-  A reimbursement framework is in place, but patients may still have out-of-pocket expenses such as co-pays, limited coverage, or financial caps on testing.
-  No formal reimbursement system exists, meaning patients must fully cover the cost of biomarker testing out-of-pocket.

Country	Reimbursement Framework	No-cost Access
United States		
United Kingdom		
Canada		
Australia		
Germany		
France		
Netherlands		
Sweden		
Italy		
Spain		
Poland		
Japan		
South Korea		
China		
India		
Singapore		
Thailand		
South Africa		
Kenya		
Nigeria		
Egypt		
Morocco		
Algeria		
Ethiopia		
Mexico		
Brazil		
Argentina		
Chile		
Colombia		
New Zealand		
Greece		
Rwanda		
Uganda		
Serbia		
Saudi Arabia		
UAE		
Syria		
Indonesia		
Vietnam		
Philippines		
Russia		
Malaysia		

Kenya



Gastric Cancer Screening



Country	Gastric Cancer Screening
United States	Annual LDCT (50-80 years, high-risk smokers)
United Kingdom	LDCT for high-risk individuals (55-74 years)
Canada	LDCT for high-risk individuals (55-74 years)
Australia	No national program, high-risk groups advised LDCT
Germany	No national program, under evaluation
France	No national LDCT screening
Netherlands	Participating in European screening studies
Sweden	No national LDCT screening
Italy	Regional pilot LDCT screening
Spain	No national LDCT program
Poland	No national program
Japan	No national LDCT program
South Korea	LDCT for high-risk individuals (50-74 years)
China	No national LDCT program
India	No national LDCT program
Singapore	No national LDCT program
Saudi Arabia	No national LDCT program; some hospital-based opportunistic screening
UAE	No national LDCT program; early-stage pilot studies ongoing in select hospitals
Syria	No national LDCT program; screening not prioritized due to conflict
Malaysia	No program; high-risk CT pilots

Country	Gastric Cancer Screening
Thailand	No national LDCT program
South Africa	No national LDCT program
Kenya	No national LDCT program
Nigeria	No national LDCT program
Egypt	No national LDCT program
Morocco	No national LDCT program
Algeria	No national LDCT program
Ethiopia	No national LDCT program
Mexico	No national LDCT program
Brazil	No national LDCT program
Argentina	No national LDCT program
Chile	No national LDCT program
Colombia	No national LDCT program
New Zealand	No national LDCT program
Greece	No national LDCT program
Rwanda	No national LDCT program
Uganda	No national LDCT program
Serbia	No national LDCT program
Indonesia	No national LDCT program; opportunistic screening in private sector
Vietnam	No national LDCT program; early pilot screening studies in Hanoi and Ho Chi Minh
Philippines	No national LDCT program; feasibility and awareness programs under discussion
Russia	No formal national LDCT program; regional pilot screening programs in large cities