

Social Networks: Assignment #1

Due on Monday, December 15, 2025

Dr. Masoud Asadpour 15:00am

Ali Dashtbozorg 810104302

Contents

1		4
(a)	Consider the below network.	4
(b)	Consider the below network.	5
(c)	Consider an undirected tree of n nodes. A particular edge in the tree joins node 1 and 2 and divides the tree into two disjoint regions of n_1 and n_2 nodes, as sketched below:	6

List of Figures

Question 1

Distances and Neighbors

(a) Consider the below network.

For each of the following scenarios, indicate which node would be the best choice, giving reasons:

1. The mayor wants to install a radio broadcast station so that, in a crisis, a single nationwide message can reach all areas. The goal is that every node's distance to the station (independently of other nodes) is as small as possible—in other words, the maximum distance from any node to the station should be minimized.

To determine the optimal location for the radio station, we identify the **Graph Center** by computing the **Eccentricity** of each node. The eccentricity of a node is the distance to the farthest node in the network; minimizing this ensures that the maximum travel time for a broadcast signal is as small as possible.

The eccentricity vector ϵ for nodes 1 through 15 is:

$$\epsilon = \begin{bmatrix} 4.0 \\ 3.0 \\ 4.0 \\ 5.0 \\ 4.0 \\ 4.0 \\ 4.0 \\ 5.0 \\ 4.0 \\ 4.0 \\ 4.0 \\ 5.0 \\ 3.0 \\ 4.0 \\ 5.0 \end{bmatrix}$$

The minimum eccentricity is $\min(\epsilon) = 3.0$, which occurs at **Node 2** and **Node 13**. Therefore, placing the radio station at either of these nodes ensures that every other node is reachable within at most 3 hops, providing the most efficient coverage across the network.

2. The mayor wants to choose a node for a bookstore so that the sum of distances from all residents to that node is minimized.

To determine the optimal node for placing the bookstore, we use **closeness centrality**, which measures how close a node is on average to all other nodes in the graph. By choosing the node with the highest closeness centrality, we ensure that the nodes can reach the bookstore as quickly as possible on average.

The formula for closeness centrality of a node v is:

$$C(v) = \frac{n-1}{\sum_{u \in V} d(v, u)}$$

where $d(v, u)$ is the shortest-path distance between nodes v and u , and n is the total number of nodes.

For our 15-node graph, the closeness centrality values are:

$$\begin{bmatrix} 0.4667 & 0.4667 & 0.4667 & 0.4 & 0.4516 \\ 0.4516 & 0.4667 & 0.4242 & 0.4516 & 0.4375 \\ 0.4118 & 0.4 & 0.5185 & 0.3684 & 0.2745 \end{bmatrix}$$

Node $i = 13$ (value 0.5185) has the highest closeness centrality, so we choose it as the location for the bookstore to ensure efficient distance to other nodes.

3. Two stores have decided to open new branches in the city. Each person (node) buys from the nearest store. If a person is at equal distance from both stores, their purchases are split equally between them. First, select the best node to open store A, then determine the best location for store B given that choice.

From the closeness centrality analysis, we know that node 13 has the highest value, so it is the best choice for opening store A. Given store A at node 13, the nodes that would maximize the number of customers for store B are those with the highest customer counts. In this case, nodes 1, 3, 7, and 8 each attract 7.0 customers, making them all equally good candidates for store B.

The number of customers each node would get if store A is at node 13 is shown below, with the best candidates for store B highlighted in bold:

1	7.0
2	6.0
3	7.0
4	6.5
5	6.5
6	5.5
7	7.0
8	7.0
9	6.5
10	5.0
11	5.5
12	6.5
14	2.0
15	1.5

Therefore, the recommended locations are: store A at node 13, and store B at any of nodes 1, 3, 7, or 8.

(b) Consider the below network.

1. What is the closeness centrality for node C ?

The closeness centrality(both standard and normalized) for node C are as follows:

$$C_C(c) = \frac{1}{\sum_{j=1}^n d_{cj}} = \frac{1}{n-1}$$

$$\tilde{C}_C(c) = \frac{n-1}{\sum_{j=1}^n d_{cj}} = \frac{n-1}{n-1} = 1$$

2. Derive the closeness centrality value for the nodes on the ring as a function of n

The closeness centrality(both standard and normalized) for nodes other than node C are as follows:

$$C_C(n_i) = \frac{1}{\sum_{j=1}^n d_{ij}} = \frac{1}{1+1+1+(n-4)(2)} = \frac{1}{2n-5}$$

$$\tilde{C}_C(n_i) = \frac{n-1}{\sum_{j=1}^n d_{ij}} = \frac{n-1}{1+1+1+(n-4)(2)} = \frac{n-1}{2n-5}$$

for any node on the ring the 2 nodes beside it and the hub node have a distance of 1 and the rest $n-4$ nodes have a distance of 2 which goes through the hub

3. Derive the betweenness centrality value for node C .

we know that C is in every shortest path except for the neighbors of each node so 2 nodes are gone and we also know that for neighbors of our neighbors we have 2 paths which one goes through C so 2 nodes are gone and for the rest of the paths they all go through C so we would have $n-1$ nodes on the ring which have the following equation:

$$\delta_{uv}(n_i) = \frac{\sigma_{uv}(n_i)}{\sigma_{uv}}$$

$$C_B(n_i) = \sum_{u \neq n_i} \sum_{v \neq n_i} \delta_{uv}(n_i)$$

$$\delta_{uv}(C) = \begin{cases} 0 & d_{uv} = 1 \\ \frac{1}{2} & d_{uv} = 2 \\ 1 & d_{uv} > 2 \end{cases} \quad \text{where } C \notin d_{uv}$$

$$C_B(C) = \frac{1}{2}(n-1)(0+0+0+\frac{1}{2}+\frac{1}{2}+(n-6)) = \frac{(n-1)(n-5)}{2}$$

(c) Consider an undirected tree of n nodes. A particular edge in the tree joins node 1 and 2 and divides the tree into two disjoint regions of n_1 and n_2 nodes, as sketched below:

Show that the closeness centrality C_1 and C_2 of the two nodes are related by:

$$\frac{1}{C_1} + \frac{n_1}{n} = \frac{1}{C_2} + \frac{n_2}{n}$$

The proof goes as follows:

$$C_C(1) = \frac{n}{\sum_{j=1}^n d_{1j}} = \frac{n}{\sum_{j=1}^{n_1} d_{1j} + n_2 + \sum_{j=n_1+1}^{n_1+n_2} d_{2j}}$$

$$C_C(2) = \frac{n}{\sum_{j=1}^n d_{2j}} = \frac{n}{\sum_{j=n_1+1}^{n_1+n_2} d_{2j} + n_1 + \sum_{j=1}^{n_1} d_{1j}}$$

$$\frac{n}{C_C(1)} = n_2 + \sum_{j=1}^{n_1} d_{1j} + \sum_{j=n_1+1}^{n_1+n_2} d_{2j}$$

$$\frac{n}{C_C(2)} = n_1 + \sum_{j=n_1+1}^{n_1+n_2} d_{2j} + \sum_{j=1}^{n_1} d_{1j}$$

$$\frac{n}{C_C(1)} - n_2 = \frac{n}{C_C(2)} - n_1$$

$$\frac{1}{C_C(1)} - \frac{n_2}{n} = \frac{1}{C_C(2)} - \frac{n_1}{n}$$

$$\frac{1}{C_C(1)} + \frac{n_1}{n} = \frac{1}{C_C(2)} + \frac{n_2}{n}$$

$$\frac{1}{C_1} + \frac{n_1}{n} = \frac{1}{C_2} + \frac{n_2}{n}$$