
Old_survival − GO (BP / MF / CC)
chemotaxis
taxis
leukocyte migration
cell chemotaxis
humoral immune response
response to molecule of bacterial origin
response to lipopolysaccharide
antimicrobial humoral response
cellular response to lipopolysaccharide
molecule of bacterial origin
leukocyte chemotaxis
myeloid leukocyte activation
cellular response to biotic stimulus
antimicrobial humoral immune response mediated by
myeloid leukocyte migration
positive regulation of killing of cells of another organism
leukocyte migration
disruption of cell in another organism
neutrophil migration
BP
extracellular matrix
external encapsulating structure
cytoplasmic vesicle lumen
vesicle lumen
endoplasmic reticulum lumen
secretory granule lumen
specific granule
secretory granule membrane
tertiary granule
external side of plasma membrane
vacuolar lumen
ficolin−1−rich granule
ficolin−1−rich granule membrane
specific granule lumen
collagen trimer
specific granule membrane
tertiary granule lumen
immunoglobulin complex
CC
cytokine activity
cytokine receptor binding
G protein−coupled receptor binding
chemokine activity
chemokine receptor binding
glycosaminoglycan binding
immune receptor activity
extracellular matrix structural constituent
sulfur compound binding
growth factor binding
growth factor activity
serine−type endopeptidase activity
serine−type peptidase activity
serine hydrolase activity
heparin binding
cytokine receptor activity
CXCR chemokine receptor binding
tumor necrosis factor receptor binding
CCR chemokine receptor binding
tumor necrosis factor receptor superfamily binding
MF
0.04
0.08
0.12
0.16
GeneRatio
p.adjust
1e−13
1e−09
1e−05
Count
5
10
15
20