

Topic: Vector orthogonal to the plane

Question: Find the vector orthogonal to the plane that includes the given vectors.

$$\overrightarrow{AB} = \langle 2, 3, 2 \rangle$$

$$\overrightarrow{AC} = \langle -2, 1, 2 \rangle$$

Answer choices:

A $\langle 1, -2, 2 \rangle$

B $\langle 1, 2, 2 \rangle$

C $\langle 1, -2, -2 \rangle$

D $\langle 1, 2, -2 \rangle$



Solution: A

Given two vectors that lie in the plane, we can find the vector orthogonal to the plane by taking the cross product of the vectors $\overrightarrow{AB} = \langle AB_1, AB_2, AB_3 \rangle$ and $\overrightarrow{AC} = \langle AC_1, AC_2, AC_3 \rangle$.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ AB_1 & AB_2 & AB_3 \\ AC_1 & AC_2 & AC_3 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} \begin{vmatrix} AB_2 & AB_3 \\ AC_2 & AC_3 \end{vmatrix} - \mathbf{j} \begin{vmatrix} AB_1 & AB_3 \\ AC_1 & AC_3 \end{vmatrix} + \mathbf{k} \begin{vmatrix} AB_1 & AB_2 \\ AC_1 & AC_2 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} (AB_2 AC_3 - AB_3 AC_2) - \mathbf{j} (AB_1 AC_3 - AB_3 AC_1) + \mathbf{k} (AB_1 AC_2 - AB_2 AC_1)$$

For the given vectors $\overrightarrow{AB} = \langle 2, 3, 2 \rangle$ and $\overrightarrow{AC} = \langle -2, 1, 2 \rangle$, we get

$$\overrightarrow{AB} \times \overrightarrow{AC} = [(3)(2) - (2)(1)] \mathbf{i} - [(2)(2) - (2)(-2)] \mathbf{j} + [(2)(1) - (3)(-2)] \mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = (6 - 2)\mathbf{i} - (4 + 4)\mathbf{j} + (2 + 6)\mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = 4\mathbf{i} - 8\mathbf{j} + 8\mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = 4(\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$$

We'll convert this into standard vector form to get the vector orthogonal to the plane.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \langle 1, -2, 2 \rangle$$



Topic: Vector orthogonal to the plane

Question: Find the vector orthogonal to the plane that includes the given points.

$$A(1, -1, 1)$$

$$B(-2, 2, -2)$$

$$C(3, 1, -2)$$

Answer choices:

A $\langle -1, 5, -4 \rangle$

B $\langle 1, -5, -4 \rangle$

C $\langle 1, 5, 4 \rangle$

D $\langle -1, -5, 4 \rangle$



Solution: C

Since we've been given three points $A(a_1, a_2, a_3)$, $B(b_1, b_2, b_3)$ and $C(c_1, c_2, c_3)$, we'll use the points to generate two vectors, \overrightarrow{AB} and \overrightarrow{AC} .

$$\overrightarrow{AB} = (b_1 - a_1)\mathbf{i} + (b_2 - a_2)\mathbf{j} + (b_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AB} = (AB_1)\mathbf{i} + (AB_2)\mathbf{j} + (AB_3)\mathbf{k}$$

and

$$\overrightarrow{AC} = (c_1 - a_1)\mathbf{i} + (c_2 - a_2)\mathbf{j} + (c_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AC} = (AC_1)\mathbf{i} + (AC_2)\mathbf{j} + (AC_3)\mathbf{k}$$

Plugging in the points we've been given, we get

$$\overrightarrow{AB} = (b_1 - a_1)\mathbf{i} + (b_2 - a_2)\mathbf{j} + (b_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AB} = (-2 - 1)\mathbf{i} + [2 - (-1)]\mathbf{j} + (-2 - 1)\mathbf{k}$$

$$\overrightarrow{AB} = -3\mathbf{i} + 3\mathbf{j} - 3\mathbf{k}$$

and

$$\overrightarrow{AC} = (c_1 - a_1)\mathbf{i} + (c_2 - a_2)\mathbf{j} + (c_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AC} = (3 - 1)\mathbf{i} + [1 - (-1)]\mathbf{j} + (-2 - 1)\mathbf{k}$$

$$\overrightarrow{AC} = 2\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$$

If we convert these into standard vector form, we get

$$AB\langle AB_1, AB_2, AB_3 \rangle = \langle -3, 3, -3 \rangle$$



$$AC\langle AC_1, AC_2, AC_3 \rangle = \langle 2, 2, -3 \rangle$$

Now that we've converted the points into vectors, we'll take the cross product of $\overrightarrow{AB} = \langle AB_1, AB_2, AB_3 \rangle$ and $\overrightarrow{AC} = \langle AC_1, AC_2, AC_3 \rangle$.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ AB_1 & AB_2 & AB_3 \\ AC_1 & AC_2 & AC_3 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} \begin{vmatrix} AB_2 & AB_3 \\ AC_2 & AC_3 \end{vmatrix} - \mathbf{j} \begin{vmatrix} AB_1 & AB_3 \\ AC_1 & AC_3 \end{vmatrix} + \mathbf{k} \begin{vmatrix} AB_1 & AB_2 \\ AC_1 & AC_2 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} (AB_2 AC_3 - AB_3 AC_2) - \mathbf{j} (AB_1 AC_3 - AB_3 AC_1) + \mathbf{k} (AB_1 AC_2 - AB_2 AC_1)$$

For the vectors $\overrightarrow{AB} = \langle -3, 3, -3 \rangle$ and $\overrightarrow{AC} = \langle 2, 2, -3 \rangle$, we get

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} [(3)(-3) - (-3)(2)] - \mathbf{j} [(-3)(-3) - (-3)(2)] + \mathbf{k} [(-3)(2) - (3)(2)]$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} [-9 - (-6)] - \mathbf{j} [9 - (-6)] + \mathbf{k} [-6 - 6]$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = -3\mathbf{i} - 15\mathbf{j} - 12\mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = -3(\mathbf{i} + 5\mathbf{j} + 4\mathbf{k})$$

We'll convert this into standard vector form to get the vector orthogonal to the plane.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \langle 1, 5, 4 \rangle$$



Topic: Vector orthogonal to the plane

Question: Find the vector orthogonal to the plane that includes the given points.

$$A(4, -5, -4)$$

$$B(7, -5, 0)$$

$$C(11, 6, 8)$$

Answer choices:

A $\langle -44, 8, 33 \rangle$

B $\langle -44, -8, 33 \rangle$

C $\langle 7, 11, 12 \rangle$

D $\langle 7, 11, -12 \rangle$



Solution: B

Since we've been given three points $A(a_1, a_2, a_3)$, $B(b_1, b_2, b_3)$ and $C(c_1, c_2, c_3)$, we'll use the points to generate two vectors, \overrightarrow{AB} and \overrightarrow{AC} .

$$\overrightarrow{AB} = (b_1 - a_1)\mathbf{i} + (b_2 - a_2)\mathbf{j} + (b_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AB} = (AB_1)\mathbf{i} + (AB_2)\mathbf{j} + (AB_3)\mathbf{k}$$

and

$$\overrightarrow{AC} = (c_1 - a_1)\mathbf{i} + (c_2 - a_2)\mathbf{j} + (c_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AC} = (AC_1)\mathbf{i} + (AC_2)\mathbf{j} + (AC_3)\mathbf{k}$$

Plugging in the points we've been given, we get

$$\overrightarrow{AB} = (b_1 - a_1)\mathbf{i} + (b_2 - a_2)\mathbf{j} + (b_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AB} = (7 - 4)\mathbf{i} + [-5 - (-5)]\mathbf{j} + [0 - (-4)]\mathbf{k}$$

$$\overrightarrow{AB} = 3\mathbf{i} + 0\mathbf{j} + 4\mathbf{k}$$

and

$$\overrightarrow{AC} = (c_1 - a_1)\mathbf{i} + (c_2 - a_2)\mathbf{j} + (c_3 - a_3)\mathbf{k}$$

$$\overrightarrow{AC} = (11 - 4)\mathbf{i} + [6 - (-5)]\mathbf{j} + [8 - (-4)]\mathbf{k}$$

$$\overrightarrow{AC} = 7\mathbf{i} + 11\mathbf{j} + 12\mathbf{k}$$

If we convert these into standard vector form, we get

$$AB\langle AB_1, AB_2, AB_3 \rangle = \langle 3, 0, 4 \rangle$$



$$AC\langle AC_1, AC_2, AC_3 \rangle = \langle 7, 11, 12 \rangle$$

Now that we've converted the points into vectors, we'll take the cross product of $\overrightarrow{AB} = \langle AB_1, AB_2, AB_3 \rangle$ and $\overrightarrow{AC} = \langle AC_1, AC_2, AC_3 \rangle$.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ AB_1 & AB_2 & AB_3 \\ AC_1 & AC_2 & AC_3 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} \begin{vmatrix} AB_2 & AB_3 \\ AC_2 & AC_3 \end{vmatrix} - \mathbf{j} \begin{vmatrix} AB_1 & AB_3 \\ AC_1 & AC_3 \end{vmatrix} + \mathbf{k} \begin{vmatrix} AB_1 & AB_2 \\ AC_1 & AC_2 \end{vmatrix}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = \mathbf{i} (AB_2 AC_3 - AB_3 AC_2) - \mathbf{j} (AB_1 AC_3 - AB_3 AC_1) + \mathbf{k} (AB_1 AC_2 - AB_2 AC_1)$$

For the vectors $\overrightarrow{AB} = \langle 3, 0, 4 \rangle$ and $\overrightarrow{AC} = \langle 7, 11, 12 \rangle$, we get

$$\overrightarrow{AB} \times \overrightarrow{AC} = [(0)(12) - (4)(11)] \mathbf{i} - [(3)(12) - (4)(7)] \mathbf{j} + [(3)(11) - (0)(7)] \mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = (0 - 44)\mathbf{i} - (36 - 28)\mathbf{j} + (33 - 0)\mathbf{k}$$

$$\overrightarrow{AB} \times \overrightarrow{AC} = -44\mathbf{i} - 8\mathbf{j} + 33\mathbf{k}$$

We'll convert this into standard vector form to get the vector orthogonal to the plane.

$$\overrightarrow{AB} \times \overrightarrow{AC} = \langle -44, -8, 33 \rangle$$

