

Topic: Curvature

Question: Find the curvature of the vector function.

$$r(t) = \cos t \mathbf{i} + \sin t \mathbf{j} + 2t \mathbf{k}$$

Answer choices:

A $\kappa(t) = \frac{1}{\sqrt{5}}$

B $\kappa(t) = \sqrt{5}$

C $\kappa(t) = \frac{1}{5}$

D $\kappa(t) = 5$



Solution: C

First we'll find the derivative of the vector function.

$$r(t) = \cos t \mathbf{i} + \sin t \mathbf{j} + 2t \mathbf{k}$$

$$r'(t) = -\sin t \mathbf{i} + \cos t \mathbf{j} + 2 \mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|r'(t)| = \sqrt{[r'(t)_1]^2 + [r'(t)_2]^2 + [r'(t)_3]^2}$$

$$|r'(t)| = \sqrt{(-\sin t)^2 + (\cos t)^2 + (2)^2}$$

$$|r'(t)| = \sqrt{(\sin^2 t + \cos^2 t) + 4}$$

$$|r'(t)| = \sqrt{(1) + 4}$$

$$|r'(t)| = \sqrt{5}$$

And now we can use everything we just found to find the unit tangent vector.

$$T(t) = \frac{r'(t)}{|r'(t)|}$$

$$T(t) = \frac{-\sin t \mathbf{i} + \cos t \mathbf{j} + 2 \mathbf{k}}{\sqrt{5}}$$

$$T(t) = -\frac{\sin t}{\sqrt{5}} \mathbf{i} + \frac{\cos t}{\sqrt{5}} \mathbf{j} + \frac{2}{\sqrt{5}} \mathbf{k}$$



$$T(t) = -\frac{1}{\sqrt{5}} \sin t \mathbf{i} + \frac{1}{\sqrt{5}} \cos t \mathbf{j} + \frac{2}{\sqrt{5}} \mathbf{k}$$

To find the curvature of the vector function, we'll first need to find the derivative of this unit tangent vector.

$$T(t) = -\frac{1}{\sqrt{5}} \sin t \mathbf{i} + \frac{1}{\sqrt{5}} \cos t \mathbf{j} + \frac{2}{\sqrt{5}} \mathbf{k}$$

$$T'(t) = -\frac{1}{\sqrt{5}} \cos t \mathbf{i} - \frac{1}{\sqrt{5}} \sin t \mathbf{j} + (0) \mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|T'(t)| = \sqrt{[T'(t)_1]^2 + [T'(t)_2]^2 + [T'(t)_3]^2}$$

$$|T'(t)| = \sqrt{\left(-\frac{1}{\sqrt{5}} \cos t\right)^2 + \left(-\frac{1}{\sqrt{5}} \sin t\right)^2 + (0)^2}$$

$$|T'(t)| = \sqrt{\frac{1}{5} \cos^2 t + \frac{1}{5} \sin^2 t}$$

$$|T'(t)| = \sqrt{\frac{1}{5} (\cos^2 t + \sin^2 t)}$$

$$|T'(t)| = \sqrt{\frac{1}{5}(1)}$$

$$|T'(t)| = \frac{1}{\sqrt{5}}$$



And now we can use everything we just found to solve for the curvature of the vector function.

$$\kappa(t) = \frac{|T'(t)|}{|r'(t)|}$$

$$\kappa(t) = \frac{\frac{1}{\sqrt{5}}}{\sqrt{5}}$$

$$\kappa(t) = \frac{1}{\sqrt{5}\sqrt{5}}$$

$$\kappa(t) = \frac{1}{5}$$

This is the curvature of the vector function.



Topic: Curvature

Question: Find the curvature of the vector function.

$$r(t) = 4t\mathbf{i} - 3\cos t\mathbf{j} - 3\sin t\mathbf{k}$$

Answer choices:

A $\kappa(t) = \frac{1}{25}$

B $\kappa(t) = \frac{3}{5}$

C $\kappa(t) = \frac{1}{5}$

D $\kappa(t) = \frac{3}{25}$



Solution: D

First we'll find the derivative of the vector function.

$$r(t) = 4t\mathbf{i} - 3 \cos t\mathbf{j} - 3 \sin t\mathbf{k}$$

$$r'(t) = 4\mathbf{i} + 3 \sin t\mathbf{j} - 3 \cos t\mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|r'(t)| = \sqrt{[r'(t)_1]^2 + [r'(t)_2]^2 + [r'(t)_3]^2}$$

$$|r'(t)| = \sqrt{(4)^2 + (3 \sin t)^2 + (-3 \cos t)^2}$$

$$|r'(t)| = \sqrt{16 + 9 \sin^2 t + 9 \cos^2 t}$$

$$|r'(t)| = \sqrt{16 + 9 (\sin^2 t + \cos^2 t)}$$

$$|r'(t)| = \sqrt{16 + 9(1)}$$

$$|r'(t)| = \sqrt{25}$$

$$|r'(t)| = 5$$

And now we can use everything we just found to find the unit tangent vector.

$$T(t) = \frac{r'(t)}{|r'(t)|}$$

$$T(t) = \frac{4\mathbf{i} + 3 \sin t\mathbf{j} - 3 \cos t\mathbf{k}}{5}$$



$$T(t) = \frac{4}{5}\mathbf{i} + \frac{3}{5}\sin t\mathbf{j} - \frac{3}{5}\cos t\mathbf{k}$$

To find the curvature of the vector function, we'll first need to find the derivative of this unit tangent vector.

$$T(t) = \frac{4}{5}\mathbf{i} + \frac{3}{5}\sin t\mathbf{j} - \frac{3}{5}\cos t\mathbf{k}$$

$$T'(t) = (0)\mathbf{i} + \frac{3}{5}\cos t\mathbf{j} + \frac{3}{5}\sin t\mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|T'(t)| = \sqrt{[T'(t)_1]^2 + [T'(t)_2]^2 + [T'(t)_3]^2}$$

$$|T'(t)| = \sqrt{(0)^2 + \left(\frac{3}{5}\cos t\right)^2 + \left(\frac{3}{5}\sin t\right)^2}$$

$$|T'(t)| = \sqrt{\frac{9}{25}\cos^2 t + \frac{9}{25}\sin^2 t}$$

$$|T'(t)| = \sqrt{\frac{9}{25}(\cos^2 t + \sin^2 t)}$$

$$|T'(t)| = \sqrt{\frac{9}{25}(1)}$$

$$|T'(t)| = \frac{3}{5}$$

And now we can use everything we just found to solve for the curvature of the vector function.



$$\kappa(t) = \frac{|T'(t)|}{|r'(t)|}$$

$$\kappa(t) = \frac{\frac{3}{5}}{5}$$

$$\kappa(t) = \frac{3}{5 \cdot 5}$$

$$\kappa(t) = \frac{3}{25}$$

This is the curvature of the vector function.



Topic: Curvature

Question: Find the curvature of the vector function.

$$r(t) = 6 \sin t \mathbf{i} - 8t \mathbf{j} + 6 \cos t \mathbf{k}$$

Answer choices:

A $\kappa(t) = \frac{3}{5}$

B $\kappa(t) = \frac{3}{50}$

C $\kappa(t) = \frac{3}{25}$

D $\kappa(t) = \frac{3}{10}$



Solution: B

First we'll find the derivative of the vector function.

$$r(t) = 6 \sin t \mathbf{i} - 8t \mathbf{j} + 6 \cos t \mathbf{k}$$

$$r'(t) = 6 \cos t \mathbf{i} - 8 \mathbf{j} - 6 \sin t \mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|r'(t)| = \sqrt{[r'(t)_1]^2 + [r'(t)_2]^2 + [r'(t)_3]^2}$$

$$|r'(t)| = \sqrt{(6 \cos t)^2 + (-8)^2 + (-6 \sin t)^2}$$

$$|r'(t)| = \sqrt{36 \cos^2 t + 64 + 36 \sin^2 t}$$

$$|r'(t)| = \sqrt{64 + 36 (\cos^2 t + \sin^2 t)}$$

$$|r'(t)| = \sqrt{64 + 36(1)}$$

$$|r'(t)| = \sqrt{100}$$

$$|r'(t)| = 10$$

And now we can use everything we just found to find the unit tangent vector.

$$T(t) = \frac{r'(t)}{|r'(t)|}$$

$$T(t) = \frac{6 \cos t \mathbf{i} - 8 \mathbf{j} - 6 \sin t \mathbf{k}}{10}$$



$$T(t) = \frac{6}{10} \cos t \mathbf{i} - \frac{8}{10} \mathbf{j} - \frac{6}{10} \sin t \mathbf{k}$$

$$T(t) = \frac{3}{5} \cos t \mathbf{i} - \frac{4}{5} \mathbf{j} - \frac{3}{5} \sin t \mathbf{k}$$

To find the curvature of the vector function, we'll first need to find the derivative of this unit tangent vector.

$$T(t) = \frac{3}{5} \cos t \mathbf{i} - \frac{4}{5} \mathbf{j} - \frac{3}{5} \sin t \mathbf{k}$$

$$T'(t) = -\frac{3}{5} \sin t \mathbf{i} - (0) \mathbf{j} - \frac{3}{5} \cos t \mathbf{k}$$

Then we'll find the magnitude of this derivative.

$$|T'(t)| = \sqrt{[T'(t)_1]^2 + [T'(t)_2]^2 + [T'(t)_3]^2}$$

$$|T'(t)| = \sqrt{\left(-\frac{3}{5} \sin t\right)^2 + (0)^2 + \left(-\frac{3}{5} \cos t\right)^2}$$

$$|T'(t)| = \sqrt{\frac{9}{25} \sin^2 t + \frac{9}{25} \cos^2 t}$$

$$|T'(t)| = \sqrt{\frac{9}{25} (\sin^2 t + \cos^2 t)}$$

$$|T'(t)| = \sqrt{\frac{9}{25} (1)}$$

$$|T'(t)| = \frac{3}{5}$$



And now we can use everything we just found to solve for the curvature of the vector function.

$$\kappa(t) = \frac{|T'(t)|}{|r'(t)|}$$

$$\kappa(t) = \frac{\frac{3}{5}}{10}$$

$$\kappa(t) = \frac{3}{5 \cdot 10}$$

$$\kappa(t) = \frac{3}{50}$$

This is the curvature of the vector function.

