

PC2232 Physics for Electrical Engineers: Tutorial 6

Question 1: Harmonic oscillator ground state

For the quantum harmonic oscillator, the potential of the system is given by

$$U(x) = \frac{1}{2}m\omega^2x^2. \quad (1)$$

Suppose that $\psi(x)$ is a solution to the quantum harmonic oscillator, i.e., the Schrödinger equation is satisfied:

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + \frac{1}{2}m\omega^2x^2\psi = E\psi. \quad (2)$$

- (a) Show that a new wavefunction $\phi(x)$ defined by

$$\phi = \left(\hbar \frac{d}{dx} + m\omega x \right) \psi \quad (3)$$

Is also a solution to the Schrödinger equation of energy $E' = E - \hbar\omega$. That is, ϕ is a state of energy lower than ψ by a value of $\hbar\omega$.

- (b) The ground state ψ_0 is defined as a state of the lowest possible energy. That means, if we use the above method to find a lower energy state ϕ by calculating

$$\phi = \left(\hbar \frac{d}{dx} + m\omega x \right) \psi_0, \quad (4)$$

does not exist. Therefore, set $\phi = 0$ and solve Eq. (4) to derive the ground state wavefunction of the harmonic oscillator.

Question 2: Step potential

Particles of energy E are incident from the left, where $U(x) = 0$, and at the origin encounter an abrupt drop in potential energy, whose depth is $-3E$.

- (a) Classically what would the particles do, and what would happen to their kinetic energy?
- (b) Applying quantum mechanics: Assume an incident wave of the form $\psi_{\text{inc}}(x) = e^{ikx}$, where the normalization constant has been given the simple value of 1. Determine completely the wave function everywhere, including numeric values for the multiplicative constants.
- (c) What is the probability that incident particles will be reflected?

Question 3: Transmission resonance

In the $E > U_0$ potential barrier, there should be no reflected wave when the incident wave is at one of the transmission resonances. Prove this by assuming that a beam of particles is incident at the first transmission resonance, $E = U_0 + \frac{\pi^2 \hbar^2}{2mL^2}$ prove that this is true by utilizing the continuity conditions. *Hint:* Your aim is to show that $B = 0$. (Note: k' is particularly simple in this special case.)

Question 4: Two-dimensional well

Consider an infinite, two-dimensional square well given by the potential

$$U(x, y) = \begin{cases} 0, & \text{if } 0 < x < L, \quad 0 < y < L \\ \infty, & \text{otherwise.} \end{cases} \quad (5)$$

- (a) Solve the Schrödinger equation and find the possible wave functions.
- (b) What are the allowed energies E ?
- (c) List the five lowest energy levels. For each level, write down the number of different states that correspond to that energy.

Question 5: Tunneling

An electron with initial kinetic energy 5.5 eV encounters a square potential barrier with height 10.0 eV. What is the width of the barrier if the electron has a 0.10% probability of tunneling through the barrier? What is the energy of the electron after it tunnels through the barrier?