**SRM**  
INSTITUTE OF SCIENCE & TECHNOLOGY  
(Established as Sri Ramakrishna Engineering College, 1983)

---

# CHEMISTRY

32<sup>nd</sup> and 33<sup>rd</sup> Class, 26-10-2021


Dr. K. Ananthanarayanan  
Associate Professor (Research)  
Department of Chemistry  
Room No 319, 3<sup>rd</sup> Floor, Research Building

Email : [ananthak@srmist.edu.in](mailto:ananthak@srmist.edu.in)  
Phone : 9840154665

---

18CYB101J-ChemistryDr K AnanthanarayananPage 1

1

**SRM**  
INSTITUTE OF SCIENCE & TECHNOLOGY  
(Established as Sri Ramakrishna Engineering College, 1983)

---

In this class..

Determination of the strength of a mixture of acetic acid and hydrochloric acid by conductometry

**Expt. No. : 5**

---

10/27/20212

2

## Experiment



### □ Aim :

To estimate the strength of the **mixture of acetic acid and hydrochloric acid present** in a given mixture by conductometry.

### □ Materials required:

Conductivity meter, conductivity cell, standard flask, pipette, burette, funnel, glass rod.

### Chemicals required :

Hydrochloric acid, acetic acid, NaOH solution

10/27/2021

3

3

## Conductometric titration




- **Electrolytic conductivity** of the reaction mixture is continuously monitored as one reactant is added.
- **The equivalence point** is the point at which the **conductivity undergoes a sudden change**. Marked increase or decrease in conductance are associated with the changing concentrations of the two **most highly conducting ions—the hydrogen and hydroxyl ions**.
- The electrical conductivity of an electrolytic solution is **dependent on the number of free ions in the solution** and the charge corresponding to each of these ions.
- The method can be used for **titrating coloured solutions** or **homogeneous suspension** which cannot be used with normal indicators.

4

4

## Apparatus




10/27/2021

5

5



- ❑ Conductivity may be measured by applying an alternating electrical current ( $I$ ) to two electrodes immersed in a solution and measuring the resulting voltage ( $V$ ).
- ❑ During this process, the cations migrate to the negative electrode, the anions to the positive electrode and the solution acts as an electrical conductor.



10/27/2021

6

6



The resistance of the solution (R) can be calculated using Ohm's law ( $V = R \times I$ ).

$$R = V/I$$

where:

V = voltage (volts)

I = current (amperes)

R = resistance of the solution (ohms)

### Conductance

Conductance (G) is defined as the reciprocal of the electrical resistance (R) of a solution between two electrodes.

$$G = 1/R \text{ (S)}$$

10/27/2021

7

7




## Ionic conductance


□ Conductivity (or **specific conductance**) of an electrolytic solution is a **measure of its ability to conduct electricity with the help of free ions in it.**

□ Conductance **of an ion depends on its size and mobility** in an aqueous solution.

□ The order of size of hydrated ionic radii of alkali metal cations is as follows:



□ Hence the ease of ionic conductance is :



10/27/2021

8

8

## Conductometric titration



- When a **mixture of acids** like a strong acid (HCl) and a weak acid (CH<sub>3</sub>COOH) is titrated against a strong base (NaOH), **HCl reacts first followed by CH<sub>3</sub>COOH.**
- When the titration of strong acid and strong base is carried out, there is a **decrease in conductivity** as highly **mobile hydrogen ions (H<sup>+</sup>) are replaced by sodium ions (Na<sup>+</sup>).**



- When the **whole strong acid is consumed**, base reacts with weak acid and **conductivity increases slightly as unionized weak acid becomes the ionized salt.**



10/27/2021

9

9

## Conductometric titration contd...



- After **both the acids are consumed**, there is a **steep increase in conductivity** which gives the end point.
- This **increase in conductivity** is due to the **fast moving hydroxide ions** from the burette solution.
- From this, **amount of base consumed for an acid and in turn, the amount of acids present is calculated**

10/27/2021

10

10


**Conductance of Cations and Anions in Aqueous Solution at 25°C (3)**

Cation	Conductance	Trend	Anion	Conductance
$\text{H}^+$	<b>349.8</b>	high	$\text{OH}^-$	<b>198.3</b>
$\text{K}^+$	73.5	↓	$\text{Br}^-$	78.1
$\text{NH}_4^+$	73.5	↓	$\text{I}^-$	76.8
$\text{Ag}^+$	61.9	↓	$\text{Cl}^-$	76.3
$\text{Na}^+$	50.1	↓	$\text{NO}_3^-$	71.5
$\text{Li}^+$	38.7	↓	$\text{F}^-$	55.4
		low	$\text{CH}_3\text{COO}^-$	40.9

10/27/2021

11


11

**Conductometric titration contd...**


10/27/2021

12

12



- ☐ Plot the graph of conductance on Y axis versus volume of NaOH on X-axis to get three straight lines.
- ☐ The point of intersection of the first and the second lines gives the volume of NaOH needed to get neutralize only HCl acid.
- ☐ The point of intersection of the second and third straight lines gives the volume of NaOH required to neutralize both HCl and CH<sub>3</sub>COOH (after drawing a perpendicular to X-axis).

10/27/2021

13

13

## Procedure

- ☐ The **given mixture of acids is diluted to 100 ml** using distilled water in standard flask.
- ☐ **20 ml of this made up solution** is pipetted out into clean beaker and 100 ml of distilled water is added.
- ☐ The conductivity cell is dipped into the test solution and titrated against NaOH with proper stirring. The **conductance is measured after each 0.5 ml addition of NaOH.**
- ☐ After complete neutralization, the amount of acid present in the given mixture is determined based on the volume **of NaOH consumed.**
- ☐ **Plot a graph between conductance and volume of base added,** where **first end point corresponds to strong acid** and **second end point corresponds to weak acid.**

10/27/2021

14

14

## Tabular column



**Table-1: Titration between mixture of acids and NaOH**

S.No.	Volume of NaOH added (ml)	Conductance ( $\text{ohm}^{-1}$ )
1		
2		
.		
.		
.		
30		

10/27/2021

15

15

## Calculation



### Strength of HCl

$$\begin{aligned}
 \text{Volume of mixture} &= 10 \text{ ml} \quad \mathbf{20 \text{ ml}} \\
 \text{Normality of HCl} &= \text{---? } N_1 \\
 \text{Volume of NaOH} &= V_1 \text{ ml [ 1<sup>st</sup> end point from graph]} \\
 \text{Normality of NaOH} &= 0.1 \text{ N} \\
 \text{Strength of HCl} &= \frac{V_1 \times 0.1}{10} \quad \mathbf{20 \text{ ml}} \\
 &= \text{----- } N.
 \end{aligned}$$

10/27/2021

16

16



## Calculation



### Strength of $\text{CH}_3\text{COOH}$

Volume of mixture	= 10 ml	<b>20 ml</b>
Normality of $\text{CH}_3\text{COOH}$	= --- ? $N_1$	
Volume of NaOH	= $V_2 - V_1$ ml	[ $V_2$ -2 <sup>nd</sup> end point from graph]
Normality of NaOH	= 0.1 N	
Strength of $\text{CH}_3\text{COOH}$	= $\frac{0.1 \times (V_2 - V_1)}{10}$	<b>20 ml</b>
	= -----N	

10/27/2021

17

17

## Result



- ☐ The strength of HCl present in the whole of the solution =
  
- ☐ The strength of  $\text{CH}_3\text{COOH}$  present in the whole of the given solution =

10/27/2021

18

18

Table 1 Standard NaOH versus Unknown mixture of acids



SI.No	Volume of NaOH added (mL)	Conductance (ohm <sup>-1</sup> )
1	0	7.86
2	0.5	7.34
3	1	6.72
4	1.5	6.25
5	2	5.52
6	2.5	5.15
7	3	4.68
8	3.5	4.13
9	4	3.62
10	4.5	3.29
11	5	2.84
12	5.5	2.37
13	6	2.22
14	6.5	2.35
15	7	2.49
16	7.5	2.58

19

17	8	2.64
18	8.5	2.72
19	9	2.88
20	9.5	2.96
21	10	3.14
22	10.5	3.21
23	11	3.64
24	11.5	3.89
25	12	4.13
26	12.5	4.31
27	13	4.69
28	13.5	4.94
29	14	5.23
30	14.5	5.51
31	15	5.87
32	15.5	6.3
33	16	6.72
34	16.5	7.16
35	17	7.59
36	17.5	7.82
37	18	8.24




20

## Problem - 6



Calculate  $\Delta H$  and  $\Delta S$  for the following reaction and decide in which direction each of these factors will drive the reaction.



Compound	$\Delta H_f^\circ(\text{kJ/mol})$	$\Delta S^\circ(\text{J/mol-K})$
$\text{N}_2(\text{g})$		191.61
$\text{H}_2(\text{g})$		130.68
$\text{NH}_3(\text{g})$	-46.11	192.45

Use the values calculated in this problem to predict whether the following reaction is spontaneous at 25 deg C and/or at 500 deg C

27 October 2021

21CYB101J

21

21

## Solution – Problem 6



$$\begin{aligned} \Delta H^\circ &= H_f^\circ(\text{products}) - H_f^\circ(\text{reactants}) \\ &= [2 \text{ mol NH}_3 \times 46.11 \text{ kJ/mol}] - [1 \text{ mol N}_2 \times 0 \text{ kJ/mol} + 3 \text{ mol H}_2 \times 0 \text{ kJ/mol}] \\ &= \mathbf{-92.22 \text{ kJ}} \end{aligned}$$

The reaction is exothermic ( $\Delta H^\circ < 0$ ), which means that the enthalpy of reaction favors the products of the reaction:

$$\begin{aligned} \Delta S^\circ &= S^\circ(\text{products}) - S^\circ(\text{reactants}) \\ &= [2 \text{ mol NH}_3 \times 192.45 \text{ J/mol-K}] - [1 \text{ mol N}_2 \times 191.61 \text{ J/mol-K} + 3 \text{ mol H}_2 \times 130.68 \text{ J/mol-K}] \\ &= \mathbf{-198.75 \text{ J/K}} \end{aligned}$$

The entropy of reaction is unfavorable, however, because there is a significant increase in the order of the system, when  $\text{N}_2$  and  $\text{H}_2$  combine to form  $\text{NH}_3$ .

27 October 2021

21CYB101J

22

22



**This reaction is favored by enthalpy but not by entropy:**

**$\Delta H^\circ = -92.22 \text{ kJ}$  (favorable)**

**$\Delta S^\circ = -198.75 \text{ J/K}$  (unfavorable)**

Before we can compare these terms to see which is larger, we have to incorporate T in our calculation the temperature at which the reaction is run:

$$T_K = 25^\circ \text{C} + 273.15 = 298.15 \text{ K}$$

We then multiply the entropy of reaction by the absolute temperature and subtract the  $T \Delta S^\circ$  term from the  $\Delta H^\circ$  term:

$$\begin{aligned} G^\circ &= \Delta H^\circ - T \Delta S^\circ \\ &= -92,220 \text{ J} - (298.15 \text{ K} \times -198.75 \text{ J/K}) \\ &= -92,220 \text{ J} + 59,260 \text{ J} \\ &= -32,960 \text{ J} \end{aligned}$$

$$G^\circ = -32.96 \text{ kJ}$$

According to this calculation, the reaction should be spontaneous at 25 deg C.

27 October 2021

21CYB101J

23

23



Before we can decide whether the reaction is still spontaneous we need to calculate the temperature of the kelvin scale:

$$T_K = 500^\circ \text{C} + 273 = 773 \text{ K}$$

We then multiply the entropy term by this temperature and subtract this quantity from the enthalpy term:

$$\begin{aligned} G^\circ_{773} &= \Delta H^\circ_{298} - T \Delta S^\circ_{298} \\ &= 92,220 \text{ J} - (773 \text{ K} \times -198.75 \text{ J/K}) \\ &= 92,220 \text{ J} - (-153,600 \text{ J}) \\ &= 61,380 \text{ J} \end{aligned}$$

$$G^\circ = 61.4 \text{ kJ}$$

**Because the entropy term becomes larger as the temperature increases, the reaction changes from one which is favorable at low temperatures to one that is unfavorable at high temperatures.**

27 October 2021

21CYB101J

24


24

## Problem – 7



$K_{sp}$  for  $MgCO_3$  at  $25^\circ C$  is  $2.0 \times 10^{-8}$ . What are the ion concentrations in a saturated solution at this temperature?

**Solution** As always, begin with a balanced equation:



Write the  $K_{sp}$  expression:

$$K_{sp} = [Mg^{2+}][CO_3^{2-}]$$

For this example, we are given the value for  $K_{sp}$  and need to find the ion concentrations.

We will let our unknown ion concentrations equal  $x$ .

**The balanced equation tells us that both  $Mg^{2+}$  and  $CO_3^{2-}$  will have the same concentration!**

27 October 2021

21CYB101J

25

25

## Solution – Problem 7



$$K_{sp}$$

$$= [Mg^{2+}][CO_3^{2-}]$$

$$x^2 = 2.0 \times 10^{-8}$$

$$x = \sqrt{(2.0 \times 10^{-8})}$$

find the square root of  $2.0 \times 10^{-8}$

$$= 1.4 \times 10^{-4} M$$

$$x = [Mg^{2+}]$$

$$= 1.4 \times 10^{-4} M$$

**Answer**

$$\text{AND } x = [CO_3^{2-}]$$

$$= 1.4 \times 10^{-4} M$$

27 October 2021

21CYB101J

26

26

## Problem – 8



The concentration of lead ions in a saturated solution of  $\text{PbI}_2$  at  $25^\circ\text{C}$  is  $1.3 \times 10^{-3}\text{M}$ . What is its  $K_{\text{sp}}$ ?

27 October 2021

21CYB101J


27

27

## Solution – Problem 8



Begin problems involving  $K_{\text{sp}}$  by writing a balanced equation:



Write the  $K_{\text{sp}}$  expression (be careful!):

$$K_{\text{sp}} = [\text{Pb}^{2+}][\text{I}^{-}]^2$$

Determine the concentration of the ions. Take care to determine the concentration of  $\text{I}^{-}$  ions:

$$[\text{PbI}_2] = 1.30 \times 10^{-3}\text{M}$$

$$[\text{Pb}^{2+}] = 1.30 \times 10^{-3}\text{M}$$

$$[\text{I}^{-}] = 2 \times 1.30 \times 10^{-3} = 2.60 \times 10^{-3}$$

Substitute values into the  $K_{\text{sp}}$  expression and solve for the unknown:

$$\begin{aligned} K_{\text{sp}} &= [\text{Pb}^{2+}][\text{I}^{-}]^2 \\ &= (1.30 \times 10^{-3})(2.60 \times 10^{-3})^2 \\ &= (1.30 \times 10^{-3})(6.76 \times 10^{-6}) \end{aligned}$$

**Watch Exponents!!!**

$$K_{\text{sp}} = 8.79 \times 10^{-9}\text{M}$$

**Answer**

27 October 2021

21CYB101J

28

28

# Thank you all for your attention

Information presented here were collected from various sources – textbooks, articles, manuscripts, internet and newsletters. All the researchers and authors of the above mentioned sources are greatly acknowledged.