



# CHEMISTRY

43<sup>rd</sup> class, Date : 10-11-2021

Dr. K. Ananthanarayanan  
Associate Professor (Research)  
Room number 319, Raman Research Park  
Department of Chemistry

Email : ananthak@srmist.edu.in  
Mobile : 9840154665

## Last class....



- ☐ Isomers in organic compounds
  
- ☐ 3-D representation

## In this class....



- ☐ Conformational analysis, continuation
  
- ☐ Stereochemistry

## Organic chemistry



- ☐ Isomerism : Structural and stereo isomerism
  
- ☐ Conformational analysis
  
- ☐ Absolute configuration : CIP rules (naming enantiomers)
  
- ☐ Reactions : Substitution, Elimination, Oxidation, Reduction, Addition, Cyclisation and C-C bond formation reactions
  
- ☐ Synthesis of pharmaceutical products, few examples

## Review of Isomerism




- ❑ **Isomers** – Compounds that have the same molecular formula but do not have identical structures. Generally classified into two types:
- ❑ **Constitutional Isomers** – differ in the way their atoms are connected
- ❑ **Stereoisomers** – differ in the way their atoms are arranged in space

21CYB101J-Chemistry


5

## Review of Isomerism

E.g.,  $C_4H_8$ 


Stereo

Constitutional



21CYB101J-Chemistry

6




## Review of Isomerism




**Structural isomerism**, or **constitutional isomerism** is a form of isomerism in which molecules with the same molecular formula have different bonding patterns and atomic organization, as opposed to stereoisomerism, in which molecular bonds are always in the same order and only spatial arrangement differs.

Two isomers of butane,  $C_4H_{10}$



n-butane



Iso-butane

### Types of Structural Isomerism

- 1) Chain Isomers
- 2) Position Isomers
- 3) Functional Isomers
- 4) Metamerism
- 5) Tautomerism

## Structural Isomers




### CHAIN ISOMERISM

In **chain isomerism**, or **skeletal isomerism**, components of the (usually carbon) **skeleton are distinctly re-ordered** to create different structures.



n-pentane  
(often called simply "pentane")



isopentane  
(2-methylbutane)





neopentane  
(dimethylpropane)

### POSITION ISOMERISM

Isomerism caused by the difference in the position of functional group in the same chain is termed as **position isomerism (regioisomerism)**

*o*-dichlorobenzene



*m*-dichlorobenzene

21CYB101J-Chemistry

9

## Hexane, C<sub>6</sub>H<sub>14</sub>



### Formulae

### IUPAC names

1. CH<sub>3</sub> - CH<sub>2</sub> - CH<sub>2</sub> - CH<sub>2</sub> - CH<sub>2</sub> - CH<sub>3</sub>    Hexane
2. CH<sub>3</sub> - CH<sub>2</sub> - CH<sub>2</sub> -  $\begin{array}{c} \text{CH} \\ | \\ \text{CH}_3 \end{array}$  - CH<sub>3</sub>    2-methyl pentane
3. CH<sub>3</sub> - CH<sub>2</sub> -  $\begin{array}{c} \text{CH} \\ | \\ \text{CH}_3 \end{array}$  - CH<sub>2</sub> - CH<sub>3</sub>    3-Methylpentane
4. CH<sub>3</sub> -  $\begin{array}{c} \text{CH} \\ | \\ \text{CH}_3 \end{array}$  -  $\begin{array}{c} \text{CH} \\ | \\ \text{CH}_3 \end{array}$  - CH<sub>3</sub>    2, 3-Dimethyl-butane
5. CH<sub>3</sub> -  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{C} \\ | \\ \text{CH}_3 \end{array}$  - CH<sub>2</sub> - CH<sub>3</sub>    2, 2-Dimethylbutane

21CYB101J-Chemistry

10

## Structural Isomers




### FUNCTIONAL ISOMERISM

Functional isomers have same molecular formula but different functional group




Ethyl Alcohol




Dimethyl Ether

### METAMERISM

This type of isomerism is due to the unequal distribution of carbon atoms on either side of functional group. Metamerism belongs to same homologous series.



Diethyl Ether



Methyl Propyl Ether

21CYB101J-Chemistry

11

## Structural Isomers



### TAUTOMERISM

Tautomers are constitutional isomers that differ from each other in the locations of a hydrogen atom and one or more double bonds and interconvert rapidly. The carbon skeleton of the compound is unchanged. A reaction which involves simple proton transfer in an intramolecular fashion is called a tautomerism

Keto-enol tautomerism is a very common process, and is acid or base catalysed. Typically the 'keto' form of the compound is more stable, but in some instances the 'enol' form can be the more stable



An equilibrium exists between the two forms

21CYB101J-Chemistry

12

## Stereoisomers



- ❑ Stereochemical isomers are molecules, which have the **same chemical formula and bond connectivity but different relative arrangement in three-dimensional space**
- ❑ Stereochemistry is the study of the **relative arrangement of atoms or groups in a molecule in three dimensional space**
- ❑ In contrast, constitutional isomers have same molecular formula but different bond connectivity





21CYB101J-Chemistry

13

## Cis-trans isomers



- ❑ Differ in the arrangement of their atoms in space



Copyright © 2006 Pearson Education, Inc.





21CYB101J-Chemistry

14

## Cis-trans isomers



## Cyclic structure






*cis*-1-bromo-3-chlorocyclobutane*trans*-1-bromo-3-chlorocyclobutane*cis*-1,4-dimethylcyclohexane*trans*-1,4-dimethylcyclohexane

Copyright © 2006 Pearson Education, Inc.

21CYB101J-Chemistry

15

## 3-D representation




21CYB101J-Chemistry


16




## Representation of 3-D-ETHANE




3D Image

Wedge – Dash or  
Flying - Wedge formula

Sawhorse projection



Newman Projection





Fischer Projection


21CYB101J-Chemistry

17


## Wedge – Dash projection



- The Flying-Wedge projection is the most common three-dimensional representation of a three dimensional molecule on a two dimensional surface (paper)
- This kind of representation is usually done for molecules containing chiral centre. In this representation, the ***ordinary lines represent bonds in the plane of the paper***. A solid Wedge (  ) represents a **bond above the plane of the paper** and a dashed wedge (  ) or a broken line represents a **bond below the plane of the paper**.




Methane



21CYB101J-Chemistry

18

## Wedge – Dash projection




21CYB101J-Chemistry

19




The Flying-Wedge projection formula of Lactic acid


$\text{CH}_3\text{CH}(\text{OH})\text{COOH}$  for example, can be shown as follows.....



21CYB101J-Chemistry

20




  
 SRM
   
INSTITUTE OF SCIENCE & TECHNOLOGY
  
(Chartered as the University with effect from 1st July 2015)

## Sawhorse projection


- ❑ In this representation, the molecule is **viewed slightly from above** and from the right and then projected on the paper
- ❑ The bond between the two carbon atoms **is drawn diagonally** and of a relatively greater length for the sake of clarity
- ❑ The **lower left hand carbon is taken as the front carbon** and the **upper right hand carbon as the back carbon**
- ❑ All **parallel bonds in sawhorse formula are eclipsed** and **all anti parallel bonds are opposite** or **trans/anti** to each other
- ❑ The sawhorse presentation of **eclipsed and staggered conformations** of ethane was shown in the previous slide

21CYB101J-Chemistry 22

## Conformations of acyclic alkanes




**Conformations** are different arrangements of atoms that are interconverted by rotation about single bonds



21CYB101J-Chemistry

23




- ☐ Names are given to two different conformations
- ☐ In the **eclipsed conformation**, the C—H bonds on one carbon are **directly aligned** with the C—H bonds on the adjacent carbon
- ☐ In the **staggered conformation**, the C—H bonds on one carbon **bisect** the H—C—H bond angle on the adjacent carbon

21CYB101J-Chemistry

24

- ❑ Rotating the atoms on one carbon by  $60^\circ$  converts an eclipsed conformation into a staggered conformation, and vice versa.
- ❑ The angle that separates a bond on one atom from a bond on an adjacent atom is called a **dihedral angle**. For ethane in the staggered conformation, the **dihedral angle for the C—H bonds is  $60^\circ$ . For eclipsed ethane, it is  $0^\circ$ .**




21CYB101J-Chemistry

25


- ❑ The staggered and eclipsed conformations of ethane **interconvert** at room temperature, **but each conformer is not equally stable**
- ❑ The **staggered conformations are more stable** (lower in energy) than the eclipsed conformations
- ❑ **Electron-electron repulsion between bonds** in the eclipsed conformation increases its energy compared with the staggered conformation, where the bonding electrons are farther apart

These C—H bonds are farther apart.



more stable

These C—H bonds are closer together.




less stable

21CYB101J-Chemistry

26

## Newman projection



21CYB101J-Chemistry


27



- End-on representations for conformations are commonly drawn using a convention called a **Newman projection**.

### How to Draw a Newman Projection:

**Step 1.** Look directly down the C—C bond (end-on), and draw a circle with a dot in the center to represent the carbons of the C—C bond.




21CYB101J-Chemistry

28

**Step 2.** Draw in the bonds.

Draw the bonds on the front C as **three lines meeting at the center of the circle.**


Draw the bonds on the back C as **three lines coming out of the edge of the circle.**



**Step 3.** Add the atoms on each bond.




=



Each C has 3 H's in ethane.


## Newman projection



- In the **staggered form**, all the bonds attached to the C1 and C2 carbon atoms are **maximum apart from each other** whereas in the **eclipsed form**, the bonds attached to the two carbon atoms are **closest to each other**.

21CYB101J-Chemistry

31




- Energetically, out of the two forms, the **staggered form is the more favored form** - bonds are maximally apart from each other – hence has the **least amount of steric strain**
- On the other hand, the bonds are closest to each other in the eclipsed form and it **has the highest energy**
- The strain existing in the eclipsed form is termed as **torsional strain (resistance to bond twisting)**. The energy barrier between the two forms in the ethane molecule has been experimentally found to be about 3 kcal/mol

21CYB101J-Chemistry


32



- ❑ Torsional strain or eclipsing strain is the increase in potential energy of a molecule **due to repulsion between electrons in bonds that do not share an atom.**



1





2

- ❑ **The smallest dihedral angle is 60° in 1; it is 0° in 2.** Thus, two C-H bonds not sharing a carbon atom that are closest to each other **(ex: C-H1 and C-H4) are closer in 2 than in 1.**
- ❑ The repulsion between electrons in bonds that do not share a carbon atom is, therefore, greater in 2 than in 1. **Consequently, torsional strain is greater in 2 than in 1.**

21CYB101J-Chemistry

33


## Newman projection




- ❑ In these Formulae the **molecule is viewed from the front.** The carbon atom **nearer to the eye is represented by a point** and the three atoms or groups are shown attached to it by **three lines at an angle of 120°** to each other.
- ❑ A **Newman Projection** can be drawn such that the groups on the front carbon are *staggered* (60 degrees apart) or *eclipsed* (directly overlapping) with the groups on the back carbon. Below are two **Newman Projections** of ethane, C<sub>2</sub>H<sub>6</sub>. The structure on the right is staggered, and the structure on the left is eclipsed.

21CYB101J-Chemistry


34



## Newman projection, n-butane



**Eclipsed**




**Highest-energy conformation**  
of butane

Front groups **eclipsed** with respect  
to back groups

CH<sub>3</sub> groups are **syn** (dihedral angle 0°)

Strain energy about **5.0 kcal/mol** above  
that of the staggered conformation  
(at right)

**Staggered**





**Lowest energy conformation**  
of butane

Front groups **staggered** with  
respect to back groups

CH<sub>3</sub> groups are **anti**  
(dihedral angle 180°)

Energy minimum





  
 SRM
   
INSTITUTE OF SCIENCE & TECHNOLOGY
  
(Chartered as the University with effect from 1st July, 2015)

### Definitions

- ☐ **Anti** - Description given to two substituents attached to adjacent atoms when their bonds are at  $180^\circ$  with respect to each other
- ☐ **Syn** - Description given to two substituents attached to adjacent atoms when their bonds are at  $0^\circ$  with respect to each other
- ☐ **Gauche** - Description given to two substituents attached to adjacent atoms when their bonds are at  $60^\circ$  with respect to each other

## Two Different Eclipsed Conformations




## Strain energy can be quantified

**TABLE 4.1** Energy Costs for Interactions in Alkane Conformers


Interaction	Cause	Energy cost	
		(kJ/mol)	(kcal/mol)
H ↔ H eclipsed	Torsional strain	4.0	1.0
H ↔ CH <sub>3</sub> eclipsed	Mostly torsional strain	6.0	1.4
CH <sub>3</sub> ↔ CH <sub>3</sub> eclipsed	Torsional plus steric strain	11.0	2.6
CH <sub>3</sub> ↔ CH <sub>3</sub> gauche	Steric strain	3.8	0.9

©2004 Thomson - Brooks/Cole

## Steric and Torsional Strain both when Eclipsed






## PE Diagram for Butane



- ❑ There are **two energy minima**, the *gauche* and *anti* forms, which are both **staggered** and thus have no **torsional strain**
- ❑ The *anti* form is the **absolute energy minimum**, since the *gauche* form has a small steric interaction between the two methyl groups. At a dihedral angle of 60 degrees, one hydrogen of each of the methyl groups is relatively close to a hydrogen of the other methyl group (van der Waals repulsion)
- ❑ Carefully note the difference between **steric strain** and **torsional strain**. The latter arises from eclipsed bonds, while the former arises from atoms which are too close to each other

43

## Gauche Interaction in Butane




© 2004 Thomson/Brooks Cole

- ❑ There are **also two energy maxima, both of which are eclipsed and thus torsionally strained**. The higher energy conformation also has steric strain
  
- ❑ **Torsional strain arises from the repulsion of electrons in bonds; steric strain arises from atoms that are too close!**

45

## PE Diagram for Butane



# Thank you all for your attention

Information presented here were collected from various sources –  
textbooks, articles, manuscripts, internet and newsletters. All the  
researchers and authors of the above mentioned sources are greatly  
acknowledged.