

3.2.1 Laplace transform of Derivatives:	159
3.3 Initial value and Final value theorems	162
3.4 Convolution Theorem	163
3.5 Laplace transform of periodic functions	171
3.6 Inverse Laplace Transform	176
3.7 Convolution Theorem	180
3.8 Application of Laplace Transform to Solve Differential Equations	194
4 ANALYTIC FUNCTIONS	211
4.1 Analytic Function or Regular Function or Holomorphic Function	212
4.1.1 Properties of analytic function:	230
4.2 Milne - Thomson Method	241
4.3 Conformal Mapping	254
4.3.1 Magnification and Rotation	255
4.3.2 Inversion and Reflection	255
4.4 Bilinear Transformation	265
5 COMPLEX INTEGRATION	274
5.1 Evaluation of line integral	276
5.2 Cauchy's Integral Theorem	281
5.2.1 Cauchy's Theorem for Multiply Connected Region	282
5.3 Taylor's Series and Laurent's Series	294
5.4 Poles and Residues	309
5.5 Cauchy's Residue Theorem	313

1 MULTIPLE INTEGRALS

INTRODUCTION AND NOTATION

1.1 Double and Triple Integrals

The symbol $\int_{y_1}^{y_2} \int_{x_1}^{x_2} f(x, y) dx dy$ denotes a double integral. It means that $f(x, y)$ is to be integrated first with respect to x , considering y temporarily, as a constant, between the limits x_1 and x_2 and the resulting expression is then to be integrated with respect to y between the limits y_1 and y_2 . Here x_1 and x_2 may be constants or functions of y alone.

(i.e.) $x_1 = f_1(y), x_2 = f_2(y), y_1$ and y_2 are constants.

Similarly the symbol $\int_{z_1}^{z_2} \int_{y_1}^{y_2} \int_{x_1}^{x_2} f(x, y, z) dx dy dz$ denotes a triple integral. This is concluded as follows. First $f(x, y, z)$ is integrated with respect to x between the limits x_1 and x_2 , keeping y and z as constants temporarily. The resulting function is integrated with respect to y between the limits y_1 and y_2 , keeping z constant. The result thus obtained is finally integrated with respect to z between the limits z_1 and z_2 . In this integral, z_1 and z_2 are constants, y_1 and y_2 are either constants or functions of z alone and x_1, x_2 are either constants or functions of y and z .

Note: In $\int \int f(x, y) dx dy$, if the right hand integral has limits involving x , then the first integral is with respect to y .

Differentiation Formulas:

$$(i) \frac{d}{dx}(k) = 0$$

$$(ii) \frac{d}{dx}[f(x) \pm g(x)] = f'(x) \pm g'(x)$$

$$(iii) \frac{d}{dx}[k \cdot f(x)] = k \cdot f'(x)$$

$$(iv) \frac{d}{dx}[f(x) \cdot g(x)] = f(x)g'(x) + g(x)f'(x)$$

$$(v) \frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$$

$$(vi) \frac{d}{dx}f(g(x)) = f'(g(x)) \cdot g'(x)$$

$$(vii) \frac{d}{dx}(x^n) = nx^{n-1}$$

$$(viii) \frac{d}{dx}(e^x) = e^x$$

$$(ix) \frac{d}{dx}(a^x) = a^x \log a, a > 0$$

$$(x) \frac{d}{dx}(\log_e x) = \frac{1}{x}, x > 0$$

$$(xi) \frac{d}{dx}(\sin x) = \cos x$$

$$(xii) \frac{d}{dx}(\cos x) = -\sin x$$

$$(xiii) \frac{d}{dx}(\tan x) = \sec^2 x$$

$$(xiv) \frac{d}{dx}(\cot x) = -\csc^2 x$$

$$(xv) \frac{d}{dx}(\sec x) = \sec x \tan x$$

$$(xvi) \frac{d}{dx}(\csc x) = -\csc x \cot x$$

$$(xvii) \frac{d}{dx}(\sinh x) = \cosh x$$

$$(xviii) \frac{d}{dx}(\cosh x) = \sinh x$$

$$(xix) \frac{d}{dx}(\tanh x) = \sec h^2 x$$

$$(xx) \frac{d}{dx}(\coth x) = -\csc h^2 x$$

$$(xxi) \frac{d}{dx}(\sec hx) = -\sec hx \tanh x$$

$$(xxii) \frac{d}{dx}(\csc hx) = -\csc hx \coth x$$

$$(xxiii) \frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}} = -\frac{d}{dx}(\cos^{-1} x)$$

$$(xxiv) \frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2} = -\frac{d}{dx}(\cot^{-1} x)$$

$$(xxv) \frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{x^2-1}} = -\frac{d}{dx}(\csc^{-1} x)$$

$$(xxvi) \frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1+x^2}}$$

$$(xxvii) \frac{d}{dx}(\cosh^{-1} x) = \frac{1}{\sqrt{x^2-1}}$$

$$(xxviii) \frac{d}{dx}(\tan^{-1} x) = \frac{1}{1-x^2}, |x| < 1$$

$$(xxix) \frac{d}{dx}(\coth^{-1} x) = -\frac{1}{x^2-1}, |x| > 1$$

$$(xxx) \frac{d}{dx}(\sec h^{-1} x) = \frac{-1}{x\sqrt{1-x^2}}$$

$$(xxxi) \frac{d}{dx}(\csc h^{-1} x) = \frac{1}{x\sqrt{x^2+1}}$$

Integration Formulas:

$$1. \int dx = x + C$$

$$2. \int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$$

$$3. \int \frac{1}{x} dx = \log x + C$$

$$4. \int e^{ax} dx = \frac{e^{ax}}{a} + C$$

$$5. \int \sin x dx = -\cos x + C$$

$$6. \int \cos x dx = \sin x + C$$

$$7. \int \sec^2 x dx = \tan x + C$$

$$8. \int \csc^2 x dx = -\cot x + C$$

$$9. \int \sec x \tan x dx = \sec x + C$$

$$10. \int \csc x \cot x dx = -\csc x + C$$

$$11. \int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x + C$$

$$12. \int \frac{dx}{1+x^2} = \tan^{-1} x + C$$

$$13. \int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x + C$$

$$14. \int \sinh x dx = \cosh x + C$$

$$15. \int \cosh x dx = \sinh x + C$$

$$16. \int \sec h^2 x dx = \tanh x + C$$

1 MULTIPLE INTEGRALS

17. $\int \csc h^2 x dx = -\coth x dx + C$

18. $\int \sec hx \tanh x dx = -\sec hx + C$

19. $\int \csc hx \coth x dx = -\csc hx + C$

20. $\int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \frac{x-a}{x+a} + C$

21. $\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \log \frac{a+x}{a-x} + C$

22. $\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$

23. $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C$

24. $\int \frac{dx}{\sqrt{x^2 + a^2}} = \sinh^{-1} \frac{x}{a} + C \text{ (or) } \log(x + \sqrt{x^2 + a^2})$

25. $\int \frac{dx}{\sqrt{x^2 - a^2}} = \cosh^{-1} \frac{x}{a} + C \text{ (or) } \log(x + \sqrt{x^2 - a^2})$

26. $\int \tan x dx = \log \sec x = -\log \cos x + C$

27. $\int \cot x dx = \log \sin x + C$

28. $\int \sec x dx = \log(\sec x + \tan x) + C$

29. $\int \csc x dx = \log(\csc x - \cot x) + C$

1 MULTIPLE INTEGRALS

30. $\int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) + C$

31. $\int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) + C$

32. $\int \sqrt{a^2 - x^2} dx = \frac{a^2}{2} \sin^{-1} \frac{x}{a} + \frac{x}{2} \sqrt{a^2 - x^2} + C$

33. $\int \sqrt{a^2 + x^2} dx = \frac{a^2}{2} \sinh^{-1} \frac{x}{a} + \frac{x}{2} \sqrt{a^2 + x^2} + C$

34. $\int \sqrt{x^2 - a^2} dx = -\frac{a^2}{2} \cosh^{-1} \frac{x}{a} + \frac{x}{2} \sqrt{x^2 - a^2} + C$

35. $\int a^x dx = \frac{a^x}{\log a} + C$

Example 1.1. Evaluate $\int_0^2 \int_0^1 4xy dx dy$

Solution: Let $I = \int_0^2 \int_0^1 4xy dx dy$

$$= 4 \int_0^2 \left(y \frac{x^2}{2} \right)_0^1 dy = 2 \left(\frac{y^2}{2} \right)_0^2 = 4$$

Example 1.2. Evaluate $\int_0^a \int_0^b \int_0^c (x + y + z) dx dy dz$

Solution: Let $I = \int_0^a \int_0^b \left(\frac{x^2}{2} + yx + zx \right)_{x=0}^{x=c} dy dz$

$$= \int_0^a \int_0^b \left(\frac{c^2}{2} + yc + zc \right) dy dz$$

$$\begin{aligned}
&= \int_0^a \left(\frac{c^2 y}{2} + \frac{y^2 c}{2} + zcy \right)_{y=0}^{y=b} dz \\
&= \int_0^a \left(\frac{c^2 b}{2} + \frac{b^2 c}{2} + zcb \right) dz \\
&= \left(\frac{c^2 bz}{2} + \frac{b^2 cz}{2} + \frac{z^2 cb}{2} \right)_{z=0}^{z=a} = \frac{abc}{2}(a+b+c)
\end{aligned}$$

Example 1.3. Evaluate $\int_0^{2\pi} \int_0^\pi \int_0^a r^4 \sin \phi dr d\phi d\theta$

Solution: The given integral

$$\begin{aligned}
&\int_0^{2\pi} d\theta \int_0^\pi d\phi \int_0^a r^4 \sin \phi dr = \int_0^{2\pi} d\theta \int_0^\pi \left(\frac{r^5}{5} \right)_0^a \sin \phi d\phi \\
&= \frac{a^5}{5} \int_0^{2\pi} d\theta \int_0^\pi \sin \phi d\phi = \frac{a^5}{5} \int_0^{2\pi} (-\cos \phi)_0^\pi d\theta \\
&= \frac{2}{5} a^5 \int_0^{2\pi} d\theta \text{ as } \cos \pi = -1, \cos 0 = 1 \\
&= \frac{4}{5} \pi a^5
\end{aligned}$$

Example 1.4. Evaluate $\int_0^1 \int_0^x dx dy$

Solution: Let I be the given integral.

$\therefore I = \int_0^1 (y)_0^x dx$ as right hand integral has a limit which is function of x.

$$I = \int_0^1 x dx = \left(\frac{x^2}{2} \right)_0^1 = \frac{1}{2}$$

Example 1.5. Evaluate $\int_0^\pi \int_0^{a \sin \theta} r d\theta dr$

$$\begin{aligned}
\text{Solution: Let } I &= \int_0^\pi \left(\frac{r^2}{2} \right)_0^{a \sin \theta} = \frac{1}{2} \int_0^\pi a^2 \sin^2 \theta d\theta \\
&= \frac{a^2}{2} \int_0^\pi \left(\frac{1 - \cos 2\theta}{2} \right) d\theta \\
&= \frac{a^2}{2} \cdot \frac{1}{2} \left(\theta - \frac{\sin 2\theta}{2} \right)_0^\pi \\
&= \frac{\pi a^2}{4} \text{ as } \sin \pi = 0, \sin 0 = 0
\end{aligned}$$

Example 1.6. Evaluate $\int_0^1 \int_0^{\sqrt{1+y^2}} \frac{dxdy}{1+x^2+y^2}$

$$\begin{aligned}
\text{Solution: The given integral} &= \int_0^1 \left[\int_0^{\sqrt{1+y^2}} \frac{1}{(1+y^2)+x^2} dx \right] dy \\
&= \int_0^1 \left[\frac{1}{\sqrt{1+y^2}} \tan^{-1} \left(\frac{x}{\sqrt{1+y^2}} \right) \right]_{x=0}^{x=\sqrt{1+y^2}} dy \\
&= \int_0^1 \left[\frac{1}{\sqrt{1+y^2}} (\tan^{-1}(1) - \tan^{-1} 0) \right] dy \\
&= \frac{\pi}{4} \int_0^1 \frac{dy}{\sqrt{1+y^2}}
\end{aligned}$$

1 MULTIPLE INTEGRALS

$$= \frac{\pi}{4} \left[\log(y + \sqrt{1+y^2}) \right]_0^1 = \frac{\pi}{4} \log(1 + \sqrt{2})$$

Example 1.7. Evaluate $\int_0^a \int_0^{\sqrt{a^2-y^2}} \sqrt{a^2-x^2-y^2} dx dy$

$$\begin{aligned} \text{Solution: Let } I &= \int_0^a \int_0^{\sqrt{a^2-y^2}} \sqrt{a^2-x^2-y^2} dx dy \\ &= \int_0^a \left[\frac{x}{2} \sqrt{(a^2-y^2)-x^2} + \left(\frac{a^2-y^2}{2} \right) \sin^{-1} \left(\frac{x}{\sqrt{a^2-y^2}} \right) \right]_0^{\sqrt{a^2-y^2}} dy \\ &= \int_0^a \left(\frac{a^2-y^2}{2} \right) \frac{\pi}{2} y = \frac{\pi}{4} \left[a^2 y - \frac{y^3}{3} \right]_0^a = \frac{\pi a^3}{6} \end{aligned}$$

Example 1.8. Evaluate $\int_0^1 \int_{y^2}^1 \int_0^{1-x} x dz dx dy$

$$\begin{aligned} \text{Solution: Let } I &= \int_0^1 \int_{y^2}^1 \int_0^{1-x} x dz dx dy \\ &= \int_0^1 \int_{y^2}^1 x(z)_0^{1-x} dx dy = \int_0^1 \int_{y^2}^1 (x-x^2) dx dy \\ &= \int_0^1 \left[\left(\frac{1}{2} - \frac{1}{3} \right) - \left(\frac{y^4}{2} - \frac{y^6}{3} \right) \right] dy \\ &= \int_0^1 \left(\frac{1}{6} - \frac{y^4}{2} + \frac{y^6}{3} \right) dy \\ &= \left(\frac{1}{6}y - \frac{1}{2}\frac{y^5}{5} + \frac{1}{3}\frac{y^7}{7} \right)_0^1 \\ &= \frac{1}{6} - \frac{1}{10} + \frac{1}{21} = \frac{4}{35} \end{aligned}$$

1 MULTIPLE INTEGRALS

Example 1.9. Evaluate $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} \frac{dxdydz}{\sqrt{1-x^2-y^2-z^2}}$

Solution:

$$\begin{aligned} \text{Let } I &= \int_0^1 \int_0^{\sqrt{1-x^2}} \left[\sin^{-1} \left(\frac{z}{\sqrt{(1-x^2-y^2)-z^2}} \right) \right]_0^{\sqrt{1-x^2-y^2}} dx dy \\ &= \int_0^1 \int_0^{\sqrt{1-x^2}} \frac{\pi}{2} dx dy = \frac{\pi}{2} \int_0^1 (y)_0^{\sqrt{1-x^2}} dx = \frac{\pi}{2} \int_0^1 \sqrt{1-x^2} dx \\ &= \frac{\pi}{2} \left[\frac{x}{2} \sqrt{1-x^2} + \frac{1}{2} \sin^{-1} x \right]_0^1 = \frac{\pi^2}{8} \end{aligned}$$

Example 1.10. Evaluate $\int_0^{\frac{\pi}{2}} \int_0^{a \cos \theta} r \sqrt{a^2 - r^2} d\theta dr$

$$\begin{aligned} \text{Solution: Let } I &= \int_0^{\frac{\pi}{2}} \left(-\frac{1}{3} \right) [(a^2 - r^2)^{3/2}]_0^{a \cos \theta} d\theta \\ &= -\frac{1}{3} \int_0^{\frac{\pi}{2}} [(a^2 - a^2 \cos^2 \theta)^{3/2} - (a^2)^{3/2}]_0^{a \cos \theta} d\theta \\ &= -\frac{1}{3} \int_0^{\frac{\pi}{2}} (a^3 \sin^3 \theta - a^3) d\theta \\ &= \frac{a^3}{3} \int_0^{\frac{\pi}{2}} (1 - \sin^3 \theta) d\theta \\ &= \frac{a^3}{3} \left[\theta + \frac{1}{4} \left(-\frac{1}{3} \cos 3\theta + 3 \cos \theta \right) \right]_0^{\frac{\pi}{2}} \end{aligned}$$

$$= \frac{a^3}{3} \left(\frac{\pi}{2} - \frac{2}{3} \right)$$

Example 1.11. Evaluate $\int_0^1 \int_0^y xye^{-x^2} dx dy$

Solution: Let $I = \int_0^1 y \left(\frac{e^{-x^2}}{2} \right)_0^y dy$

$$= -\frac{1}{2} \int_0^1 (ye^{-y^2} - y) dy$$

$$= -\frac{1}{2} \left[\frac{e^{-y^2}}{-2} - \frac{y^2}{2} \right]_0^1$$

$$= \frac{1}{4} [(e^{-1} + 1) - (1)] = \frac{1}{4e}$$

Example 1.12. Evaluate $\int_0^a \int_0^{a-x} \int_0^{a-x-y} x^2 dx dy dz$

Solution: Let $I = \int_0^a \int_0^{a-x} [x^2 z]_0^{a-x-y} dy dx$

$$= \int_0^a \int_0^{a-x} x^2 (a - x - y) dy dx$$

$$= \int_0^a \left[x^2 (a - x)y - x^2 \cdot \frac{y^2}{2} \right]_0^{a-x} dx$$

$$= \frac{1}{2} \int_0^a x^2 (a - x)^2 dx$$

$$= \frac{1}{2} [a^2 \frac{x^3}{3} - 2a \frac{x^4}{4} + \frac{x^5}{5}]_0^a = \frac{a^5}{60}$$

Example 1.13. Evaluate $\int_0^a \int_0^x \int_0^y xyz dz dy dx$

Solution: Let $I = \int_0^a \int_0^x \left[\frac{xyz^2}{2} \right]_0^y dy dx$

$$= \frac{1}{2} \int_0^a \int_0^x xy^3 dy dx = \frac{1}{2} \int_0^a \left[\frac{xy^4}{4} \right]_0^x dx$$

$$= \frac{1}{8} \left[\frac{x^6}{6} \right]_0^a = \frac{a^6}{48}$$

Example 1.14. Evaluate $\int_0^{1/2} \int_0^1 \frac{x}{\sqrt{1 - x^2 y^2}} dy dx$

Solution: Let $I = \int_0^{1/2} x \left[\frac{\sin^{-1} xy}{x} \right]_0^1$

$$= \int_0^{1/2} \sin^{-1} x dx$$

$$= \left[x \sin^{-1} x + \sqrt{1 - x^2} \right]_0^{1/2} = \frac{\pi}{12} + \frac{\sqrt{3}}{2} - 1$$

Example 1.15. Evaluate $\int_0^1 \int_0^{1-z} \int_0^{1-y-z} xyz dx dy dz$

Solution: Let $I = \int_0^1 \int_0^{1-z} yz \left(\frac{x^2}{2} \right)_0^{1-y-z} dy dz$

$$= \frac{1}{2} \int_0^1 \int_0^{1-z} yz (1 - y - z)^2 dy dz$$

$$= \frac{1}{2} \int_0^1 yz [(1 - z)^2 - 2(1 - z)y + y^2] dy dz$$

$$\begin{aligned}
&= \frac{1}{2} \int_0^1 \left[z(1-z)^2 \frac{y^2}{2} - 2z(1-z) \frac{y^3}{3} + z \frac{y^4}{4} \right]_0^{1-z} dz \\
&= \frac{1}{2} \int_0^1 \left(\frac{1}{2} z(1-z)^4 - \frac{2}{3} z(1-z)^4 + \frac{1}{4} z(1-z)^4 \right) dz \\
&= \frac{1}{2} \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) \int_0^1 z(1-z)^4 dz \\
&= \frac{1}{24} \int_0^1 [1 - (1-z)](1-z)^4 dz \\
&= \frac{1}{24} \left[\frac{(1-z)^5}{-5} + \frac{(1-z)^6}{6} \right] \\
&= \frac{1}{24} \left(\frac{1}{5} - \frac{1}{6} \right) = \frac{1}{720}
\end{aligned}$$

Example 1.16. Evaluate $\int_0^{\log 2} \int_0^x \int_0^{x-y} e^{x+y+z} dx dy dz$

Solution: Let $I = \int_0^{\log 2} \int_0^x \int_0^{x-y} e^{x+y+z} dx dy dz$

$$\begin{aligned}
&= \int_0^{\log 2} dx \int_0^x dy e^{x+y} (e^z)_{z=0}^{z=x+y} \\
&= \int_0^{\log 2} dx \int_0^x (e^{2x+2y} - e^{x+y}) dy \\
&= \int_0^{\log 2} \left[e^{2x} \frac{e^{2y}}{2} - e^x e^y \right]_{y=0}^{y=x} dx \\
&= \int_0^{\log 2} \left(\frac{e^{4x}}{2} - \frac{3e^{2x}}{2} + e^x \right) dx
\end{aligned}$$

$$= \left[\frac{1}{8} e^{4x} - \frac{3e^{2x}}{4} + e^x \right]_0^{\log 2} = \frac{5}{8}$$

Example 1.17. Evaluate $\iint xy dx dy$ over the region bounded by the x -axis, ordinate at $x = 2a$ and the parabola $x^2 = 4ay$.

Solution:

In the region y varies from 0 to $\frac{x^2}{4a}$ and x varies from 0 to $2a$.




Fig. 1

$$\begin{aligned}
\therefore I &= \int_0^{2a} \int_0^{x^2/4a} xy dy dx = \int_0^{2a} \left[x \frac{y^2}{2} \right]_0^{x^2/4a} dx \\
&= \int_0^{2a} \frac{x}{2} \cdot \frac{x^4}{16a^2} dx = \frac{1}{32a^2} \int_0^{2a} x^5 dx
\end{aligned}$$

$$= \frac{1}{32a^2} \left[\frac{x^6}{6} \right]_0^{2a} = \frac{a^4}{3}.$$

Example 1.18. Evaluate $\int \int y dx dy$ over the area bounded by $x = 0, y = x^2, x + y = 2$ in the first quadrant.

Solution: We have to find the integral over the region OAB.




Fig. 2

$$I = \int_0^1 \int_{x^2}^{2-x} y dy dx = \int_0^1 \left[\frac{y^2}{2} \right]_{x^2}^{2-x} dx$$

$$= \frac{1}{2} \int_0^1 [(2-x)^2 - x^4] dx$$

$$= \frac{1}{2} \int_0^1 [4 - 4x + x^2 - x^4] dx$$

$$\begin{aligned} &= \frac{1}{2} \left[4x - 2x^2 + \frac{x^3}{3} - \frac{x^5}{5} \right]_0^1 \\ &= \frac{1}{2} \left[4 - 2 + \frac{1}{3} - \frac{1}{5} \right] = \frac{16}{15} \end{aligned}$$

Example 1.19. Evaluate $\int \int (x^2 + y^2) dx dy$ over the area of the triangle whose vertices are $(0, 1), (1, 1), (1, 2)$.

Solution:




Fig. 3

The equation of the line AC is $\frac{y-2}{1} = \frac{x-1}{1}$

(i.e.) $y = x + 1$.

$$\therefore I = \int_0^1 dx \int_1^{x+1} (x^2 + y^2) dy = \int_0^1 \left[x^2 y + \frac{y^3}{3} \right]_1^{x+1} dx$$

1 MULTIPLE INTEGRALS


$$= \int_0^1 \left\{ \left(x^2(x+1) + \frac{(x+1)^3}{3} \right) - \left(x^2 + \frac{1}{3} \right) \right\} dx$$

$$= \frac{1}{3} \int_0^1 (4x^3 + 3x^2 + 3x) dx = \frac{1}{3} \left[x^4 + x^3 + \frac{3x^2}{2} \right]_0^1$$

$$= \frac{1}{3} \left[1 + 1 + \frac{3}{2} \right] = \frac{7}{6}$$

Example 1.20. Evaluate $\iint e^{y/x} dxdy$ where D is the region bounded by the straight lines $y = x$, $y = 0$ and $x = 1$.

Solution: The region D is a triangle as shown in the figure.



In this region x varies from 0 to 1. For each fixed x , y varies from 0 to x .


$$\therefore I = \int_0^1 \int_0^x e^{y/x} dy dx = \int_0^1 [xe^{y/x}]_0^x$$

1 MULTIPLE INTEGRALS

$$= \int_0^1 x(e-1) dx = \frac{1}{2}(e-1)$$

Example 1.21. Evaluate $\iint_D x^2 y^2 dxdy$ where D is the circular disc $x^2 + y^2 \leq 1$.

Solution: In D , x varies from -1 to 1. For a fixed x , y varies from $-\sqrt{1-x^2}$ to $\sqrt{1-x^2}$.



$$\therefore \iint_D x^2 y^2 dxdy = \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} x^2 y^2 dy dx$$

$$= 4 \int_0^1 \int_0^{\sqrt{1-x^2}} x^2 y^2 dy dx \quad [\text{As integrand is even } \int_{-a}^a f(x)dx = 2 \int_0^a f(x)dx \text{ if } f(x) \text{ is even}]$$

$$= 4 \int_0^1 \left[\frac{x^2 y^3}{3} \right]_0^{\sqrt{1-x^2}}$$

$$= \frac{4}{3} \int_0^1 x^2 (1-x^2)^{3/2} dx$$

$$= \frac{4}{3} \int_0^{\pi/2} \sin^2 \theta \cos^4 \theta d\theta \text{ put } x = \sin \theta, dx = \cos \theta d\theta$$

$$= \frac{4}{3} \left(\frac{1.3.1}{2.4.6} \right) \left(\frac{\pi}{2} \right) = \frac{\pi}{24}$$

Example 1.22. Evaluate $I = \iint_D xy dxdy$ where D is the region bounded by the curve $x = y^2$, $x = 2 - y$, $y = 0$ and $y = 1$.

Solution: The given region bounded by the curves is shown in the figure.




Fig. 5

In this region x varies from 0 to 2 when $0 \leq x \leq 1$ for fixed x, y varies from 0 to \sqrt{x} . When $1 \leq x \leq 2$, y varies from 0 to $2 - x$.

\therefore The region D can be subdivided into two regions D_1 and D_2 as shown in the figure.

$$I = \iint_D xy dxdy = \iint_{D_1} xy dxdy + \iint_{D_2} xy dxdy$$

$$= \int_0^1 \int_0^{\sqrt{x}} xy dy dx + \int_1^2 \int_0^{2-x} xy dy dx$$

$$= \int_0^1 \left[\frac{xy^2}{2} \right]_0^{\sqrt{x}} + \int_1^2 \left[\frac{xy^2}{2} \right]_0^{2-x} dx$$

$$= \frac{1}{2} \int_0^1 x^2 dx + \frac{1}{2} \int_1^2 x(2-x)^2 dx$$

$$= \frac{1}{2} \int_0^1 x^2 dx + \frac{1}{2} \int_1^2 (x^3 - 4x^2 + 4x) dx$$

$$= \left[\frac{x^3}{6} \right]_0^1 + \frac{1}{2} \left[\frac{x^4}{4} - \frac{4x^3}{3} + \frac{4x^2}{2} \right]_1^2 = \frac{3}{8}$$

Example 1.23. Evaluate $I = \iiint_D xyz dxdydz$ where D is the region bounded by the positive octant of the sphere $x^2 + y^2 + z^2 = a^2$.

Solution:




Fig. 6

The projection of the given region in the xy plane ($z = 0$) is the

region bounded by the circle $x^2 + y^2 = a^2$ and lying in the 1st quadrant as shown in the figure.

In the given region x varies 0 to a . For a fixed x , y varies from 0 to $\sqrt{a^2 - x^2}$. For a fixed (x, y) , z varies from 0 to $\sqrt{a^2 - x^2 - y^2}$

$$\therefore I = \int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2-y^2}} xyz dz dy dx$$

$$= \int_0^a \int_0^{\sqrt{a^2-x^2}} xy \left(\frac{z^2}{2} \right)_0^{\sqrt{a^2-x^2-y^2}} dy dx$$

$$= \frac{1}{2} \int_0^a \int_0^{\sqrt{a^2-x^2}} xy(a^2 - x^2 - y^2) dy dx$$

$$= \frac{1}{2} \int_0^a \int_0^{\sqrt{a^2-x^2}} x(a^2y - x^2y - y^3) dy dx$$

$$= \frac{1}{2} \int_0^a x \left[a^2 \frac{y^2}{2} - x^2 \frac{y^2}{2} - \frac{y^4}{4} \right]_0^{\sqrt{a^2-x^2}} dx$$

$$= \frac{1}{8} \int_0^a x(a^2 - x^2)^2 dx$$

$$= \frac{1}{8} \int_0^a (a^4x - 2a^2x^3 + x^5) dx$$

$$= \frac{1}{8} \left[a^4 \frac{x^2}{2} - 2a^2 \frac{x^4}{4} + \frac{x^6}{6} \right]_0^a = \frac{a^6}{48}$$

Example 1.24. Evaluate $I = \iiint_D \frac{dxdydz}{(1+x+y+z)^3}$ where D is the region bounded by the planes $x = 0, y = 0, z = 0$ and $x + y + z = 1$.

Solution: The given region is a tetrahedron. The projection of the given region in the xy plane is the triangle bounded by the lines $x = 0, y = 0$ and $x + y = 1$ as shown in the figure. In the




Fig. 7

given region x varies 0 to 1. For each fixed x, y varies from 0 to $1 - x$. For each fixed (x, y) , z varies from 0 to $1 - x - y$.

$$\therefore I = \int_0^1 \int_0^{1-x} \int_0^{1-x-y} \frac{dxdydz}{(1+x+y+z)^3} dy dx$$

$$= -\frac{1}{2} \int_0^1 \int_0^{1-x} [(1+x+y+z)^{-2}]_0^{1-x-y} dy dx$$

$$= -\frac{1}{2} \int_0^1 \int_0^{1-x} \left[\frac{1}{4} - (x+y+1)^{-2} \right] dy dx$$

$$= -\frac{1}{2} \int_0^1 \left[\frac{1}{4}y + (x+y+1)^{-1} \right]_0^{1-x} dx$$

1 MULTIPLE INTEGRALS

$$= -\frac{1}{2} \int_0^1 \left[\frac{1}{4}(1-x) + \frac{1}{2} - (x+1)^{-1} \right] dx$$

$$= -\frac{1}{2} \left[\frac{1}{4}x - \frac{1}{8}x^2 + \frac{1}{2}x - \log(x+1) \right]$$

$$= \frac{1}{2} \log 2 - \frac{5}{16}$$

EXERCISE

1. Evaluate $\int_0^{\pi/2} \int_{\pi/2}^{\pi} \cos(x+y) dx dy$ [Ans: -2]

2. Evaluate $\int_0^1 \int_0^x e^{x+y} dy dx$ [Ans: $\frac{1}{2}(e-1)^2$]

3. Evaluate $\int_0^{\pi} \int_0^{a(1+\cos\theta)} r d\theta dr$ [Ans: $\frac{3}{4}\pi a^2$]

4. Evaluate $\int_0^1 dx \int_0^2 dy \int_1^2 x^2 y z dz$ [Ans: 1]

5. Find the value of $\int_2^3 \int_1^2 \frac{dxdy}{xy}$ [Ans: $\log 2 \log \frac{3}{2}$]

6. Evaluate $\int_0^1 \int_0^2 xy^2 dy dx$. [Ans: 4/3]

7. Find $I = \int_0^{\pi} \int_0^{\pi/2} \int_0^1 r^2 \sin \theta dr d\theta d\phi$ [Ans: $\frac{\pi}{3}$]

8. Find the value of $\int_0^2 \int_1^3 \int_1^2 xy^2 z dz dy dx$ [Ans: 26]

9. Evaluate $\int_0^1 \int_0^z \int_0^{y+z} dz dy dx$ [Ans: 1/2]


1 MULTIPLE INTEGRALS

10. Find the limits of $\iint_R f(x,y) dx dy$, where R is the 1st quadrant bounded by (a) $x = 0, y = 0, x + y = 1$ (b) $x = 0, x = y, y = 1$.

$$\left[\text{Ans: a) } \int_0^1 \int_0^{1-x} f(x,y) dx dy \text{ b) } \int_0^1 \int_0^{1-y} f(x,y) dy dx \right]$$


11. Sketch the region by integration to $\int_0^a \int_0^{\sqrt{a^2-x^2}} f(x,y) dx dy$.

Ans:



12. Sketch the region by integration to $\int_0^1 \int_0^x f(x,y) dx dy$.

Ans:



13. Evaluate $\int_0^1 \int_0^x dx dy$

14. Evaluate $\int_1^2 \int_0^x \frac{1}{x^2+y^2} dx dy$ [Ans: $\frac{\pi}{4} \log 2$]

1 MULTIPLE INTEGRALS

15. Evaluate $\int_0^{\pi/2} \int_0^{a \sin \theta} \int_0^{(a^2 - r^2)/a} r d\theta dr dz$ [Ans: $\frac{5}{64}\pi a^3$]

16. Evaluate $\int_0^2 \int_0^x \int_0^{x+y} e^{x+y+z} dx dy dz$ [Ans: $\frac{e^8 - 3}{8} - \frac{3}{4}e^4 + e^2$]

17. Evaluate $\iint (x^2 - y^2) dx dy$ over the positive quadrant of the circle $x^2 + y^2 = a^2$ [Ans: $\frac{a^5}{15}$]

18. Evaluate $\iint (x + y)^2 dx dy$ over the area bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ [Ans: $\frac{\pi ab}{4}(a^2 + b^2)$]

19. Evaluate $\iint dx dy$ throughout the area bounded by

$$y = x^2, x + y = 2 \quad \left[\text{Ans: } -\frac{10}{3} \right]$$

20. Evaluate $\iint x^2 y^2 dx dy$ over the area of the circle

$$x^2 + y^2 = 1 \quad \left[\text{Ans: } \frac{\pi}{24} \right]$$

21. Evaluate $\iint (x^2 + y^2) dx dy$ over the area bounded by the parabola $y^2 = 4x$ and its latusrectum. [Ans: $\frac{344}{105}$]

22. Evaluate $\int_0^1 \int_x^1 \frac{y dx dy}{x^2 + y^2}$ and also sketch the region of integration. [Ans: $\frac{\pi}{4}$]

1 MULTIPLE INTEGRALS

23. Evaluate $\iint xy dx dy$ over the region in the positive quadrant bounded by the line $2x + 3y = 6$ [Ans: $\frac{3}{2}$]

24. Evaluate $\iint \iint_V (xy + yz + zx) dx dy dz$, where V is the region of sphere bounded by $x = 0, x = 1, y = 0, y = 2, z = 0, z = 3$ [Ans: $\frac{33}{2}$]

25. Evaluate $\iint \iint_D x^2 y z dx dy dz$, where D is the tetrahedron bounded by the planes $x = 0, y = 0, z = 0$ and

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1. \quad \left[\text{Ans: } \frac{a^2 b^2 c^2}{2520} \right]$$

26. Evaluate $\int_0^{\pi/4} \int_0^{a \sin \theta} \frac{r dr d\theta}{\sqrt{a^2 - r^2}}$ [Ans: $\frac{a(\pi - 3)}{6}$]

27. Evaluate $\int_0^4 \int_{y^2/4}^y \frac{y dx dy}{x^2 + y^2}$ and also sketch the region of integration. [Ans: $2 \log 2$]

28. Evaluate $\iint_R x^2 dx dy$, where R is the region bounded by the hyperbola $xy = 4, y = 0, x = 1$ and $x = 2$. [Ans: 6]

29. Evaluate $\iint \iint_V dx dy dz$, where V is the finite region of space formed by the planes $x = 0, y = 0, z = 0$ and $2x + 3y + 4z = 12$. [Ans: 12]

30. Evaluate $\iint_R xy dx dy$, where R is the region bounded by the parabola $y^2 = x$ and the lines $y = 0$ and $x + y = 2$, lying in the 1st quadrant. [Ans: 3/8]

1.2 Change of Order of Integration

Sometimes a double integral may be conveniently evaluated by changing the order of integration. When the limits of the integration are variables, the change of order of integration will involve a change in the limit also. In such cases, it will be convenient to draw a figure indicating the original region of integration.

Example 1.25. Evaluate $\int_0^a \int_y^a \frac{x dx dy}{x^2 + y^2}$ by changing the order of integration.

Solution: The region D of integration is bounded by the lines $x = y$, $x = a$, $y = 0$ and $y = a$ as shown in the figure.




Fig. 8

We have to change the order of integration as $dydx$.

Here in the region D, x varies from 0 to a and for fixed x , y varies from 0 to x .

$$\therefore \int_0^a \int_y^a \frac{x dx dy}{x^2 + y^2} = \int_0^a \int_0^x \frac{x dx dy}{x^2 + y^2} = \int_0^a x \left[\frac{1}{x} \tan^{-1} \left(\frac{y}{x} \right) \right]_0^x$$

$$\begin{aligned} &= \int_0^a x \left[\tan^{-1} \left(\frac{y}{x} \right) \right]_0^x dx = \int_0^a [\tan^{-1} 1 - \tan^{-1} 0] dx \\ &= \int_0^a [\tan^{-1} 1 - \tan^{-1} 0] dx = \frac{\pi}{4} \int_0^a dx \\ &= \frac{\pi}{4} [x]_0^a = \frac{\pi a}{4} \end{aligned}$$

Example 1.26. Evaluate $I = \int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx$ by changing the order of integration.

Solution: The region of integration is as shown. When we change the order x varies from 0 to y and y varies from 0 to ∞ .




Fig. 9

$$\begin{aligned} I &= \int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx = \int_0^\infty \int_0^y \frac{e^{-y}}{y} dx dy \\ &= \int_0^\infty \left[\frac{e^{-y}}{y} x \right]_0^y dy = \int_0^\infty e^{-y} dy \end{aligned}$$

$$= \left[\frac{e^{-y}}{-1} \right]_0^\infty = 1 \left[\frac{e^{-\infty} - e^0}{-1} \right] = \frac{0 - 1}{-1} = 1$$

Example 1.27. Change the order of integration in $\int_0^a \int_x^a (x^2 + y^2) dx dy$ and hence evaluate the same.

Solution: The region of integration is OAB bounded by $x = 0, x = y, y = a$ and $x = a$.

On changing the order of integration $x = 0, x = y$. Then the strip slides from $y = 0$ to $y = a$.




Fig. 10

$$\therefore \int_0^a \int_x^a (x^2 + y^2) dy dx = \int_0^a \int_0^y (x^2 + y^2) dy dx$$

$$= \int_0^a \left[\frac{x^3}{3} + y^2 x \right]_0^y = \int_0^a \left[\frac{y^3}{3} + y^3 \right] dy$$

$$= \frac{4}{3} \int_0^a y^3 dy = \frac{1}{3} [y^4]_0^a = \frac{a^4}{3}$$

Example 1.28. By changing the order of integration, prove that $\int_0^\infty \int_0^y ye^{-y^2/x} dx dy = \frac{1}{2}$

Solution: The region of integration is as shown:




Fig. 11

It is bounded by $x = 0, x = y, y = 0, y = \infty$. On changing the order of integration y ranges from $y = x, y = \infty$ and x ranges from $x = 0$ to $x = \infty$.

$$\int_0^\infty \int_0^y ye^{-y^2/x} dx dy = \int_{x=0}^\infty \int_{y=x}^\infty ye^{-y^2/x} dy dx$$

$$= \int_{x=0}^\infty \left[\int_{y=x}^\infty e^{-y^2/x} d\left(\frac{y^2}{2}\right) \right] dx$$

$$= \int_0^\infty \frac{1}{2} \left[\frac{e^{-y^2/x}}{-1/x} \right]_x^\infty = \int_0^\infty \frac{-x}{2} (0 - e^x) dx$$

$$= \frac{1}{2} \int_0^\infty x e^{-x} dx$$

$$= \frac{1}{2} \left[x \left(\frac{e^{-x}}{-1} \right) - 1 \cdot \frac{e^{-x}}{1} \right]_0^\infty, \text{ on integrating by parts}$$

$$= \frac{1}{2}(0+1) = \frac{1}{2}$$

Example 1.29. Change the order of integration and evaluate
 $I = \int_0^1 \int_0^{\sqrt{1-x^2}} \frac{e^y}{(e^y + 1)\sqrt{1-x^2-y^2}} dx dy$

Solution: The integration is with respect to y between $y = 0, y = \sqrt{1-x^2}$. (i.e.) $y = 0, x^2 + y^2 = 1$ and the next integration is w.r.to x between 0 and 1. The region of integration is as shown here as OAB .




Fig. 12

We now change the order of integration. Integration I w.r.to x between $x = 0, x = \sqrt{1-y^2}$. To cover OAB , y slides from 0 to 1.

$$\begin{aligned} I &= \int_0^1 \int_0^{\sqrt{1-x^2}} \frac{e^y}{(e^y + 1)\sqrt{1-x^2-y^2}} dx dy \\ &= \int_0^1 \int_0^{\sqrt{1-x^2}} \frac{e^y}{(e^y + 1)\sqrt{1-x^2-y^2}} dy dx \end{aligned}$$

$$\begin{aligned} &= \int_0^1 \int_0^{\sqrt{1-y^2}} \frac{e^y}{(e^y + 1)\sqrt{1-x^2-y^2}} dx dy \\ &= \int_0^1 \int_0^{\sqrt{1-y^2}} \left[\frac{e^y}{(e^y + 1)} \frac{dx}{\sqrt{(1-y^2)-x^2}} \right] dy \\ &= \int_0^1 \frac{e^y}{(e^y + 1)} \left[\sin^{-1} \frac{x}{\sqrt{1-y^2}} \right]_0^{\sqrt{1-y^2}} dy = \int_0^1 \frac{e^y}{(e^y + 1)} \frac{\pi}{2} dy \\ &= \frac{\pi}{2} [\log(e^y + 1)]_0^1 = \frac{\pi}{2} \log \left(\frac{e+1}{2} \right) \end{aligned}$$

Example 1.30. By changing the order of integration, evaluate $\int_0^3 \int_1^{4-y} (x+y) dx dy$.

Solution: The region D of integration is bounded by the line $y = 0, y = 3, x = 1$ and by the parabola $x^2 = 4 - y$ as shown in the figure.




Fig. 13

In this region, x varies from 1 to 2 and y varies from 0 to $4 - x^2$.

$$\begin{aligned} \int_0^3 \int_1^{\sqrt{4-y}} (x+y) dx dy &= \int_1^2 \int_0^{4-x^2} (x+y) dy dx \\ &= \int_1^2 \left[xy + \frac{y^2}{2} \right]_0^{4-x^2} = \int_1^2 \left[x(4-x^2) + \frac{(4-x^2)^2}{2} \right] dx \\ &= \int_1^2 \left[\frac{x^4}{2} - x^3 - 4x^2 + 4x + 8 \right] dx \\ &= \left[\frac{x^5}{10} - \frac{x^4}{4} - 4\frac{x^3}{3} + 2x^2 + 8x \right]_1^2 = \frac{241}{60} \end{aligned}$$

Example 1.31. Change the order of integration and evaluate
 $I = \int_0^4 \int_{x^2/4}^{2\sqrt{x}} dy dx$

Solution: The region of integration is bounded by $y = \frac{x^2}{4}$, $y = 2\sqrt{x}$. (i.e) $x^2 = 4y$ and $y^2 = 4x$ which are parabolas and $x = 0$ and 4. Solving $y^2 = 4x$ and $x^2 = 4y$, we get $(0,0), (4,4)$.

After changing the order of integration

$$\begin{aligned} I &= \int_0^4 \int_{x^2/4}^{2\sqrt{x}} dy dx = \int_0^4 \int_{y^2/4}^{2\sqrt{y}} dx dy \\ &= \int_0^4 [x]_{y^2/4}^{2\sqrt{y}} dy = \int_0^4 (2\sqrt{y} - (y^2/4)) dy \\ &= \left[\frac{4}{3}y^{3/2} - \frac{y^3}{12} \right]_0^4 = \frac{32}{3} - \frac{16}{3} = \frac{16}{3} \end{aligned}$$

Example 1.32. Change the order of integration and evaluate




Fig. 14

$$I = \int_0^1 dx \int_x^{\sqrt{2-x^2}} \frac{x}{\sqrt{x^2+y^2}} dy$$

Solution: The limits for y are x and $\sqrt{2-x^2}$ and those for x are 0 and 1. We, therefore, draw the curve $y = x$ which is a straight line and $y = \sqrt{2-x^2}$ which is the upper half of the circle $x^2+y^2 = 2$. The region of integration is OACD. Solving the equation $y = x$ and $x^2+y^2 = 2$, we get the points of intersection $A(1,1)$ and $B(-1,-1)$. If we consider the strip parallel to the x-axis the region has to be divided into two points OAD and ADC. In the region ODA, x varies from 0 to y and y varies from 0 to 1. In the region ADC, x varies from 0 to $\sqrt{2-y^2}$ and y varies from 1 to $\sqrt{2}$.

$$\therefore I = \int_0^1 dx \int_x^{\sqrt{2-x^2}} \frac{x}{\sqrt{x^2+y^2}} dy$$




Fig. 15

$$= \int_0^1 dy \int_0^y \frac{x}{\sqrt{x^2 + y^2}} dx + \int_1^{\sqrt{2}} dy \int_0^{\sqrt{2-y^2}} \frac{x}{\sqrt{x^2 + y^2}} dx = I_1 + I_2$$

$$\text{Now } I_1 = \int_0^1 dy [\sqrt{x^2 + y^2}]_0^y = \int_0^1 (\sqrt{2}y - y) dy$$

$$(\text{i.e.}) \quad I_1 = (\sqrt{2} - 1) \left[\frac{y^2}{2} \right]_0^1 = \frac{1}{2}(\sqrt{2} - 1)$$

$$I_2 = \int_1^{\sqrt{2}} dy [\sqrt{x^2 + y^2}]_0^{\sqrt{2-y^2}} = \int_1^{\sqrt{2}} (\sqrt{2}y - y) dy$$

$$= \left[\sqrt{2}y - \frac{y^2}{2} \right]_1^{\sqrt{2}} = \frac{3}{2} - \sqrt{2}$$

$$I = I_1 + I_2 = 1 - \frac{1}{\sqrt{2}}$$

Example 1.33. Change the order of integration and evaluate
 $I = \int_0^5 dx \int_{2-x}^{2+x} dxdy$

Solution: The I integration is w.r.to y, where $y = 2 - x$ and $y = 2 + x$. The II integration is w.r.to x where $x = 0$ to 5. The region integration is as shown in the figure as ABC. Split ABC as ADC, ADB.




Fig. 16

$$I = I_1 + I_2, \text{ where } I_1 = \int_{-3}^2 \int_{2-y}^5 dx dy \text{ and } I_2 = \int_2^7 \int_{y-2}^5 dx dy$$

$$\text{Now } I_1 = \int_{-3}^2 [x]_{2-y}^5 dy = \int_{-3}^2 (3+y) dy$$

$$= \left[3y + \frac{y^2}{2} \right]_{-3}^2 = 17 - \frac{9}{2} = \frac{25}{2}$$

$$\text{and } I_2 = \int_2^7 [x]_{y-2}^5 dy = \int_2^7 (7-y) dy$$

$$= \left[7y - \frac{y^2}{2} \right]_2^7 = 37 - \frac{49}{2} = \frac{25}{2}$$

$$\therefore I = I_1 + I_2 = \frac{25}{2} + \frac{25}{2} = 25$$

Example 1.34. Change the order of integration and evaluate $\int_0^a \int_{x^2/a}^{2a-x} xy dxdy$

Solution: The I integration is w.r.to y with limits $y = \frac{x^2}{2}$ and $y = 2a - x$. The II integration is w.r.to x between the limits $x = 0, x = a$.

The region of integration is as shown in the figure as OAC. We split this into two regions OAB, ABC. $I = I_1 + I_2$ where I_1 corresponds to OAB and I_2 corresponds to ABC.




Fig. 17

$$I = \int_0^a \int_0^{\sqrt{ay}} xy dxdy + \int_a^{2a} \int_0^{2a-y} xy dxdy$$

$$= \int_0^a y \left(\frac{x^2}{2} \right)_0^{\sqrt{ay}} dy + \int_a^{2a} y \left(\frac{x^2}{2} \right)_0^{2a-y} dy$$

$$\begin{aligned} &= \int_0^a \frac{a}{2} y^2 dy + \frac{1}{2} \int_a^{2a} (4a^2 y - 4ay^2 + y^3) dy \\ &= \frac{a}{2} \left(\frac{y^3}{3} \right)_0^a + \frac{1}{2} \left[2a^2 y^2 - 4a \frac{y^3}{3} + \frac{y^4}{4} \right]_0^a \\ &= \frac{a^4}{6} + \frac{5a^4}{24} = \frac{3a^4}{8} \end{aligned}$$

EXERCISE

1. Change the order of integration in $\int_{-a}^a \int_0^{\sqrt{a^2-y^2}} f(x, y) dxdy$

$$\left[\text{Ans: } \int_0^a \int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} f(x, y) dydx \right]$$

2. Change the order of integration in $\int_0^a \int_y^a \frac{x}{x^2+y^2} dxdy$

$$\left[\text{Ans: } \int_0^a \int_0^x \frac{x}{x^2+y^2} dydx \right]$$

3. Change the order of integration in $\int_0^\infty \int_0^y f(x, y) dxdy$

$$\left[\text{Ans: } \int_0^\infty \int_x^\infty f(x, y) dxdy \right]$$

4. Change the order of integration in $\int_0^1 \int_0^{2\sqrt{x}} \phi(x, y) dydx$

$$\left[\text{Ans: } \int_0^2 \int_{y^2/4}^1 \phi(x, y) dxdy \right]$$

5. Change the order of integration in $\int_0^1 \int_0^x dydx$

1 MULTIPLE INTEGRALS

$\left[\text{Ans: } \int_0^1 \int_y^1 dx dy \right]$

6. Change the order of integration in $\int_0^a \int_{x^2/a}^x f(x, y) dy dx$

$\left[\text{Ans: } \int_0^a \int_y^{\sqrt{ay}} f(x, y) dy dx \right]$

7. Change the order of integration in $\int_0^\infty \int_0^{1/y} f(x, y) dx dy$

$\left[\text{Ans: } \int_0^\infty \int_0^{1/y} f(x, y) dx dy \right]$

8. Change the order of integration and show that $\int_0^1 \int_0^x dy dx = \frac{1}{2}$.

9. Change the order of integration and hence evaluate $\int_1^4 \int_{\sqrt{y}}^2 (x^2 + y^2) dx dy$ $\left[\text{Ans: } \frac{1006}{105} \right]$

10. Change the order of integration and hence evaluate

$\int_0^a \int_0^{\sqrt{ax}} x^2 dy dx$ $\left[\text{Ans: } \frac{2a^4}{7} \right]$

11. Evaluate $\int_0^a \int_0^{\sqrt{1-x^2}} y^2 dy dx$ by interchanging the order of integration. $\left[\text{Ans: } \frac{\pi}{16} \right]$

12. Change the order of integration in $\int_0^a \int_{a-y}^{\sqrt{a^2-y^2}} y dx dy$ and then evaluate it. $\left[\text{Ans: } \frac{a^3}{6} \right]$

1 MULTIPLE INTEGRALS

13. Change the order of integration in $\int_0^{2a} \int_{x^2/4a}^a (x + y) dy dx$ and hence evaluate it. $\left[\text{Ans: } \frac{9a^3}{5} \right]$

14. Evaluate $\int_1^2 \int_0^{4/x} xy dy dx$ by changing the order of integration. $\left[\text{Ans: } 8 \log 2 \right]$

15. Change the order of integration in $\int_0^1 \int_y^{2-y} xy dx dy$ and then evaluate it. $\left[\text{Ans: } \frac{1}{3} \right]$

16. Change the order of integration in $\int_1^2 \int_1^{x^2} \frac{x^2}{y} dy dx$ and hence evaluate the same. $\left[\text{Ans: } \frac{2}{9} \log 2 - 7 \right]$

17. Change the order of integration in $\int_0^a \int_{y^2/a}^y \frac{y dy dx}{(a-x)\sqrt{ax-y^2}}$ $\left[\text{Ans: } \frac{\pi a}{2} \right]$

18. Evaluate $\int_3^5 \int_0^{4/x} xy dx dy$ after reversing the order of integration. $\left[\text{Ans: } 8 \log \frac{5}{3} \right]$

19. Change the order of integration in $\int_0^\infty \int_0^\infty e^{-x^2(1+t^2)} x dx dt$ and then evaluate it. $\left[\text{Ans: } \frac{\pi}{4} \right]$

20. Change the order of integration in each of the double integrals $\int_0^1 \int_1^2 \frac{dx dy}{x^2 + y^2}$ and $\int_1^2 \int_y^2 \frac{dx dy}{x^2 + y^2}$ and hence express

their sum as one double integral and evaluate it. [Ans: $\frac{\pi}{4} \log 2$]

21. Evaluate $\int_0^1 \int_{\sqrt{x}}^1 e^{x/y} dx dy$ by interchanging the order of integration. [Ans: 1/2]

1.3 Plane Area as Double Integral

Plane area enclosed by one or more curves can be expressed as a double integral. Let R be the plane region, the area of which is required. Let us divide the area into a large number of elemental areas $ABCD$, by dividing lines parallel to the y -axis of intervals Δx and the lines parallel to the x -axis, at intervals of Δy .




Fig. 18

$$\text{Area of the rectangle} = \Delta x \Delta y$$

$$\text{Rectangle of area } \Delta = \lim_{\Delta x \rightarrow 0} [\sum \sum \Delta x \Delta y] = \int \int_R dx dy$$

Note: In polar system $A = \int \int_D r dr d\theta$

Example 1.35. Show, by double integration, that the area between the parabolas $y^2 = 4ax$ and $x^2 = 4ay$ is $\frac{16}{3}a^2$.

Solution: The region by integration is as shown.

Solving $y^2 = 4ax$ and $x^2 = 4ay$ are set $(0, 0), (4a, 4a)$. Take a strip parallel to y axis implies limits for $y = \frac{x^2}{4a}$ and $y = 2\sqrt{ax}$ and then x varies from 0 to $4a$.




Fig. 19

$$\therefore \text{Area} = \int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} dx dy = \int_0^{4a} [y]_{x^2/4a}^{2\sqrt{ax}} dx$$

$$= \int_0^{4a} \left(2\sqrt{ax}^{1/2} - \frac{x^2}{4a} \right) dx = \left[2\sqrt{a} \frac{x^{1/2} + 1}{\frac{1}{2} + 1} - \frac{1}{4a} \frac{x^3}{3} \right]_0^{4a}$$

1 MULTIPLE INTEGRALS

$$\begin{aligned}
 &= \left[\frac{4}{3} \sqrt{a} x^{3/2} - \frac{1}{12a} x^3 \right]_0^{4a} = \frac{4}{3} \sqrt{a} (4a^{3/2}) - \frac{1}{12a} (4a)^3 \\
 &= \frac{4}{3} a^2 (4 \times 2) - \frac{1}{12a} \times 64a^3 \\
 &= \frac{32}{3} a^2 - \frac{16}{3} a^2 = \frac{16}{3} a^2 \text{ Sq.units.}
 \end{aligned}$$

Example 1.36. Find by double integration, the area enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Solution: The curve is symmetrical about both axes.




Fig. 20

\therefore Area = 4 area in I quadrant.

$$\begin{aligned}
 &= 4 \int \int_A dy dx = 4 \int_0^a \int_0^{b\sqrt{1-x^2/a^2}} dy dx \\
 &= 4 \int_0^a [y]_0^{b\sqrt{1-x^2/a^2}} = \frac{4b}{a} \int_0^a \sqrt{a^2 - x^2} dx
 \end{aligned}$$

1 MULTIPLE INTEGRALS

$$\begin{aligned}
 &= \frac{4b}{a} \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]_0^a \\
 &= \frac{4b}{a} \times \frac{a^2}{2} \times \frac{\pi}{2} = \pi ab \text{ Sq.units.}
 \end{aligned}$$

Example 1.37. Find by double integration, smaller of the areas bounded by the circle $x^2 + y^2 = 9$ and $x + y = 3$.

Solution: The region of integration is as shown y varies from $3 - x$ to $\sqrt{9 - x^2}$ and x varies from 0 to 3.




Fig. 21

$$\text{Area} = \int \int dy dx$$

$$\begin{aligned}
 &= \int_0^3 \int_{3-x}^{\sqrt{9-x^2}} dy dx = \int_0^3 [\sqrt{9-x^2} - (3-x)] dx \\
 &= \left[\frac{x}{2} \sqrt{9-x^2} + \frac{9}{2} \sin^{-1} \frac{x}{3} \right]_0^3 - 3[x]_0^3 + \left[\frac{x^2}{2} \right]_0^3
 \end{aligned}$$

$$= \frac{9\pi}{2} - 9 + \frac{9}{2} = \frac{9\pi}{4} - \frac{9}{2} = \frac{9}{4}(\pi - 2) \text{ sq.units.}$$

Example 1.38. Find the area between the parabola $y = 4x - x^2$ and the line $y = x$.

Solution:

The two curves intersect at points whose abscissa are given by $4x - x^2 = x$ or $x^2 - 3x = 0$ (i.e.) $x = 0, 3$. Using vertical strips, the required area lies between $x = 0, x = 3$ and $y = x, y = 4x - x^2$.




Fig. 22

$$\begin{aligned}\therefore \text{Required area} &= \int_0^3 \int_x^{4x-x^2} dy dx = \int_0^3 [y]_x^{4x-x^2} dx \\ &= \int_0^3 (4x - x^2 - x) dx = \int_0^3 (3x - x^2) dx \\ &= \left[\frac{3x^2}{2} - \frac{x^3}{3} \right]_0^3 = 4.5 \text{ Sq.units.}\end{aligned}$$

Example 1.39. Find by double integration, the area bounded by the parabola $y = x^2$ and the line $y = 2x + 3$.

Solution: The region of integration is as shown.




Fig. 23

Solving $y = 2x + 3$ we get $x^2 - 2x - 3 = 0$. (i.e.) $x = 3, -1$.

Required area = $\int \int dy dx$ where y varies from $y = x^2$ and $y = 2x + 3$. Further x varies from -1 to 3.

$$\therefore \text{Required area} = \int_{-1}^3 \int_{x^2}^{2x+3} dy dx = \int_{-1}^3 [y]_{x^2}^{2x+3} dx$$

$$= \int_{-1}^3 (2x + 3 - x^2) dx = \left[2\left(\frac{x^2}{2}\right) + 3x - \frac{x^3}{3} \right]_{-1}^3$$

$$= \frac{32}{3} = 10\frac{2}{3} \text{ Sq.units.}$$

Example 1.40. Find the area of the circle using double integral.

Solution: Area of the circle $A = 4 \int \int_D dx dy$ where D is the

1 MULTIPLE INTEGRALS

bounded region in the I quadrant of xy -plane. In D , x varies from 0 to r , and for fixed x , y varies from 0 to $\sqrt{r^2 - x^2}$.




Fig. 24

$$A = 4 \int_0^r \int_0^{\sqrt{r^2 - x^2}} dy dx = 4 \int_0^r \sqrt{r^2 - x^2} \text{ put } x = r \sin \theta$$

$$= 4r^2 \int_0^{\pi/2} \cos^2 \theta d\theta = 4r^2 \int_0^{\pi/2} \frac{1 + \cos 2\theta}{2} d\theta$$

$$= 2r^2 \left[\theta + \frac{\sin 2\theta}{2} \right]_0^{\pi/2} = \pi r^2 \text{ as } \sin \pi = 0, \sin 0 = 0$$

Example 1.41. Find the area of the region D bounded by the parabolas $y = x^2$ and $x = y^2$.

Solution: The point of intersection of the curves are given by $(0,0)$ and $(1,1)$. The required region D is as shown below.

$$\text{The required area } A = \int \int_D dx dy = \int_0^1 \int_{x^2}^{\sqrt{x}} dy dx$$

1 MULTIPLE INTEGRALS




Fig. 25

$$= \int_0^1 [y]_{x^2}^{\sqrt{x}} dx = \int_0^1 (\sqrt{x} - x^2) dx$$

$$= \left[2 \frac{x^{3/2}}{3} - \frac{x^3}{3} \right]_0^1 = \frac{1}{3}$$

Example 1.42. Find by double integration, the area lying inside the circle $r = a \sin \theta$ and outside the cardioid $r = a(1 - \cos \theta)$.

Solution: Eliminating r between the equations of two curves $\sin \theta = 1 - \cos \theta$ or $\sin \theta + \cos \theta = 1$.

Squaring $1 + \sin 2\theta = 1$ or $\sin 2\theta = 0 \therefore 2\theta = 0$ or π

$$\text{(i.e.) } \theta = 0 \text{ or } \frac{\pi}{2}.$$

For the required area, r varies from $a(1 - \cos \theta)$ to $a \sin \theta$ and θ

varies from 0 to $\frac{\pi}{2}$.

$$\begin{aligned}\text{Required area} &= \int_0^{\pi/2} \int_{a(1-\cos\theta)}^{a\sin\theta} r dr d\theta \\ &= \int_0^{\pi/2} \left[\frac{r^2}{2} \right]_{a(1-\cos\theta)}^{a\sin\theta} d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} a^2 [\sin^2\theta - (1 - \cos\theta)^2] d\theta \\ &= \frac{a^2}{2} \int_0^{\pi/2} (2\cos\theta - 2\cos^2\theta) d\theta = a^2 \left(1 - \frac{\pi}{4} \right)\end{aligned}$$




Fig. 26

EXERCISE

- Find the smaller of the areas bounded by the ellipse $4x^2 + 9y^2 = 36$ and the straight line $2x + 3y = 6$. [Ans: $\frac{3}{2}(\pi - 2)$]
- Find by double integration, the area bounded by the parabolas $y^2 = 4 - x$ and $y^2 = 4 - 4x$ [Ans: 8]
- Find the area bounded by the circle $r = 2 \sin\theta$ and $r = 4 \sin\theta$ [Ans: 3π]
- Find the area enclosed by the curve $r^2 = a^2 \cos 2\theta$, by double integration. [Ans: a^2]
- Find the area bounded by the parabola $y^2 = 4 - x$ and $y^2 = x$ by double integration. [Ans: $\frac{16}{3}\sqrt{2}$]
- Find the area common to the parabola $y^2 = x$ and $x^2 + y^2 = 2$ by double integration. [Ans: $\frac{\pi}{2} + \frac{1}{3}$]
- Find the area that lies outside the circle $r = a \cos\theta$ and inside the circle $r = 2a \cos\theta$ [Ans: $\frac{3}{4}\pi a^2$]
- Find the area common to the two circles $r = a$ and $r = 2a \cos\theta$. [Ans: $a^2 \left(\frac{2\pi}{3} - \frac{\sqrt{3}}{2} \right)$]

1.4 Change of Variable from Cartesian to Polar Co-ordinates

Let $\int \int_R f(x, y) dx dy$ be the double integral. $x = r \cos \theta, y = r \sin \theta$ is the transformation from Cartesian to Polar coordinates.

Then $dx dy = |J| dr d\theta$ where $J = \frac{\partial(x, y)}{\partial(r, \theta)}$ is the Jacobian of the transformation and

$$J = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix}$$

$$= r(\cos^2 \theta + \sin^2 \theta) = r$$

$$\therefore \int \int_R f(x, y) dx dy = \int \int_R f(r, \theta) r dr d\theta$$

Example 1.43. Change to polar coordinates and evaluate $\int_0^a \int_y^a x dx dy$

Solution:

The region of integration is $x = y, x = a, y = 0, y = a$.

(i.e.) The triangle OAB putting $x = r \cos \theta, y = r \sin \theta$, the line $x = y$ becomes $r \cos \theta = r \sin \theta$

$$(i.e.) \tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$$




Fig. 27

$$\text{Hence in polar form } I = \int_0^{\frac{\pi}{4}} \int_0^{a/\cos \theta} r^2 \cos \theta dr d\theta$$

$$\begin{aligned} &= \int_0^{\frac{\pi}{4}} \cos \theta \left(\frac{r^3}{3} \right)_0^{a/\cos \theta} d\theta = \frac{a^3}{3} \int_0^{\frac{\pi}{4}} \sec^2 \theta d\theta \\ &= \frac{a^3}{3} [\tan \theta]_0^{\frac{\pi}{4}} = \frac{a^3}{3} \end{aligned}$$

Example 1.44. Change to polar coordinates and evaluate

$$\int_0^a \int_0^{\sqrt{a^2 - x^2}} e^{-(x^2 + y^2)} dx dy$$

Solution:

Putting $x = r \cos \theta, y = r \sin \theta$, the given limit $y^2 = a^2 - x^2$.

(i.e.) The circle $x^2 + y^2 = a^2$ changes to $r = a$ and $y = 0$.




Fig. 28

(i.e.) The x-axis changes to initial line $\theta = 0$. Hence, in the given region r changes from 0 to a and θ changes from 0 to $\frac{\pi}{2}$.

$$I = \int_0^{\frac{\pi}{2}} \int_0^a e^{-r^2} r dr d\theta$$

$$= \int_0^{\frac{\pi}{2}} \left(-\frac{1}{2} e^{-r^2} \right)_0^a d\theta$$

$$= -\frac{1}{2} \int_0^{\frac{\pi}{2}} (e^{-a^2} - 1) d\theta = \frac{\pi}{4} (1 - e^{-a^2})$$

Example 1.45. Evaluate $\int_0^{4a} \int_{y^2/4a}^y dy dx$ by changing to polar coordinates.

Solution:

The region of integration is bounded by the parabola $x = y^2/4a$.




Fig. 29

(i.e.) $y^2 = 4ax$ and the line $x = y$.

By putting $x = r \cos \theta, y = r \sin \theta$, the parabola becomes $r^2 \sin^2 \theta = 4ar \cos \theta$.

(i.e.) $r = \frac{4a \cos \theta}{\sin^2 \theta}$ and the line becomes $x \cos \theta = r \sin \theta$

(i.e.) $\theta = \frac{\pi}{4}$

Hence r varies from 0 to $\frac{4a \cos \theta}{\sin^2 \theta}$ and θ varies from $\frac{\pi}{4}$ to $\frac{\pi}{2}$.

$$I = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^{\frac{4a \cos \theta}{\sin^2 \theta}} r d\theta dr = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \left(\frac{r^2}{2} \right)_0^{\frac{4a \cos \theta}{\sin^2 \theta}}$$

$$\begin{aligned}
 &= \frac{1}{2} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \left[\frac{16a^2 \cos^2 \theta}{\sin^4 \theta} \right] d\theta = 8a^2 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cot^2 \theta \csc^2 \theta d\theta \\
 &= 8a^2 \left[-\frac{\cot^3 \theta}{3} \right]_{\frac{\pi}{4}}^{\frac{\pi}{2}} = -\frac{8a^2}{3}[0 - 1] = \frac{8a^2}{3}
 \end{aligned}$$

Example 1.46. Express the following integral in polar coordinates and evaluate $\int_0^a \int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} \frac{dxdy}{\sqrt{a^2-x^2-y^2}}$

Solution: The limits of y are $\sqrt{ax-x^2}$ and $\sqrt{a^2-x^2}$.




Fig. 30

(i.e.) Upper half of the circles i) $x^2 + y^2 - ax = 0$; $\left(x - \frac{a}{2}\right)^2 + (y - 0)^2 = \left(\frac{a}{2}\right)^2$ and ii) $x^2 + y^2 = a^2$

To change the given integral to polar coordinates, we put $x = r \cos \theta$, $y = r \sin \theta$ and $dxdy = rdrd\theta$.

The equations of the circles now become

i) $r^2 - ar \cos \theta = 0$ (i.e.) $r = a \cos \theta$

ii) $r^2 = a^2$ (i.e.) $r = a$

Hence r changes from $r = a \cos \theta$ to a and θ changes from 0 to $\frac{\pi}{2}$

$$\begin{aligned}
 \therefore I &= \int_0^{\frac{\pi}{2}} \int_{a \cos \theta}^a \frac{rdrd\theta}{\sqrt{a^2 - r^2}} = \int_0^{\frac{\pi}{2}} [-\sqrt{a^2 - r^2}]_{a \cos \theta}^a d\theta \\
 &= \int_0^{\frac{\pi}{2}} a \sin \theta d\theta = [a \cos \theta]_0^{\frac{\pi}{2}} = a
 \end{aligned}$$

Example 1.47. Evaluate $\int \int_R \frac{xydxdy}{\sqrt{x^2+y^2}}$ where R is the region in the first quadrant enclosed by the circles $x^2 + y^2 = a^2$ and $x^2 + y^2 = 4a^2$, by converting into polar coordinates.

Solution: We put $x = r \cos \theta$, $y = r \sin \theta$ $\therefore dxdy = rdrd\theta$

r varies from $r = a$ to $r = 2a$ and θ varies from 0 to $\frac{\pi}{2}$

$$\begin{aligned}
 \text{Hence } I &= \int_0^{\frac{\pi}{2}} \int_a^{2a} \frac{r^3 \sin \theta \cos \theta drd\theta}{\sqrt{r^2}} \\
 &= \left(\int_a^{2a} r^2 dr \right) \left(\frac{1}{2} \int_0^{\frac{\pi}{2}} \sin 2\theta d\theta \right) \\
 &= \frac{1}{2} \left(\frac{r^3}{3} \right)_0^{2a} \left(\frac{-\cos 2\theta}{2} \right)_0^{\frac{\pi}{2}} \\
 &= -\frac{1}{12}(8a^3 - a^3)(\cos \pi - \cos 0)
 \end{aligned}$$




Fig. 31

$$= -\frac{1}{12}(7a^3)(-2) = \frac{7}{6}a^3$$

Example 1.48. Evaluate $\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy$ by using polar coordinates and hence evaluate $\int_0^\infty e^{-(x^2)} dx$.

Solution:

$\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy = \int \int_A e^{-(x^2+y^2)} dA$, where A is the positive quadrant.

Put $x = r \cos \theta, y = r \sin \theta \therefore dx dy = r dr d\theta$

In the positive quadrant A, given the limits for r varies from 0 to ∞ and θ varies from 0 to $\frac{\pi}{2}$

$$I = \int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy = \int_0^{\frac{\pi}{2}} \int_0^\infty e^{-(r^2)} r dr d\theta$$




Fig. 32

$$\text{Put } r^2 = R \Rightarrow 2r dr = dR$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{1}{2} \left(\frac{e^{-R}}{-1} \right)_0^\infty d\theta = \frac{1}{2} \int_0^{\frac{\pi}{2}} d\theta = \frac{1}{2} \times \frac{\pi}{2} = \frac{\pi}{4}$$

$$\text{Since } \int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy = \int_0^\infty e^{-x^2} dx * \int_0^\infty e^{-y^2} dy$$

$$= I \times I = I^2$$

$$I^2 = \frac{\pi}{4}$$

$$\therefore I = \int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$$

Example 1.49. Evaluate $\int_{-a}^a \int_0^{\sqrt{a^2-x^2}} (x^2+y^2) dx dy$ by changing to polar coordinates.

Solution:




Fig. 33

Put $x = r \cos \theta, y = r \sin \theta \therefore dxdy = rdrd\theta$

$$I = \int_{-a}^a \int_0^{\sqrt{a^2 - x^2}} (x^2 + y^2) dxdy$$

$$= \int_0^\pi \int_0^a r^2 \cdot r dr d\theta = \left(\frac{r^4}{4} \right)_0^a [\theta]_0^\pi = \frac{\pi a^4}{4}$$

Example 1.50. Evaluate $\int_0^{2a} \int_0^{\sqrt{2ax - x^2}} (x^2 + y^2) dy dx$.

Solution:

Here y varies from 0 to $\sqrt{2ax - x^2}$

(i.e.) $y = 0$ to $x^2 + y^2 = 2ax$ which is a circle centre $(a, 0)$ and radius a . Also the circle passes through $(0, 0)$. Since x varies from 0 to $2a$, the area of integration is the area above the x-axis and inside the circle. The polar equation of the circle is $x = 2a \cos \theta$.

Changing to polar coordinates $x = r \cos \theta, y = r \sin \theta$




Fig. 34

$$\therefore dxdy = rdrd\theta$$

$$I = \int_0^{\pi/2} \int_0^{2a \cos \theta} r^2 r dr d\theta = \int_0^{\pi/2} \left(\frac{r^4}{4} \right)_0^{2a \cos \theta} d\theta$$

$$= 4a^4 \int_0^{\pi/2} \cos^4 \theta d\theta = 4a^4 \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \frac{3}{4} \pi a^4$$

$$\left[\int_0^{\pi/2} \cos^n \theta d\theta = \begin{cases} \frac{n-1}{n} \times \frac{n-3}{n-2} \times \dots \times \frac{\pi}{2}, & n \text{ is even} \\ \frac{n-1}{n} \times \frac{n-3}{n-2} \times \dots \times 1, & n \text{ is odd} \end{cases} \right]$$

EXERCISE

Changing to polar coordinates evaluate the following:

$$1. \int \int x^2 y^2 dxdy \text{ over the circle } x^2 + y^2 = a^2 \quad \left[\text{Ans: } \frac{\pi}{24} a^6 \right]$$

$$2. \int_0^a \int_y^a \frac{x dx dy}{x^2 + y^2} \quad \left[\text{Ans: } \frac{\pi a}{4} \right]$$

3. $\int_0^a \int_0^x \frac{x^3 dx dy}{\sqrt{x^2 + y^2}}$ [Ans: $\frac{a^4}{4} \log(1 + \sqrt{2})$]

4. $\int_0^a \int_0^{\sqrt{a^2 - x^2}} y^2 \sqrt{x^2 + y^2} dx dy$ over the region bounded by
 $y = 0, x = y, x = 1$ [Ans: $\frac{1}{20} \pi a^5$]

5. $\iint \frac{dxdy}{(x^2 + y^2)^{3/2}}$ over the region bounded by $y = 0, x = y, x = 1$ [Ans: $\frac{\pi}{12}$]

1.5 Volume as a Triple Integral

Triple integrals can be used to evaluate volume V of a finite region D in space.

The volume $V = \iiint_D dxdydz$

Example 1.51. Find the volume of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1 \text{ by triple integrals.}$$

Solution:

Volume = 8 × volume in the first octant.

$$\begin{aligned} V &= 8 \times \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} \int_0^{c\sqrt{1-\frac{x^2}{a^2}-\frac{y^2}{b^2}}} dz dy dx \\ &= 8 \times \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} [z]_0^{c\sqrt{1-\frac{x^2}{a^2}-\frac{y^2}{b^2}}} dy dx \end{aligned}$$

$$= 8c \int_0^a \int_0^{b\sqrt{1-\frac{x^2}{a^2}}} \left[\sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}} \right] dy dx$$

$$\text{Put } r^2 = \left(1 - \frac{x^2}{a^2} \right) b^2$$

$$= \frac{8c}{b} \int_0^a \int_0^r \sqrt{r^2 - y^2} dy dx$$

$$= \frac{8c}{b} \int_0^a \left[\frac{r^2}{2} \sin^{-1} \frac{y}{r} + \frac{y}{2} \sqrt{r^2 - y^2} \right]_0^r dr$$

$$= \frac{2c\pi}{b} \int_0^a r^2 dx = \frac{2c\pi}{b} \int_0^a \left(1 - \frac{x^2}{a^2} \right) b^2 dx$$

$$= 2cb\pi \left(x - \frac{x^3}{3a^2} \right)_0^a = \frac{4\pi}{3} abc$$

Example 1.52. Find the volume bounded by the cylinder $x^2 + y^2 = 4$ and the planes $y + z = 4$ and $z = 0$.

Solution:

z varies from $z = 0$ to $z = 4 - y$ and x, y varies over all points of the circle $x^2 + y^2 = 4$.

$$\begin{aligned} \text{Volume } V &= \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_0^{4-y} dz dy dx \\ &= \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [z]_0^{4-y} dy dx = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (4-y) dy dx \end{aligned}$$




Fig. 35

$$= \int_{-2}^2 \left[4y - \frac{y^2}{2} \right]_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} dx = 8 \times 2 \int_0^2 \sqrt{4-x^2} dx$$

$$V = 16 \left[\frac{x}{2} \sqrt{4-x^2} + \frac{4}{2} \sin^{-1} \frac{x}{2} \right] = 16\pi$$

Example 1.53. Find the volume of the tetrahedron bounded by the coordinate planes and $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$

Solution:

Volume required = $\iiint dxdydz$ with limits.

$$= \int_0^a \int_0^{b(1-\frac{x}{a})} \int_0^{c(1-\frac{x}{a}-\frac{y}{b})} dz dy dx$$

$$= c \int_0^a \int_0^{b(1-\frac{x}{a})} \left(1 - \frac{x}{a} - \frac{y}{b} \right) dy dx$$




Fig. 36

$$\begin{aligned} &= c \int_0^a \left[\left(1 - \frac{x}{a} \right) y - \frac{y^2}{2b} \right]_0^{b(1-\frac{x}{a})} dx \\ &= c \int_0^a \left[b \left(1 - \frac{x}{a} \right)^2 - \frac{b^2}{2b} \left(1 - \frac{x}{a} \right)^2 \right] dx \\ &= \frac{bc}{2} \int_0^a \left(1 - \frac{x}{a} \right)^2 dx = \frac{bc}{2} \left[\frac{\left(1 - \frac{x}{a} \right)^2}{3} \times \left(\frac{-a}{1} \right) \right] \end{aligned}$$

$$= \frac{abc}{6}$$

Example 1.54. Find the volume of the sphere $x^2 + y^2 + z^2 = a^2$ using the triple integrals.

Solution:

1 MULTIPLE INTEGRALS

Since the sphere $x^2 + y^2 + z^2 = a^2$ is symmetric about the coordinate planes, the volume of the sphere = $8 \times$ volume of the first octant.

$$= 8 \times \int \int \int_V dx dy dz$$

$$= 8 \int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2-y^2}} dz dy dx$$

$$= 8 \int_0^a \int_0^{\sqrt{a^2-x^2}} [z]_0^{\sqrt{a^2-x^2-y^2}} dy dx$$

$$= 8 \int_0^a \int_0^{\sqrt{a^2-x^2}} \sqrt{a^2 - x^2 - y^2} dy dx$$

$$\therefore \left[\int \sqrt{(a^2 - x^2)} dx = \frac{x}{2} \sqrt{(a^2 - x^2)} + \frac{a^2}{2} \sin^{-1} \frac{x}{2} \right]$$

$$= 8 \int_0^a \left[\frac{y}{2} \sqrt{(a^2 - x^2) - y^2} + \frac{a^2 - x^2}{2} \sin^{-1} \left(\frac{y}{\sqrt{a^2 - x^2}} \right) \right]_0^{\sqrt{a^2-x^2}} dx$$

$$= 2\pi \int_0^a (a^2 - x^2) dx$$

$$= 2\pi \left[a^2 x - \frac{x^3}{3} \right] = \frac{4\pi}{3} a^3$$

EXERCISE

1. Find the volume bounded by xy -plane, the cylinder $x^2 + y^2 = 4$ and the plane $x + y + z = 3$ $\left[\text{Ans: } \frac{1}{3\pi} (9\pi - 4) \right]$

2. Find the volume in the positive octant bounded by the

1 MULTIPLE INTEGRALS

plane $x + 2y + 3z = 4$ and the coordinate planes. $\left[\text{Ans: } \frac{16}{9} \right]$

3. Evaluate $\int \int \int_V dx dy dz$, where V is the volume enclosed by the cylinder $x^2 + y^2 = 1$ and the planes $z = 0, z = 2 - x$ $\left[\text{Ans: } 2\pi - \frac{4}{3} \right]$