

Box Product

EE25BTECH11008 - Anirudh M Abhilash

October 5, 2025

Problem Statement

If **a**, **b** and **c** are three non-coplanar vectors, then find the value of

$$(\mathbf{a} + \mathbf{b} + \mathbf{c}) \cdot [(\mathbf{a} + \mathbf{b}) \times (\mathbf{a} + \mathbf{c})].$$

Solution

We know that the scalar triple product can be written as a determinant:

$$[\mathbf{p} \ \mathbf{q} \ \mathbf{r}] = \mathbf{p} \cdot (\mathbf{q} \times \mathbf{r}) = \begin{vmatrix} \mathbf{p} & \mathbf{q} & \mathbf{r} \end{vmatrix}.$$

$$(\mathbf{a} + \mathbf{b} + \mathbf{c}) \cdot [(\mathbf{a} + \mathbf{b}) \times (\mathbf{a} + \mathbf{c})] = \begin{vmatrix} \mathbf{a} + \mathbf{b} + \mathbf{c} & \mathbf{a} + \mathbf{b} & \mathbf{a} + \mathbf{c} \end{vmatrix}. \quad (1)$$

Solution (cont..)

$$\begin{vmatrix} \mathbf{a} + \mathbf{b} + \mathbf{c} & \mathbf{a} + \mathbf{b} & \mathbf{a} + \mathbf{c} \end{vmatrix} = \begin{vmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} \end{vmatrix} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix}. \quad (2)$$

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix} = -1. \quad (3)$$

Solution (cont..)

Therefore

$$(\mathbf{a} + \mathbf{b} + \mathbf{c}) \cdot [(\mathbf{a} + \mathbf{b}) \times (\mathbf{a} + \mathbf{c})] = \begin{vmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} \end{vmatrix} \cdot (-1) \quad (4)$$

$$= -[\mathbf{a} \ \mathbf{b} \ \mathbf{c}]. \quad (5)$$

$$(\mathbf{a} + \mathbf{b} + \mathbf{c}) \cdot [(\mathbf{a} + \mathbf{b}) \times (\mathbf{a} + \mathbf{c})] = -[\mathbf{a} \ \mathbf{b} \ \mathbf{c}]$$

C Code (Computations)

https:

`//github.com/Anirudh-EE25BTECH11008/ee1030-2025/tree/
main/EE25BTECH11008/MATGEO/2.10.37/Codes/verify.c`

Python Code (Verifying)

https:

`//github.com/Anirudh-EE25BTECH11008/ee1030-2025/tree/
main/EE25BTECH11008/MATGEO/2.10.37/Codes/verify.py`