

RESEARCH ARTICLE

Distinguishing internally club and approachable on an Infinite interval

Hannes Jakob  | Maxwell Levine Mathematisches Institut, University of
Freiburg, Freiburg im Breisgau, Germany**Correspondence**Maxwell Levine, Ernst-Zermelo-Strasse 1,
79100 Freiburg im Breisgau, Germany.
Email: [maxwell.levine@mathematik.
uni-freiburg.de](mailto:maxwell.levine@mathematik.uni-freiburg.de)**Abstract**

Krueger showed that PFA implies that for all regular $\Theta \geq \aleph_2$, there are stationarily many $[H(\Theta)]^{\aleph_1}$ that are internally club but not internally approachable. From countably many Mahlo cardinals, we force a model in which, for all positive $n < \omega$ and $\Theta \geq \aleph_{n+1}$, there is a stationary subset of $[H(\Theta)]^{\aleph_n}$ consisting of sets that are internally club but not internally approachable. The theorem is obtained using a new variant of Mitchell forcing. This answers questions of Krueger.

MSC 2020

03E35, 03E55 (primary)

1 | INTRODUCTION

Skolem originally regarded what is now known as the Löwenheim–Skolem theorem as an illustration of the paradoxical nature of infinite cardinals, particularly when uncountable cardinals are considered as elements of countable models. Despite this original motivation, this theorem is nowadays used as an essential tool for proving numerous positive results in combinatorial set theory. Foreman and Todorčević deepened the study of the Löwenheim–Skolem theorem and introduced variations of the concept of internal approachability, which considers the internal composition of elementary submodels [4]. We are left with a situation in which the formal construction of sets gives rise to its own sort of geometry in which subtle distinctions can be made regarding the structure of such models.

Following the work of Foreman and Todorčević, Krueger wrote a series of papers exploring variations of internal approachability, in particular proving that the variations are distinct [10]. He showed that these distinctions can be obtained using mixed-support iterations, which resemble

the forcings Mitchell used to obtain the tree property at double successor cardinals. Notable developments in the study of the tree property pertain to obtaining the tree property simultaneously on long intervals of cardinals, and this area of research requires analyses of variants of Mitchell's forcing. In this spirit, Krueger raised the question of whether these properties could be separated for successive cardinals, or even an infinite sequence of cardinals [12]. We studied the case in which internally stationary is distinguished from internally club by using forcings that accomplish the work of mixed support iterations while more explicitly resembling Mitchell's forcing [7, 13].

In this paper, we will demonstrate the robustness of this idea by addressing the separation of internally club from internally approachable. We introduce a new version of Mitchell forcing, for which we must consider somewhat elaborate termspaces. The benefit is derived from having an Abraham-style projection analysis. We hope that this concept will be useful for points in the literature where mixed support iterations are called for (see [5], for example).

The concepts we study here are framed in terms of the notion of stationarity for spaces of the form $[X]^{\leq \mu}$, which was formulated by Jech (see [9]). For a regular cardinal μ , we say that some $N \in [X]^{\leq \mu}$ is:

- *internally unbounded* if $[N]^{< \mu} \cap N$ is unbounded in $[N]^{< \mu}$,
- *internally stationary* if $[N]^{< \mu} \cap N$ is stationary in $[N]^{< \mu}$,
- *internally club* if $[N]^{< \mu} \cap N$ contains a club in $[N]^{< \mu}$,
- *internally approachable* if there is a continuous sequence $\langle N_i : i < \mu \rangle$ consisting of elements of $[N]^{< \mu}$ such that for all $i < \mu$, $\langle N_j : j \leq i \rangle \in N$ and $N = \bigcup_{i < \mu} N_i$.

For clarity, let $\text{ICNIA}(\Theta, \mu)$ be the statement that $\Theta \geq \mu^+$ and that there exist stationarily many $N \in [H(\Theta)]^{\leq \mu}$ that are internally club but not internally approachable. In all cases, we will be assuming that μ is regular and uncountable for the following reason: If μ is singular, there cannot be stationarily many elementary substructures $N < H(\Theta)$ with $|N| = \mu$ such that $[N]^{< \mu} \cap N$ is unbounded in $[N]^{< \mu}$, since $N \cap \mu^+$ (which is an ordinal for club many N) will always have cofinality $< \mu$. Additionally, the property can never hold for $\mu = \omega$ because any countable $M < H(\Theta)$ is internally approachable. Thus, if we work in a model without weakly inaccessible cardinals, the optimal global result can be achieved by just forcing the property $\text{ICNIA}(\Theta, \mu)$ to hold for all successor cardinals μ . Krueger showed that PFA implies $\text{ICNIA}(\Theta, \aleph_1)$ for all $\Theta \geq \aleph_2$ [10] and later showed that $\text{ICNIA}(\mu^+, \mu)$ is consistent from a Mahlo cardinal for regular μ [12]. We extend that result here.

Theorem 1.1. *Assume that there are countably many Mahlo cardinals in V . Then, there is a forcing extension in which, for all $1 \leq n < \omega$, $\text{ICNIA}(\Theta, \aleph_n)$ holds for all $\Theta \geq \aleph_{n+1}$.*

This resolves a case of [12, Question 12.9], where the projection analysis allows us to obtain consecutive instances of $\text{ICNIA}(\Theta, \aleph_n)$. It also resolves a case of [12, Question 12.7], where the idea of the solution is more or less that the image of the Mostowski collapse cannot distinguish between $H(\Theta)$'s for large Θ .

2 | THE NEW FORCING

In this section, we will define the new forcing and present a simple application before moving on to the proof of our main theorem.

2.1 | Defining the forcing

The idea of our forcing is to take the two-step iteration used to establish ICNIA(Θ, \aleph_1) and build it into a variant of Mitchell forcing that enjoys some of the nice properties of the more standard variants. First, we need the collapse that Krueger used, which forces a chain through a stationary set.

Definition 2.1 (See [10]). Let μ be cardinals and $S \subseteq [X]^{<\mu}$ be stationary for some set X . The *chain forcing* $\mathbb{P}(S)$ consists of closed sequences of length $< \mu$ through S , that is, it consists of sequences s such that $\text{dom}(s)$ is a successor ordinal below μ , $s(\alpha) \in S$ for all $\alpha \in \text{dom}(s)$, and $s(\gamma) = \bigcup_{\alpha < \gamma} s(\alpha)$ for all limit $\gamma \in \text{dom}(s)$.

The poset $\mathbb{P}(S)$ is used because it allows us to collapse $|X|$ while retaining both the approximation property and the clubness of the “old” sets.

Fact 2.2. Let $\mu \leq \delta$ be cardinals and $S \subseteq [H(\delta)]^{<\mu}$ stationary.

- (1) $\mathbb{P}(S)$ adds an increasing and continuous sequence of elements $\langle S_i : i < \mu \rangle$ of S with union $H(\delta)^V$, thus collapsing δ to have cardinality μ .
- (2) If $\delta^{<\delta} = \delta$, then $\mathbb{P}(S)$ has cardinality δ and therefore is δ^+ -c.c.

Note that to obtain $< \mu$ -distributivity of $\mathbb{P}(S)$, S needs to satisfy some additional assumptions.

Now we are ready to define our Mitchell forcing. We need to take some care regarding the model in which the Cohen sets are defined. This is analogous to constructions in which the tree property holds on an interval of cardinals (see [3]), and is done here in anticipation of the iteration used to prove Theorem 1.1. We will therefore use the following basic fact from here on without comment.

Fact 2.3. Suppose that $W \subseteq V$ are models of set theory such that τ is regular and κ is inaccessible in W . Suppose also that τ and κ are cardinals in V and that the extension $W \subseteq V$ has the κ -covering property. Then, $\text{Add}^W(\tau, \kappa)$, the version of $\text{Add}(\tau, \kappa)$ defined in W , has the κ -Knaster property in V . (See [3, Lemma 2.6] and [1].)

Definition 2.4. Let $\tau < \mu < \kappa$ be cardinals in V such that $\tau^{<\tau} = \tau$ and κ is inaccessible. Let $W \subseteq V$ be models of ZFC containing the ordinals such $(\text{Add}(\tau, \kappa))^W$ is μ -Knaster. Then, we define $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ in V to be the poset consists of pairs (p, q) such that:

- (1) $p \in (\text{Add}(\tau, \kappa))^W$,
- (2) q is a $< \mu$ -sized function on κ such that
 - (a) for each $\alpha \in \text{dom}(q)$, $\alpha = \delta + 1$ for an inaccessible cardinal δ ,
 - (b) $q(\alpha)$ is an $(\text{Add}(\tau, \alpha))^W$ -name for an element in the chain forcing $\mathbb{P}([H(\delta)]^{<\mu} \cap V[(\text{Add}(\tau, \delta))^W])$.

We let $(p', q') \leq_{\mathbb{M}^\oplus(\tau, \mu, \kappa, W)} (p, q)$ if

- (1) $p' \leq_{(\text{Add}(\tau, \kappa))^W} p$,
- (2) $\text{dom}(q') \supseteq \text{dom}(q)$ and for all $\alpha \in \text{dom}(q)$,

$$p' \restriction \alpha \Vdash q'(\alpha) \leq_{\mathbb{P}([H(\delta)]^{<\mu} \cap V[(\text{Add}(\tau, \delta))^W])} q(\alpha).$$

For simplicity, we define $\mathbb{M}^\oplus(\tau, \mu, \kappa) := \mathbb{M}^\oplus(\tau, \mu, \kappa, V)$.

Given our definition of the Mitchell forcing, it then becomes clear that we can define a termspace forcing, which is essentially the main benefit of this presentation.

Definition 2.5. Let $\mathbb{T} = \mathbb{T}(\mathbb{M}^\oplus(\tau, \mu, \kappa, W))$ be the termspace of $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$. For the sake of explicitness, this is the poset consisting of conditions q such that:

- (2) q is a μ -sized function such that for each $\alpha \in \text{dom}(q)$:
 - (a) $\alpha = \delta + 1$ for an inaccessible cardinal δ ,
 - (b) $q(\alpha)$ is an $(\text{Add}(\tau, \alpha))^W$ -name for an element in the chain forcing $\mathbb{P}([H(\delta)]^{<\mu} \cap V[(\text{Add}(\tau, \delta))^W])$.

Most importantly, we let $q' \leq q$ if and only if:

- (1) $\text{dom}(q') \supseteq \text{dom}(q)$,
- (2) for all $\alpha \in \text{dom}(q)$, $\Vdash_{\mathbb{P}([H(\delta)]^{<\mu} \cap V[(\text{Add}(\tau, \delta))^W])} "q'(\alpha) \leq q(\alpha)"$.

Next, we will establish strategic closure properties of our forcing.

Definition 2.6. Let \mathbb{P} be a forcing order, δ an ordinal. The *completeness game* $G(\mathbb{P}, \delta)$ on \mathbb{P} with length δ has players COM (complete) and INC (incomplete) playing elements of \mathbb{P} with COM playing at even ordinals (i.e., limit ordinals and ordinals of the form $\alpha + n$ for α a limit and $n < \omega$) and INC playing at odd ordinals. COM starts by playing $1_{\mathbb{P}}$, afterward p_α has to be a lower bound of $(p_\beta)_{\beta < \alpha}$. INC wins if either player is unable to play at some point $< \delta$. Otherwise, COM wins.

A poset \mathbb{P} is δ -strategically closed if COM has a winning strategy for the game $G(\mathbb{P}, \delta)$. We say that \mathbb{P} is *strongly δ -strategically closed* if COM has a winning strategy for the version of the game where they play at odd ordinals and INC plays at even ordinals (see [2] for background on these definitions).

The subtlety here is that, even though $\mathbb{P}(S)$ is in most cases not μ -strategically closed since it destroys the stationarity of a subset of $[\delta]^{<\mu}$, the term ordering on $\text{Add}(\tau) * \mathbb{P}([\delta]^{<\mu} \cap V)$ is strongly μ -strategically closed.

Lemma 2.7. Let $\tau < \mu < \delta$ be cardinals such that $\tau^{<\tau} = \tau$. Then, the term forcing $\mathbb{T}(\mathbb{M}^\oplus(\tau, \mu, \kappa, W))$ is strongly μ -strategically closed.

Proof. Since products of strongly μ -strategically closed forcings are strongly μ -strategically closed, it is sufficient to argue that the direct extension ordering on $\text{Add}(\tau, \gamma + 1)^W * \mathbb{P}([\delta]^{<\mu} \cap V[\text{Add}(\tau, \gamma)^W])$, that is, the ordering \leq^* for which $(p, q) \leq^* (p', q')$ holds if and only if $p = p'$ and $q \leq q'$, is strongly μ -strategically closed.[†] In order to do so, we will first show that this is true for $\gamma = 0$. We will suppress notation for the inner model W in this proof for the sake of readability.

We give a winning strategy for COM by describing a play of the game of the form $\langle (p, q_\gamma) : \gamma < \mu \rangle$ where $p \in \text{Add}(\tau)$. At any odd stage γ , COM will play \dot{q}_γ such that the following holds.

- (1) There is ν_γ such that $p \Vdash \text{dom}(\dot{q}_\gamma) = \check{\nu}_\gamma + 1$.
- (2) There is $x_\gamma \in V$ such that $p \Vdash \dot{q}_\gamma(\check{\nu}_\gamma) = \check{x}_\gamma$.

We will argue both that this choice will be possible at every stage and that this is sufficient to keep the game going.

[†] The strategic closure of the direct extension ordering was first noticed by Krueger with a different proof [11].

Suppose that γ is a limit ordinal: If COM has played according to the strategy until γ , we let $\nu_\gamma := \bigcup \{\nu_\alpha : \alpha < \gamma, \alpha \in \text{Odd}\}$ and $x_\gamma := \bigcup \{x_\alpha : \alpha < \gamma, \alpha \in \text{Odd}\}$. Then, we can find a lower bound: Let \dot{q}_γ be a name for a condition with domain $\nu_\gamma + 1$ such that $\dot{q}_\gamma(\alpha) = \dot{q}_\beta(\alpha)$ for some $\beta < \alpha$ whenever $\alpha < \nu_\gamma$ and such that $\dot{q}_\gamma(\nu_\gamma) = x_\gamma$. In particular, this works because we have made it explicit that $x_\gamma \in V$.

Now assume $\gamma = \beta + 1$ is a successor ordinal and INC has just played \dot{q}_β . Because $\text{Add}(\tau)$ is μ -Knaster and in particular has the $< \mu$ -covering property, $\nu'_\gamma := \sup\{\nu \mid \exists p' \leq p(p' \Vdash \text{dom}(\dot{q}_\beta) = \check{\nu})\}$ is below μ and $x_\gamma := \{\epsilon \mid \exists p' \leq p(p' \Vdash \epsilon \in \bigcup \dot{q}_\beta)\}$ has size $< \mu$. Let \dot{q}_γ be a function with domain $\nu'_\gamma + 1$ extending \dot{q}_β and such that $\dot{q}_\gamma(\nu'_\gamma) = \check{x}_\gamma$.

We show that \dot{q}_γ is as required: \dot{q}_γ is obviously forced to extend \dot{q}_β . Furthermore, the values of \dot{q}_γ are forced to be elements of V : Until ν'_γ , this holds because \dot{q}_β is forced to be in $\mathbb{P}([\delta]^{<\mu} \cap S)$. At ν'_γ , it holds because $\dot{q}_\gamma(\nu'_\gamma)$ is the check-name \check{x}_γ . Lastly, \dot{q}_γ is continuous at every limit and increasing. Because $\text{dom}(\dot{q}_\gamma) = \nu_\gamma + \check{\alpha} + 1$ and $\dot{q}_\gamma(\check{\nu}_\gamma + \alpha) = \check{x}_\gamma$, we are done.

Now let us show how the preceding fact shows that the direct extension ordering on $\text{Add}(\tau, \gamma + 1) * \mathbb{P}([\delta]^{<\mu} \cap V[\text{Add}(\tau, \gamma)])$ is μ -strategically closed. The former poset is isomorphic to $\text{Add}(\tau, \gamma) * (\text{Add}(\tau) * \mathbb{P}([\delta]^{<\mu} \cap V))$ (where we treat V as the new ground model). Ergo $\text{Add}(\tau, \gamma)$ forces that there exists a winning strategy $\dot{\sigma}$ for COM in the completeness game on the direct extension ordering on $\text{Add}(\tau) * \mathbb{P}([\delta]^{<\mu} \cap V)$. This allows us to define a winning strategy σ for COM in the completeness game on the direct extension ordering on $(\text{Add}(\tau, \gamma) * \text{Add}(\tau)) * \mathbb{P}([\delta]^{<\mu} \cap V[\text{Add}(\tau, \gamma)])$ by letting

$$\sigma(((p \upharpoonright \gamma, p(\gamma)), q_\alpha)_{\alpha < \beta}) := (p \upharpoonright \gamma, \dot{\sigma}((p(\gamma), q_\alpha))_{\alpha < \beta}).$$

□

In particular, by Easton's lemma, $\mathbb{P}([\delta]^{<\mu} \cap V)$ is $< \mu$ -distributive (actually strongly $< \mu$ -distributive, see [7], Def. 3.1) in $V[\text{Add}(\tau)]$.

We note that what we have given is actually a winning *tactic*, that is, in successor stages, the play by COM depends only on the last play of INC, not on the plays before that (see [16]).

Now we can show that \mathbb{M}^\oplus has similar properties to more standard versions of Mitchell forcing.

Proposition 2.8. *Let $\mathbb{M}^\oplus = \mathbb{M}^\oplus(\tau, \mu, \kappa, W)$:*

- (1) \mathbb{M}^\oplus is κ -Knaster,
- (2) \mathbb{M}^\oplus is a projection of the product $\text{Add}(\tau, \kappa)^W \times \mathbb{T}(\mathbb{M}^\oplus(\tau, \mu, \kappa, W))$,
- (3) \mathbb{M}^\oplus forces $\kappa = 2^\tau = \mu^+ = \tau^{++}$.

Sketch. Recall that the first point follows from a Δ -system argument, the second point uses some mixing of forcing names and the third point uses the first two points along with Easton's Lemma.

The next point will be the crux of what is needed to bring Krueger's arguments into our context.

Lemma 2.9. $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ has the $< \mu$ -approximation property.

The argument uses the fact that $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ is *iteration-like*, meaning that we can mix the conditions in the second coordinate to “move disagreements into the first coordinate.”[†] Our argument here uses ideas of Usuba [15] and Unger [14].

[†] See [7] for a generalization that uses the notion of strong distributivity, due to the first author.

Proof of Lemma 2.9. Let us abbreviate $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ as \mathbb{M} .

Claim 2.10. Suppose that $(p, q) \in \mathbb{M}$ forces $\dot{x} \in V$ but that there is no $y \in V$ such that (p, q) forces $\dot{x} = \check{y}$. Then, there are $q', p_0, p_1 \leq p$ and $y_0 \neq y_1$ such that $q' \leq_{\mathbb{T}} q$ and $y_i \in V$ and $(p_i, q') \Vdash \dot{x} = \check{y}_i$ for $i \in 2$.

Proof. We consider two possible cases:

Case 1: There are q^* and y_0 such that $(p, q^*) \leq_{\mathbb{M}} (p, q)$ and $(p, q^*) \Vdash \dot{x} = \check{y}_0$.

Then choose $(p_1, q^{**}) \leq_{\mathbb{M}} (p, q)$ and some y_1 such that $(p_1, q^{**}) \Vdash \dot{x} = \check{y}_1$. Strengthen if necessary to assume that p_1 is strictly below p and choose $p_0 \leq p$ incompatible with p_1 . Using standard arguments for the construction of names, there is q' such that $q' \leq_{\mathbb{T}} q$ and such that for all $\alpha \in \text{dom } q^{**}$, $p_1 \Vdash q^{**}(\alpha) = q'(\alpha)$ and for all $\alpha \in \text{dom } q^*$, $p_0 \Vdash q^*(\alpha) = q'(\alpha)$. Then, we have $(p_0, q') \leq_{\mathbb{M}} (p_0, q^*) \leq_{\mathbb{M}} (p, q^*) \leq_{\mathbb{M}} (p, q)$ and $(p_1, q') \leq_{\mathbb{M}} (p_1, q^{**}) \leq_{\mathbb{M}} (p, q)$, and so, have this case of the claim.

Case 2: For all q^* with $(p, q^*) \leq_{\mathbb{M}} (p, q)$, $(p, q^*) \nVdash \dot{x} = \check{y}_0$ for any y_0 .

Then choose $(p_0, q^*) \leq_{\mathbb{M}} (p, q)$ forcing $\dot{x} = \check{y}_0$ for some y_0 . Using the mixing of names, we can assume that $q^* \leq_{\mathbb{T}} q$, and hence that $(p, q^*) \leq_{\mathbb{M}} (p, q)$. The present case implies that $(p, q^*) \nVdash \dot{x} = \check{y}_0$, so there is some $(p_1, q') \leq_{\mathbb{M}} (p, q^*)$ forcing $\dot{x} = \check{y}_1$ for some $y_1 \neq y_0$. Again, we can assume that $q' \leq_{\mathbb{T}} q^*$. Therefore, $(p_0, q') \leq_{\mathbb{M}} (p_0, q^*) \leq_{\mathbb{M}} (p, q^*) \leq_{\mathbb{M}} (p, q)$ and $(p_1, q') \leq_{\mathbb{M}} (p, q^*) \leq_{\mathbb{M}} (p, q)$. \square

Now suppose for contradiction that the lemma is false. Let \dot{f} be an \mathbb{M} -name such that some (p, q) forces every $< \mu$ -approximation to be in V , but \dot{f} itself to be outside of V . For simplicity, assume $(p, q) = 1_{\mathbb{M}}$.

We will use the winning strategy for COM in the completeness game of length μ played on \mathbb{T} . More precisely, the values of q_γ chosen for $\gamma \in \text{Even}$ are chosen by INC, and the construction continues because of the winning strategy for COM.

We will construct $(p_\gamma^0, p_\gamma^1, q_\gamma, y_\gamma)_{\gamma \in \text{Even}}$ such that

- (1) $y_\gamma \in [V]^{<\mu} \cap V$, the sequence $(y_\gamma)_{\gamma \in \text{Even}}$ is \subseteq -increasing,
- (2) the q_γ 's are $\leq_{\mathbb{T}}$ -decreasing,
- (3) (p_γ^0, q_γ) and (p_γ^1, q_γ) decide $\dot{f} \restriction \check{y}_\alpha$ the same way for $\alpha < \gamma$, but differently for $\alpha = \gamma$.

Assume the game has been played until some even ordinal $\gamma < \mu$. Let $y'_{\gamma+1} := \bigcup_{\alpha < \gamma} y_\alpha$, which has size $< \mu$. Because \dot{f} is forced not to be in V , there is $y_{\gamma+1} \supseteq y'_{\gamma+1}$ of size $< \mu$ such that $(1_{\mathbb{A}}, q_\gamma)$ does not decide $\dot{f} \restriction \check{y}_{\gamma+1}$. Thus, we find all required objects by appealing to Claim 2.10. Formally, we can choose the splitting below (p_γ^0, q_γ) at every step.

We claim that $\{(p_\gamma^0, p_\gamma^1) \mid \gamma \in \text{Even}\}$ is an antichain in $\mathbb{A} \times \mathbb{A}$ where $\mathbb{A} := \text{Add}(\tau, \kappa)^W$, obtaining a contradiction since it is known that $\mathbb{A} \times \mathbb{A}$ has the μ -chain condition. To this end, assume $(p^0, p^1) \leq_{\mathbb{A} \times \mathbb{A}} (p_{\gamma'}^0, p_{\gamma'}^1), (p_{\gamma'}^0, p_{\gamma'}^1)$ with $\gamma > \gamma'$. Because $p^0 \leq_{\mathbb{A}} p_{\gamma'}^0$ and $p^1 \leq_{\mathbb{A}} p_{\gamma'}^1$, $(p^0, q_\gamma) \leq_{\mathbb{M}} (p^0, q_{\gamma'}) \leq_{\mathbb{M}} (p_{\gamma'}^0, q_{\gamma'})$ and $(p^1, q_\gamma) \leq_{\mathbb{M}} (p^1, q_{\gamma'}) \leq_{\mathbb{M}} (p_{\gamma'}^1, q_{\gamma'})$ decide $\dot{f} \restriction \check{y}_{\gamma'}$ differently, but because $p^0 \leq_{\mathbb{A}} p_{\gamma'}^0$ and $p^1 \leq_{\mathbb{A}} p_{\gamma'}^1$, $(p^0, q_\gamma) \leq_{\mathbb{M}} (p_{\gamma'}^0, q_\gamma)$ and $(p^1, q_\gamma) \leq_{\mathbb{M}} (p_{\gamma'}^1, q_\gamma)$ decide $\dot{f} \restriction \check{y}_{\gamma'}$ the same way, a contradiction. \square

2.2 | Considering quotients

As is common when working with variants of Mitchell forcing, we give an explicit description of the quotient forcing. In this subsection, we will define and state what we need in order to carry

out the proof of our main theorem, leaving out some of the details that are addressed elsewhere in the literature.

Definition 2.11. Let $\tau < \mu < \nu < \kappa$ be cardinals such that $\text{Add}(\tau, \kappa)^W$ is μ -Knaster. Let $G \subseteq \mathbb{M}^\oplus(\tau, \mu, \nu, W)$ be a generic filter. In $V[G]$, define $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ to consist of (p, q) such that

- (1) $p \in \text{Add}(\tau, \kappa \setminus \nu)^W$,
- (2) q is a partial function on $\kappa \setminus \nu$ of size $< \mu$ such that for each $\alpha \in \text{dom}(q)$, $\alpha = \nu + 1$ for an inaccessible cardinal ν and $q(\alpha)$ is an $\text{Add}(\tau, \alpha \setminus \nu)$ -name for an element of $\mathbb{P}([\nu]^{<\mu} \cap V[G][\text{Add}(\tau, \nu \setminus \nu)])$.

We let $(p', q') \leq (p, q)$ if

- (1) $p' \leq p$,
- (2) $\text{dom}(q') \supseteq \text{dom}(q)$ and for all $\alpha \in \text{dom}(q)$,

$$p' \restriction \alpha \Vdash q'(\alpha) \leq q(\alpha).$$

The next lemma follows similarly to other known variants of Mitchell forcing.

Lemma 2.12. Let $\tau < \mu < \nu < \kappa$ be regular cardinals such that $\tau^{<\tau} = \tau$ and ν, κ are inaccessible. There is a dense embedding from $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ into $\mathbb{M}^\oplus(\tau, \mu, \nu, W) * \mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$.

Proof. As in other versions of Mitchell forcing, we define

$$(p, q) \mapsto (p \restriction \nu, q \restriction \nu, \text{op}(p \restriction (\check{\kappa} \setminus \nu), \bar{q})),$$

where \bar{q} reimagines $q \restriction (\kappa \setminus \nu)$ as an $\mathbb{M}^\oplus(\tau, \mu, \nu, W)$ -name. □

Similarly, we have the following.

Proposition 2.13. Let $\tau < \mu < \nu < \kappa$ be cardinals such that $\tau^{<\tau} = \tau$ and ν, κ are inaccessible and let G be $\mathbb{M}^\oplus(\tau, \mu, \nu, W)$ -generic over V and let H be $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ -generic over $V[G]$. Then, there is a filter K_A that is $\text{Add}(\tau)$ -generic over $V[G]$ and a filter K_C that is $\mathbb{P}([\nu]^{<\mu} \cap V[G])$ -generic over $V[G][K_A]$ such that $V[G][H]$ is a forcing extension of $V[G][K_A][K_C]$.

Proof. Use a map similar to the one from the previous lemma. This is where we use the fact that $\text{dom}(q)$ consists of ordinal successors of inaccessibles for $(p, q) \in \mathbb{M}^\oplus(\tau, \mu, \kappa, W)$. Here, we also note that $\mathbb{P}([\nu]^{<\mu} \cap V[G]) = \mathbb{P}([\nu]^{<\mu} \cap V[A])$ where A is the $\text{Add}(\tau, \nu)$ -generic induced by G . □

In $V[G]$, $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ has similar properties to $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ using arguments similar to the ones we detailed:

Lemma 2.14. Let $\tau < \mu < \nu < \kappa$ be cardinals such that $\text{Add}(\tau, \kappa)^W$ is μ -Knaster and ν, κ are inaccessible. Let G be $\mathbb{M}^\oplus(\tau, \mu, \nu, W)$ -generic. The following holds in $V[G]$.

- (1) $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ is κ -Knaster.
- (2) The term ordering on $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ is strongly μ -strategically closed.

It is crucial for us to obtain the approximation property for quotients, which we can obtain from trivial modifications of the proof of Lemma 2.9.

Lemma 2.15. *Let $\tau < \mu < \nu < \kappa$ be cardinals such that $\tau^{<\tau} = \tau$ and ν, κ are inaccessible. Let G be $\mathbb{M}^\oplus(\tau, \mu, \nu, W)$ -generic. In $V[G]$, $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ has the $< \mu$ -approximation property.*

2.3 | Distinguishing internally club and approachable for a single cardinal

In this subsection, we show that $\mathbb{M}^\oplus(\tau, \mu, \kappa, V)$ forces $\text{ICNIA}(\kappa, \mu)$ to hold at $\kappa = \mu^+$. Technically, the next theorem will become redundant after giving the proof of Lemma 3.6. However, the proof serves as a gentle introduction to these arguments.

Definition 2.16 [6]. Let Θ be a sufficiently large regular cardinal. We say that $M < H(\Theta)$ is *rich* or *rich with respect to κ* if the following hold:

- (1) $\kappa \in M$;
- (2) $\bar{\kappa} := M \cap \kappa \in \kappa$ and $\bar{\kappa} < \kappa$;
- (3) $\bar{\kappa}$ is an inaccessible cardinal;
- (4) The cardinality of M is $\bar{\kappa}$;
- (5) M is closed under $< \bar{\kappa}$ -sequences.

It is easy to show that:

Fact 2.17. *If κ is Mahlo and Θ is a sufficiently large regular cardinal, then for all $a \in [H(\Theta)]^{<\kappa}$, there is a model $M < K$ such that $a \subseteq M$ and M is rich with respect to κ .*

Theorem 2.18. *$\mathbb{M}^\oplus(\tau, \mu, \kappa, V)$ forces that there exist stationarily many $N \in [H(\kappa)]^{\leq \mu}$ such that N is internally club but not internally approachable.*

Proof. To aid in the legibility, we let $\mathbb{M}^\oplus := \mathbb{M}^\oplus(\tau, \mu, \kappa, V)$.

Let \dot{C} be an \mathbb{M}^\oplus -name for a club. Let \dot{F} be an \mathbb{M}^\oplus -name for a function $[H(\kappa)]^{<\omega} \rightarrow [H(\kappa)]^{<\kappa}$ such that the closure points of \dot{F} are contained in \dot{C} . Let Θ be a cardinal such that $\dot{F} \in H(\Theta)$ and let $M' < H(\Theta)$ be rich with respect to κ such that $\dot{F}, \mathbb{M}^\oplus, \tau, \mu, \kappa \in M', \mu \subseteq M'$. Let $\nu := M' \cap \kappa$.

Let G be \mathbb{M}^\oplus -generic and consider $M'[G]$: Since $\dot{F}^G \in M'[G]$, $M'[G] \cap H(\kappa)$ is closed under \dot{F}^G and thus $M'[G] \cap H(\kappa) \in \dot{C}^G$. Furthermore, because \mathbb{M}^\oplus is κ -c.c., $M'[G] \cap H(\kappa) = (M' \cap H(\kappa)^V)[G] =: M[G]$. We will show that $M[G]$ is as required.

Let $\pi : M \rightarrow N$ be the Mostowski-Collapse of M . Because \mathbb{M}^\oplus is κ -cc., $M[G] \cap V = M$ and thus π extends to $\pi : M[G] \rightarrow N[G']$, where $G' := \pi[G]$. Looking at the proof of Lemma 2.12, G' is also equal to the $\mathbb{M}^\oplus(\tau, \mu, \nu)$ -generic filter induced by G and there is an $\mathbb{M}^\oplus(G', \tau, \mu, \kappa \setminus \nu)$ -generic filter (over $V[G']$) G'' such that $G = G' * G''$.

Claim 2.19. $M[G]$ is internally club.

Proof. We first show that $N[G']$ is internally club. We have $N[G'] \subseteq V[G']$. Additionally, the reverse inclusion holds for many sets.

Subclaim 2.20. If $x \in [N[G']]^{<\mu} \cap V[G']$, $x \in N[G']$.

Proof. If $x \in [N[G']]^{<\mu} \cap V[G']$, x has been added by $\text{Add}(\tau, \nu)$. Let \dot{x} be an $\text{Add}^V(\tau, \nu)$ -name for x . By the μ -c.c. of $\text{Add}^V(\tau, \nu)$, we can assume that \dot{x} is a $<\mu$ -sized subset of N (since $\dot{x}(\alpha)$ is an element of $N[G']$ for every α). Then, $\dot{x} \in N$ and thus $\dot{x}^{G'} \in N[G']$. \square

Subclaim 2.21. $N[G']$ is internally club.

Proof. By the previous claim, $[N[G']]^{<\mu} \cap V[G'] = [N[G']]^{<\mu} \cap N[G']$. $\mathbb{M}^\oplus(\nu + 2)$ collapses ν by adding a continuous and cofinal sequence into $[\nu]^{<\mu} \cap V[\text{Add}^V(\tau, \nu)]$ by Proposition 2.13. This is isomorphic to $[N[G']]^{<\mu} \cap V[G']$ since $|N[G']| = |N| = \nu$. Hence, $\mathbb{M}^\oplus(\nu + 2)$ forces that we can write $N[G'] = \bigcup_{i < \mu} N_i$ where $N_i \in [N[G']]^{<\mu} \cap V[G'] = [N[G']]^{<\mu} \cap N[G']$ for every $i < \mu$. \square

Since π is an “internal” isomorphism of $M[G]$ and $N[G']$, M is also internally club: Write $N[G'] = \bigcup_{i < \mu} N_i$ such that $N_i \in [N[G']]^{<\mu} \cap N[G']$ for every i . Then, $M[G] = \bigcup_{i < \mu} \pi^{-1}[N_i]$ and $\pi^{-1}[N_i] = \pi^{-1}(N_i) \in [M[G]]^{<\mu} \cap M[G]$ for every i (since $\text{otp}(N_i) < \mu < \text{crit}(\pi^{-1})$). \square

Thus, we are finished after showing.

Claim 2.22. $M[G]$ is not internally approachable.

Proof. Again, we show the following first:

Subclaim 2.23. $N[G']$ is not internally approachable.

Proof. Assume toward a contradiction that $N[G'] = \bigcup_{i < \mu} N_i$ such that for each j , $(N_i)_{i < j} \in N[G']$. In particular, $(N_i)_{i < j} \in V[G']$. Because $G = G' * G''$ and G'' is generic for an ordering with the $<\mu$ -approximation property (2.15), $(N_i)_{i < \mu} \in V[G']$. However, this implies that $N[G']$ has size μ in $V[G']$, a contradiction as $\mathbb{M}^\oplus(\tau, \mu, \nu)$ is ν -c.c. and $|N[G']| = |N| = |\nu|$. \square

Now assume $M[G] = \bigcup_{i < \mu} M_i$ such that for each $j < \mu$, $(M_i)_{i < j} \in M[G]$. Then, $N[G'] = \bigcup_{i < \mu} \pi[N_i]$ and for each $j < \mu$, $(\pi[N_i])_{i < j} = (\pi(N_i))_{i < j} = \pi((N_i)_{i < j}) \in N[G']$, since $\pi(\mu) = \mu$, giving us a contradiction. \square

Thus, we have produced a set in \dot{C}^G that is internally club but not internally approachable. \square

3 | DISTINGUISHING INTERNALLY CLUB AND APPROACHABLE ON AN INFINITE INTERVAL

In this section, we apply the previous results to obtain the distinction between internally club and approachable on the interval $[\aleph_2, \aleph_\omega)$, thus obtaining our main theorem.

3.1 | Preservation of the distinction

First, we do some preliminary work by establishing some conditions under which $\text{ICNIA}(\Theta, \aleph_n)$ is preserved by sufficiently well-behaved forcings.

To obtain the model for Theorem 1.1, we will make use of a projection analysis, showing that, for a given n , the distinction holds in an outer model of the target model. With this intention, we introduce a slight strengthening of ICNIA that is more easily preserved downward.

Definition 3.1. Let $\text{ICNIA}^+(\Theta, \mu)$ be the statement that $\Theta \geq \mu^+$ and there exist stationarily many $N \in [H(\Theta)]^{\leq \mu}$ such that

- (1) N is internally club.
- (2) There is no sequence $(X_i)_{i < \mu}$ of elements of $[\Theta]^{< \mu}$ such that $\bigcup_{i < \mu} X_i = N \cap \Theta$ and $(X_i)_{i < j} \in N$ for all $j < \mu$.

We say that N is *not ordinal-internally approachable* if clause (2) holds.

We easily see that $\text{ICNIA}^+(\Theta, \mu)$ implies $\text{ICNIA}(\Theta, \mu)$: if N is internally approachable, simply intersect the approaching sequence with the class of ordinals.

Proposition 3.2. Assume that W is a forcing extension of V by a forcing notion \mathbb{P} that is $< \mu^+$ -distributive. If $\text{ICNIA}^+(\Theta, \mu)$ holds in W , $\text{ICNIA}^+(\Theta, \mu)$ holds in V .

Proof. In V , let C be club in $[H(\Theta)^V]^{\leq \mu}$. Then, in $W := V[G]$, C is club in $[H(\Theta)^V]^{\leq \mu}$ by the distributivity. Let Θ' be larger than Θ and at least so large that $\mathbb{P} \in H(\Theta')$. We have the following statement whose form connects it to notions of properness.

Claim 3.3. In $V[G]$, the set

$$D' := \{M \in [H(\Theta')^V]^\mu \mid M[G] \cap V = M\}$$

is club in $[H(\Theta')^V]^\mu$.

Proof. For closure, notice that

$$\left(\bigcup_{i < \mu} M_i \right) [G] \cap V = \left(\bigcup_{i < \mu} M_i [G] \right) \cap V = \bigcup_{i < \mu} (M_i [G] \cap V).$$

For unboundedness, let $M_0 \in [H(\Theta')^V]^\mu$ be arbitrary. Inductively define $M_{n+1} := M_n \cup (M_n [G] \cap V)$. Then,

$$\left(\bigcup_{n \in \omega} M_n \right) [G] \cap V = \bigcup_{n \in \omega} M_n$$

since, given some $\tau \in M_n$ with $\tau^G \in V$, $\tau^G \in M_n [G] \cap V = M_{n+1}$. □

Additionally, $C' := \{M \in [H(\Theta')^V]^\mu \mid M \cap H(\Theta)^V \in C\}$ is club in $[H(\Theta')^V]^\mu$. Thus

$$E' := \{M[G] \mid M \in D' \cap C'\}$$

is club in $[H(\Theta')^V[G]]^\mu$ that equals $[H(\Theta')^W]^\mu$ by the size of Θ' . This implies that the set

$$E := \{M[G] \cap H(\Theta)^W \mid M[G] \in E'\}$$

contains a club in $[H(\Theta)^W]^\mu$.

Thus, there exists $M \in D' \cap C'$ such that $M[G] \cap H(\Theta)^W$ is internally club but μ^+ is not ordinal internally approachable in $M[G] \cap H(\Theta)^W$. We aim to show that, in V , $M \cap H(\Theta)^V$ is internally club but μ^+ is not approachable in $M \cap H(\Theta)^V$.

Claim 3.4. $M \cap H^V(\Theta)$ is internally club in the model V .

Proof. We can write $M[G] \cap H^W(\Theta) = \bigcup_{i < \mu} M_i$, where the union is continuous and increasing and each M_i is in $[M[G] \cap H(\Theta)]^{<\mu} \cap M[G] \cap H(\Theta)$. Because $M \in D'$, $M = M[G] \cap V$, so

$$\begin{aligned} M \cap H^V(\Theta) &= (M[G] \cap V) \cap H^W(\Theta) = (M[G] \cap H^W(\Theta)) \cap V = \\ &= \bigcup_{i < \mu} M_i \cap V = \bigcup_{i < \mu} M_i \cap H^V(\Theta), \end{aligned}$$

using that $H^V(\Theta) = H(\Theta) \cap V$ as \mathbb{P} does not collapse cardinals. As $M[G] \cap V = M$, $M_i \cap H^V(\Theta') = M_i \cap H^V(\Theta) \in M[G]$ for every $i < \mu$. Additionally, $M_i \cap H^V(\Theta)$ is a subset of $H^V(\Theta)$ of size $< \mu$, so $M_i \cap H^V(\Theta) \in H^V(\Theta)$: $M_i \in V$ by distributivity and has hereditary size $< \Theta$. Again, as $M \in D$, $M_i \cap H^V(\Theta) \in M[G] \cap V = M$, so in summary $M_i \cap H^V(\Theta) \in M \cap H^V(\Theta)$. \square

Claim 3.5. $M \cap H^V(\Theta)$ is not ordinal-internally approachable in the model V .

Proof. Since $M[G] \cap V = M$, $(M[G] \cap H^W(\Theta)) \cap \Theta = (M \cap H^V(\Theta)) \cap \Theta$. Thus, if $M \cap H^V(\Theta)$ were ordinal-internally approachable in the model V , the same would be the case in the model W (witnessed by the same sequence), a contradiction. \square

Thus, we have produced an element of C that is as required. \square

3.2 | Proving the main theorem

Now we will set up the proof of Theorem 1.1. Let $(\kappa_n)_{n \in \omega}$ be a sequence of Mahlo cardinals. We force with the full support iteration $\mathbb{I} = \langle \mathbb{P}_n : n < \omega \rangle$ where $\mathbb{P}_0 = \mathbb{M}^\oplus(\omega, \omega_1, \kappa_0, V)$ and given \mathbb{P}_n we let

$$\mathbb{P}_{n+1} = \mathbb{P}_n * \dot{\mathbb{M}}^\oplus(\kappa_{n-1}, \kappa_n, \kappa_{n+1}, V[\mathbb{P}_{n-1}]),$$

where $\kappa_{-2} = \omega$, $\kappa_{-1} = \omega_1$, and $V[\mathbb{P}_{-1}] = V$ for simplicity. Observe that the iteration will turn κ_n into \aleph_{n+2} for all $0 \leq n < \omega$.

We start with a small improvement of Theorem 2.18.

Lemma 3.6. *Let $\tau < \mu < \kappa$ be cardinals such that $\tau^{<\tau} = \tau$, $\mu = \mu^{<\mu}$ and κ is Mahlo. If γ is any ordinal, $\mathbb{M}^\oplus(\tau, \mu, \kappa, W) \times \text{Add}(\mu, \gamma)$ forces $\text{ICNIA}^+(\Theta, \mu)$ for all regular $\Theta \geq \kappa$.*

Proof of Lemma 3.6. We modify the proof of Theorem 2.18.

Define $\mathbb{Q} := \mathbb{M}^\oplus(\tau, \mu, \kappa, W) \times \text{Add}(\mu, \gamma)$. We will abbreviate this product as $\mathbb{M} \times \mathbb{A}_{\text{big}}$. We will write the product that projects onto \mathbb{M} as $\mathbb{T} \times \mathbb{A}_{\text{small}}$.

Let \dot{C} be a \mathbb{Q} -name for a club in $[H(\Theta)]^{\leq \mu}$ and \dot{F} a name for the corresponding function. Let $\Theta' > \Theta$ be such that $H(\Theta')$ contains \dot{F} and choose a rich model $M \prec H(\Theta')$ with respect to κ of cardinality ν such that M contains $\dot{q}, \mathbb{Q}, \Theta$, and \dot{F} .

Let $\bar{\mathbb{M}} = \mathbb{M} \cap M = \pi_M(\mathbb{M}) = \mathbb{M}(\tau, \mu, \nu, W)$. Let $\bar{\mathbb{A}}_{\text{big}} = \mathbb{A}_{\text{big}} \cap M = \pi_M(\mathbb{M})$.

Now we will argue that we can choose the generics in a way that will suit us. Let $G' \times H'$ be a $\bar{\mathbb{M}} \times \bar{\mathbb{A}}_{\text{big}}$ -generic filter containing \dot{q} . We can find $G_1'' \times G_2''$, a product of filters that are $\mathbb{T} \times \mathbb{A}_{\text{small}}$ -generic over $V[G']$ such that \mathbb{T} induces a generic G_0'' for \mathbb{M}/G using Lemma 2.14. Now we let $G = G' * G''$. Since $\mathbb{A}_{\text{big}} = \pi_M(\mathbb{A}_{\text{big}}) \times \mathbb{A}'_{\text{big}}$ where \mathbb{A}'_{big} is a remainder, there is H'' such that $H = H' \times H''$ is \mathbb{A}_{big} -generic and $\pi_M(G \times H) = G' \times H'$. (This can be formulated in terms of j_M , the reverse of the Mostowski collapse π_M , and applying Silver's classical lifting criterion.)

We will argue that $(M \cap H(\Theta))[G][H]$ is internally club and $X := \dot{X}^{G \times H}$ is not internally approachable in $(M \cap H(\Theta))[G][H]$ in the model $V[G][H]$. Since $(M \cap H(\Theta))[G][H]$ is closed under $\dot{F}^{G \times H}$, this suffices. We will argue using N , the image $\pi_M : M \rightarrow N$.

Claim 3.7. $(M \cap H(\Theta))[G][H]$ is internally club.

Proof. This holds as in the proof of Theorem 2.18: The Mostowski-Collapse of $(M \cap H(\Theta))[G][H]$ is equal to $\pi(H(\Theta))[G'][H']$ that is closed under $< \nu$ -sequences in $V[G'][H']$. As before, $G \times H$ adds a club in $[\pi(H(\Theta))[G'][H']]^{< \mu}$ consisting of elements of $\pi(H(\Theta))[G'][H']$. \square

The slightly harder claim is as follows.

Claim 3.8. $(M \cap H(\Theta))[G][H]$ is not ordinal-internally approachable.

Proof. Assume toward a contradiction that there is a sequence $(X_i)_{i < \mu}$ of elements of $[\Theta]^{< \mu}$ such that $(X_i)_{i < j} \in (M \cap H(\Theta))[G][H]$ for every $j < \mu$ and $\bigcup_{i < \mu} X_i = (M \cap H(\Theta))[G][H] \cap \Theta = \nu$. It follows that, for every $j < \mu$, $\pi((X_i)_{i < j}) = (\pi[X_i])_{i < j} = (X_i)_{i < j} \in N[G'][H'] \subseteq V[G'][H']$. The pair $(V[G'][H], V[G][H])$ has the $< \mu$ -approximation property (see the remark after the proof): In $V[G']$, the poset $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ has the $< \mu$ -approximation property because its term ordering is μ -strategically closed and $\text{Add}(\tau, \kappa \setminus \nu)^W$ is μ -Knaster. This remains true in $V[G'][H]$ that is an extension by the μ -distributive forcing poset $\text{Add}(\mu, \gamma)^V$. Thus, $\mathbb{M}^\oplus(G, \tau, \mu, \kappa \setminus \nu, W)$ has the $< \mu$ -approximation property in $V[G'][H]$ and because $V[G][H]$ is an extension of $V[G'][H]$ using precisely that forcing, the statement follows.

In particular, $(X_i)_{i < j} \in V[G'][H]$ for any $j < \mu$. Therefore, the whole sequence $(X_i)_{i < \mu}$ is in $V[G'][H]$. However, this implies that $\Theta \geq \nu$ has size μ in $V[G'][H]$, a contradiction, as $G' \times H$ is generic for the ν -Knaster forcing $\mathbb{M}^\oplus(\tau, \mu, \nu, W) \times \text{Add}(\mu, \gamma)$. \square

Remark 3.9. It is clear that the pair $(V[G'][H'], V[G][H])$ does not have the $< \mu$ -approximation property because it contains a factor of $\text{Add}(\mu, \gamma)$ that certainly adds fresh sets with ordertype μ .

Again, we have produced $(M \cap H(\Theta))[G][H] \in \dot{C}^{G \times H}$ that is internally club but not internally approachable. This contradicts the choice of \dot{q} . \square

Now we can finish the proof of Theorem 1.1: Let $n \in \omega$ be arbitrary.

To obtain $\text{ICNIA}(\Theta, \kappa_{n-1})$, we will view the iteration as a factorization $\mathbb{P}_{\text{low}}^n * \dot{\mathbb{P}}_{\text{next}}^n * \dot{\mathbb{P}}_{\text{high}}^n$, where

- $\mathbb{P}_{\text{low}}^n := \mathbb{P}_{n-1}$,
- $\dot{\mathbb{P}}_{\text{next}}^n$ is a \mathbb{P}_{n-1} -name for

$$\begin{aligned} & \mathbb{M}^\oplus(\kappa_{n-2}, \kappa_{n-1}, \kappa_n, V[\mathbb{P}_{n-2}]) * \\ & \dot{\mathbb{M}}^\oplus(\kappa_{n-1}, \kappa_n, \kappa_{n+1}, V[\mathbb{P}_{n-1}]) * \\ & \dot{\mathbb{M}}^\oplus(\kappa_n, \kappa_{n+1}, \kappa_{n+2}, V[\mathbb{P}_n]), \end{aligned}$$

and

- $\dot{\mathbb{P}}_{\text{high}}^n$ is a $\mathbb{P}_{\text{low}}^n * \dot{\mathbb{P}}_{\text{next}}^n$ -name for

$$\langle \mathbb{M}^\oplus(\kappa_{k-2}, \kappa_{k-1}, \kappa_k, V[\mathbb{P}_{k-2}]) : n+3 \leq k < \omega \rangle.$$

We want to show that \mathbb{P} forces $\text{ICNIA}(\Theta, \kappa_{n-1})$. Let G_{low} be $\mathbb{P}_{\text{low}}^n$ -generic over V and work in $V[G_{\text{low}}]$. Because $|\mathbb{P}_{\text{low}}^n| < \kappa_n$, κ_n remains Mahlo in this model.

Now we need to perform a term-space argument. Here, we will use the notation in which $A(\mathbb{P}_1, \dot{\mathbb{P}}_2)$ is the term-space forcing in which $\dot{\mathbb{P}}_2$ is the underlying forcing and the ordering is taken with respect to what is forced by the empty condition of \mathbb{P}_1 (see [2, Section 22]).

By standard term-space arguments, $\dot{\mathbb{P}}_{\text{next}}^n * \dot{\mathbb{P}}_{\text{high}}^n$ is a projection of $\mathbb{P}_{\text{next}}^n \times A(\mathbb{P}_{\text{next}}^n, \dot{\mathbb{P}}_{\text{high}}^n)$. Since $\mathbb{P}_{\text{next}}^n$ forces $\dot{\mathbb{P}}_{\text{high}}^n$ to be $< \kappa_n$ -strategically closed (using similar arguments to [3]), $A(\mathbb{P}_{\text{next}}^n, \dot{\mathbb{P}}_{\text{high}}^n)$ is $< \kappa_n$ -strategically closed. Now we focus on $\mathbb{P}_{\text{next}}^n$. Writing $\mathbb{M}^\oplus(\tau, \mu, \kappa, W)$ as $\text{Add}(\tau, \kappa)^W * \mathbb{T}(\tau, \mu, \kappa)$, we have

$$\begin{aligned} \mathbb{P}_{\text{next}}^n &= (\mathbb{M}^\oplus(\kappa_{n-2}, \kappa_{n-1}, \kappa_n, V[\mathbb{P}_{n-2}]) \times \text{Add}(\kappa_{n-1}, \kappa_{n+1})) \\ & * (\mathbb{T}(\kappa_{n-1}, \kappa_n, \kappa_{n+1}) * \text{Add}(\kappa_n, \kappa_{n+2})^{V[\mathbb{P}_n]} * \mathbb{T}(\kappa_n, \kappa_{n+1}, \kappa_{n+2})). \end{aligned}$$

Let

$$\mathbb{P}_{\text{mid}}^n := \mathbb{M}^\oplus(\kappa_{n-2}, \kappa_{n-1}, \kappa_n, V[\mathbb{P}_{n-2}]) \times \text{Add}(\kappa_{n-1}, \kappa_{n+1})$$

and

$$\begin{aligned} \mathbb{T}_{\text{next}}^n &:= \mathbb{T}(\mathbb{M}^\oplus(\kappa_{n-1}, \kappa_n, \kappa_{n+1})) \times A(\mathbb{M}^\oplus(\kappa_{n-2}, \kappa_{n-1}, \kappa_n), \text{Add}(\kappa_n, \kappa_{n+2})^{V[\mathbb{P}_n]}) \\ & \times A(\mathbb{M}^\oplus(\kappa_{n-2}, \kappa_{n-1}, \kappa_n) * \mathbb{M}^\oplus(\kappa_{n-1}, \kappa_n, \kappa_{n+1}), \mathbb{T}(\mathbb{M}^\oplus(\kappa_n, \kappa_{n+1}, \kappa_{n+2}))), \end{aligned}$$

which is $< \kappa_n$ -strategically closed. Then, $\mathbb{P}_{\text{next}}^n$ is easily seen to be a projection of $\mathbb{P}_{\text{mid}}^n \times \mathbb{T}_{\text{next}}^n$.

So, in summary, $\mathbb{P}_{\text{next}}^n * \dot{\mathbb{P}}_{\text{high}}^n$ is a projection of $\mathbb{P}_{\text{mid}}^n \times \mathbb{T}_{\text{high}}^n$, where

$$\mathbb{T}_{\text{high}}^n := \mathbb{T}_{\text{next}}^n \times A(\mathbb{P}_{\text{next}}^n, \dot{\mathbb{P}}_{\text{high}}^n).$$

We can consider any extension by $\mathbb{P}_{\text{mid}}^n \times \mathbb{T}_{\text{high}}^n$ as an extension first by $\mathbb{T}_{\text{high}}^n$ and then by $\mathbb{P}_{\text{mid}}^n$. In such an extension, $\text{ICNIA}^+(\Theta, \kappa_{n-1})$ holds: $\mathbb{T}_{\text{high}}^n$ preserves the Mahloness of κ_n by its strategic closure and does not add any new conditions to $\mathbb{P}_{\text{mid}}^n$. Therefore, by Lemma 3.6, $\mathbb{P}_{\text{mid}}^n$ forces $\text{ICNIA}^+(\Theta, \kappa_{n-1})$. Furthermore, any $<\kappa_n$ -sequence added by $\mathbb{P}_{\text{mid}}^n \times \mathbb{T}_{\text{high}}^n$ has been added by $\mathbb{P}_{\text{mid}}^n$, so $\text{ICNIA}^+(\Theta, \kappa_{n-1})$ also holds in any extension by $\mathbb{P}_{\text{next}}^n * \mathbb{P}_{\text{high}}^n$ by Proposition 3.2.

Remark 3.10. The first author obtained a proof of Theorem 1.1 using a product rather than an iteration [8].

Here is a question related to the technical aspects of this paper.

Question 3.11. Suppose that \mathbb{P} is a ν^+ -closed forcing and $S \subseteq P_\nu(H(\Theta))$ is a stationary set of internally club sets. Is S stationary in an extension by \mathbb{P} ?

JOURNAL INFORMATION

The *Bulletin of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

ORCID

Hannes Jakob  <https://orcid.org/0009-0006-4563-1369>

Maxwell Levine  <https://orcid.org/0000-0001-7150-102X>

REFERENCES

1. U. Abraham, *Aronszajn trees on \aleph_2 and \aleph_3* , Ann. Pure Appl. Logic **24** (1983), no. 3, 213–230.
2. J. Cummings, *Iterated forcing and elementary embeddings*, Handbook of set theory, M. Foreman and A. Kanamori (eds.), Springer, Dordrecht, 2010, pp. 775–883.
3. J. Cummings and M. Foreman, *The tree property*, Adv. Math. **133** (1998), no. 1, 1–32.
4. M. Foreman and S. Todorcevic, *A new Löwenheim-Skolem theorem*, Trans. Amer. Math. Soc. **357** (2005), no. 5, 1693–1715.
5. S. Fuchino and A. O. M. Rodrigues, *Reflection principles, generic large cardinals, and the continuum problem*, Symposium on Advances in Mathematical Logic, Springer, 2018, pp. 1–25.
6. L. Harrington and S. Shelah, *Some exact equiconsistency results in set theory*, Notre Dame J. Form. Log. **26** (1985), no. 2, 178–188.
7. H. Jakob, *Disjoint stationary sequences on an interval of cardinals*, arXiv:2309.01986, 2023.
8. H. Jakob, *Variants of Mitchell forcing*, Ph.D. thesis, University of Freiburg, 2023–2025.
9. T. Jech, *Set Theory*, Springer Monographs in Mathematics, Springer, Berlin, the third millennium, revised and expanded edition, 2003.
10. J. Krueger, *Internally club and approachable*, Adv. Math. **213** (2007), no. 2, 734–740.
11. J. Krueger, *A general Mitchell style iteration*, MLQ Math. Log. Q. **54** (2008), no. 6, 641–651.
12. J. Krueger, *Some applications of mixed support iterations*, Ann. Pure Appl. Logic **158** (2009), no. 1–2, 40–57.
13. M. Levine, *On disjoint stationary sequences*, Pac. J. Math. **332** (2024), no. 1, 147–165. <https://doi.org/10.2140/pjm.2024.332.147>
14. S. Unger, *Fragility and indestructibility ii*, Ann. Pure Appl. Logic **166** (2015), no. 11, 1110–1122.
15. T. Usuba, *The approximation property and the chain condition*, RIMS Kokyuroku **1985** (2014), 130–134.
16. Y. Yoshinobu, *The*-variation of the Banach–Mazur game and forcing axioms*, Ann. Pure Appl. Logic **168** (2017), no. 6, 1335–1359.