

MA 556 PRESENTATION

ANKAN MUKHERJEE

IIT BOMBAY

November 3, 2021

Sectional Curvature

Definition of Sectional Curvature

Let M be a Riemann manifold. Let σ be a two dimensional subspace of $T_p M$. Let (v, w) be a basis of σ . Then the scalar curvature is defined as

$$K(v, w) = \frac{\langle R(v, w)w, v \rangle}{|v \wedge w|^2} \quad (1)$$

Sectional Curvature

Definition of Sectional Curvature

Let M be a Riemann manifold. Let σ be a two dimensional subspace of $T_p M$. Let (v, w) be a basis of σ . Then the scalar curvature is defined as

$$K(v, w) = \frac{\langle R(v, w)w, v \rangle}{|v \wedge w|^2} \quad (1)$$

Note that the above definition of the sectional curvature is independent of the choice of basis (v, w) and is thus a property of the plane σ alone. This will be proven in the slides to come.

Intuitive Version of Sectional Curvature

Here we will intuitively see why the sectional curvature takes the form as given in 1. Note that the intuitive version presented in these slides is a culmination of various sources on the internet presenting several viewpoints on the topic.

Intuitive Version of Sectional Curvature

Here we will intuitively see why the sectional curvature takes the form as given in 1. Note that the intuitive version presented in these slides is a culmination of various sources on the internet presenting several viewpoints on the topic. I will be using geodesics as the starting point for the intuition behind curvature. Note that the definition of geodesic is a curve on a surface along which the distance between two points is the shortest. This can also be interpreted as the path that parallel transport of a vector along itself takes.

Intuitive Version of Sectional Curvature

Here we will intuitively see why the sectional curvature takes the form as given in 1. Note that the intuitive version presented in these slides is a culmination of various sources on the internet presenting several viewpoints on the topic. I will be using geodesics as the starting point for the intuition behind curvature. Note that the definition of geodesic is a curve on a surface along which the distance between two points is the shortest. This can also be interpreted as the path that parallel transport of a vector along itself takes. We will use a vector field of the tangent vectors on the family of geodesics, denoted by V and the vector field orthogonal to it, denoted by S . Note that we can use the vectors in S to estimate the distance between geodesics at a given point.

Figure for Intuitive Version of Sectional Curvature




Figure: Geodesics on some arbitrary surface. Note that v indicates the tangent vector to the geodesic s is the orthogonal vector that measures the distance between two geodesics.

Getting into the Math behind Curvature

We will start with some basic assumptions. Let (M, g) be our Riemann Manifold.

- We choose ∇ to be the connection whose torsion is 0.
- We will let the lie brackets of the vector fields V and S vanish.

Getting into the Math behind Curvature

We will start with some basic assumptions. Let (M, g) be our Riemann Manifold.

- We choose ∇ to be the connection whose torsion is 0.
- We will let the lie brackets of the vector fields V and S vanish.

Recall that in multi-variable calculus, we often used the second derivative to check curvature. This works because the second derivative measures the change of change of function. Change of function can occur even without curvature since the change can be uniform (linear). We will apply an analogous concept here by replacing the double derivative with double covariant derivative of the vector field V .

Getting into the Math behind Curvature

We will start with some basic assumptions. Let (M, g) be our Riemann Manifold.

- We choose ∇ to be the connection whose torsion is 0.
- We will let the lie brackets of the vector fields V and S vanish.

Recall that in multi-variable calculus, we often used the second derivative to check curvature. This works because the second derivative measures the change of change of function. Change of function can occur even without curvature since the change can be uniform (linear). We will apply an analogous concept here by replacing the double derivative with double covariant derivative of the vector field V . Note that the geodesic equation is given as the vector field whose parallel transport along itself is zero (i.e., covariant derivative with respect to itself is zero). Thus, we have

$$\nabla_{\vec{V}}\vec{V} = 0 \tag{2}$$

Math behind Curvature

Continuing,

$$\nabla_{\vec{v}}\vec{v} = 0 \quad (3)$$

Taking covariant derivative with respect to \vec{s} on both sides,

$$\nabla_{\vec{s}}\nabla_{\vec{v}}\vec{v} = 0 \quad (4)$$

$$\Rightarrow \nabla_{\vec{s}}\nabla_{\vec{v}}\vec{v} - \nabla_{\vec{v}}\nabla_{\vec{s}}\vec{v} + \nabla_{\vec{v}}\nabla_{\vec{s}}\vec{v} = 0 \quad (5)$$

Using the definition of Riemann Curvature, the above equation reduces to

$$R(\vec{s}, \vec{v})\vec{v} + \nabla_{\vec{v}}\nabla_{\vec{v}}\vec{s} = 0 \quad (6)$$

More Figures




Figure: Case I: We have straight geodesics




Figure: Case II: We have converging geodesics (positive curvature)




Figure: Case III: We have diverging geodesics (negative curvature)

Figures Explained

Note that $\nabla_{\vec{v}}\nabla_{\vec{v}}\vec{s}$ indicates the rate of change of how fast the \vec{s} vector changes with respect to travelling on the geodesic. First we consider the sign. In case I, $\nabla_{\vec{v}}\nabla_{\vec{v}}\vec{s} = 0$ (since \vec{s} is uniform throughout the geodesic). In case II, we have \vec{s} decreasing in magnitude and this decrease decreases continuously. Thus, $\nabla_{\vec{v}}\nabla_{\vec{v}}\vec{s} = 0$ is opposite to \vec{s} . This yields $\langle \nabla_{\vec{v}}\nabla_{\vec{v}}\vec{s}, \vec{s} \rangle < 0$. Using equation 6, we get $\langle R(\vec{s}, \vec{v})\vec{v}, \vec{s} \rangle > 0$. Likewise, for case III $\langle R(\vec{s}, \vec{v})\vec{v}, \vec{s} \rangle < 0$. This is the intuition behind curvature.

Is This It?

We have intuitively concluded the form of curvature as $K(\vec{s}, \vec{v}) = \langle R(\vec{s}, \vec{v})\vec{v}, \vec{s} \rangle$, which holds true upto a constant. Note that curvature must be independent of the choice of basis. This can be done by suitable normalization.

Is This It?

We have intuitively concluded the form of curvature as $K(\vec{s}, \vec{v}) = \langle R(\vec{s}, \vec{v})\vec{v}, \vec{s} \rangle$, which holds true upto a constant. Note that curvature must be independent of the choice of basis. This can be done by suitable normalization. Now onwards, we will drop the $\vec{\cdot}$ symbol.

Consider a new set of basis $(\tilde{v}, \tilde{w}) = (av + bw, cv + dw)$ for $a, b, c, d \in \mathbb{R}$. The curvature should not change in this new basis. We will use the fact that

$$\|\tilde{v} \wedge \tilde{w}\| = \|av + bw \wedge cv + dw\| \quad (7)$$

$$= |ad - bc| \|v \wedge w\| \quad (8)$$

Normalization: Part I

Now,

$$R(\tilde{v}, \tilde{w}) = R(av + bw, cv + dw) \quad (9)$$

$$= R(av, cv) + R(av, dw) + R(bw, cv) + R(bw, dw) \quad (10)$$

$$= acR(v, v) + adR(v, w) + bcR(w, v) + bdR(w, w) \quad (11)$$

$$= (ad - bc)R(v, w) \quad (12)$$

where equations 10 and 11 use the multi linearity of tensors and equation 12 uses the fact that $R(v, w) = -R(w, v)$.

Normalization: Part II

Using $K(v, w) = \langle R(v, w)w, v \rangle$ from the previous slides,

$$K(\tilde{v}, \tilde{w}) = \langle R(\tilde{v}, \tilde{w})\tilde{w}, \tilde{v} \rangle \quad (13)$$

$$= (ad - bc) \langle R(v, w)\tilde{w}, \tilde{v} \rangle \quad (14)$$

$$= (ad - bc) \langle R(v, w)av + bw, cv + dw \rangle \quad (15)$$

$$= (ad - bc)^2 \langle R(v, w)v, w \rangle \quad (16)$$

where equation 14 follows from equation 12 and equation 16 follows from the axioms of inner product of vectors.

Normalization: Part II

Using $K(v, w) = \langle R(v, w)w, v \rangle$ from the previous slides,

$$K(\tilde{v}, \tilde{w}) = \langle R(\tilde{v}, \tilde{w})\tilde{w}, \tilde{v} \rangle \quad (13)$$

$$= (ad - bc) \langle R(v, w)\tilde{w}, \tilde{v} \rangle \quad (14)$$

$$= (ad - bc) \langle R(v, w)av + bw, cv + dw \rangle \quad (15)$$

$$= (ad - bc)^2 \langle R(v, w)v, w \rangle \quad (16)$$

where equation 14 follows from equation 12 and equation 16 follows from the axioms of inner product of vectors.

Equation 16 shows that we need to scale down the curvature by $(ad - bc)^2$ in order to normalize it.

Final Result

Thus, we can define curvature as

$$K(v, w) = \frac{\langle R(v, w)w, v \rangle}{\|v \wedge w\|^2} \quad (17)$$

We can show that the above quantity is indeed a tensor and invariant under a change of basis.

Problem 8 of Notes Chapter 5

Problem 8 of Notes Chapter 5

Show that the sectional curvature of the sphere at all points is 1.

Problem 8: Motivation

We will derive the sectional curvature of a spherical surface of radius R . Note that while the calculations are straight-forward using the formulae for Christoffel symbols and Riemann tensor, the calculations are rather cumbersome and non-trivial. We aim to provide a detailed calculation of the various parameters, most of which may be skipped over due to want of time. The slides are however there to stay and will contain the detailed derivations.

Problem 8: Motivation

We will derive the sectional curvature of a spherical surface of radius R . Note that while the calculations are straight-forward using the formulae for Christoffel symbols and Riemann tensor, the calculations are rather cumbersome and non-trivial. We aim to provide a detailed calculation of the various parameters, most of which may be skipped over due to want of time. The slides are however there to stay and will contain the detailed derivations. Our basic layout is as follows:



Figure: Flowchart of our calculations

Problem 8: Setting Up

Some assumptions that we will make:

- We will assume the metric is given (this is one of the defining parameters of the manifold).

Problem 8: Setting Up

Some assumptions that we will make:

- We will assume the metric is given (this is one of the defining parameters of the manifold).
- We will follow the convention given in Do Carmo's book.

Problem 8: Setting Up

Some assumptions that we will make:

- We will assume the metric is given (this is one of the defining parameters of the manifold).
- We will follow the convention given in Do Carmo's book.
- We will use the formula for Christoffel symbols from metric.

$$\Gamma_{\mu\nu}^{\lambda} = \frac{1}{2}g^{\lambda\alpha}(g_{\alpha\mu,\nu} + g_{\alpha\nu,\mu} - g_{\mu\nu,\alpha}) \quad (18)$$

where α is a dummy summation index.

Problem 8: Setting Up

Some assumptions that we will make:

- We will assume the metric is given (this is one of the defining parameters of the manifold).
- We will follow the convention given in Do Carmo's book.
- We will use the formula for Christoffel symbols from metric.

$$\Gamma_{\mu\nu}^{\lambda} = \frac{1}{2}g^{\lambda\alpha}(g_{\alpha\mu,\nu} + g_{\alpha\nu,\mu} - g_{\mu\nu,\alpha}) \quad (18)$$

where α is a dummy summation index.

- We will use the formula for Riemann curvature from Christoffel symbols (Lemma 5.5 of notes).

$$R_{\lambda\mu\nu\sigma} = g_{\lambda\alpha}(\partial_{\sigma}\Gamma_{\mu\nu}^{\alpha} - \partial_{\nu}\Gamma_{\mu\sigma}^{\alpha} + \Gamma_{\sigma\gamma}^{\alpha}\Gamma_{\mu\nu}^{\gamma} - \Gamma_{\nu\gamma}^{\alpha}\Gamma_{\mu\sigma}^{\gamma}) \quad (19)$$

where α and γ are dummy summation indices.

Problem 8: Specific Case of Sphere

Note that our independent basis is formed by (θ, ϕ) that span the surface of the sphere. The sphere is of radius R .

Problem 8: Specific Case of Sphere

Note that our independent basis is formed by (θ, ϕ) that span the surface of the sphere. The sphere is of radius R .

The distance metric is

$$g = \begin{pmatrix} R^2 & 0 \\ 0 & R^2 \sin^2 \theta \end{pmatrix} \quad (20)$$

i.e., $g_{\theta\theta} = R^2, g_{\theta\phi} = g_{\phi\theta} = 0, g_{\phi\phi} = R^2 \sin^2 \theta$.

Problem 8: Specific Case of Sphere

Note that our independent basis is formed by (θ, ϕ) that span the surface of the sphere. The sphere is of radius R .

The distance metric is

$$g = \begin{pmatrix} R^2 & 0 \\ 0 & R^2 \sin^2 \theta \end{pmatrix} \quad (20)$$

i.e., $g_{\theta\theta} = R^2, g_{\theta\phi} = g_{\phi\theta} = 0, g_{\phi\phi} = R^2 \sin^2 \theta$.

Calculating the inverse

$$g^{-1} = \begin{pmatrix} \frac{1}{R^2} & 0 \\ 0 & \frac{1}{R^2 \sin^2 \theta} \end{pmatrix} \quad (21)$$

i.e., $g^{\theta\theta} = \frac{1}{R^2}, g^{\theta\phi} = g^{\phi\theta} = 0, g^{\phi\phi} = \frac{1}{R^2 \sin^2 \theta}$.

Problem 8: Computing the Christoffel Symbols

We will use equation 18 to calculate the Christoffel symbols. We will also use the fact that the cross terms in the metric are zero.

$$\Gamma_{\theta\theta}^{\theta} = \frac{1}{2}g^{\theta\theta}(g_{\theta\theta,\theta} + g_{\theta\theta,\theta} - g_{\theta\theta,\theta}) = 0 \quad (22)$$

$$\Gamma_{\theta\phi}^{\theta} = \frac{1}{2}g^{\theta\theta}(g_{\theta\theta,\phi} + g_{\theta\phi,\theta} - g_{\theta\phi,\theta}) = 0 \quad (23)$$

$$\Gamma_{\phi\theta}^{\theta} = \frac{1}{2}g^{\theta\theta}(g_{\theta\phi,\theta} + g_{\theta\theta,\phi} - g_{\phi\theta,\theta}) = 0 \quad (24)$$

$$\Gamma_{\phi\phi}^{\theta} = \frac{1}{2}g^{\theta\theta}(g_{\theta\phi,\phi} + g_{\theta\phi,\phi} - g_{\phi\phi,\theta}) = -\frac{1}{2}g^{\theta\theta}g_{\phi\phi,\theta} = -\sin\theta\cos\theta \quad (25)$$

Problem 8: Continuing the Computation of the Christoffel Symbols

$$\Gamma_{\theta\theta}^{\phi} = \frac{1}{2}g^{\phi\phi}(g_{\phi\theta,\theta} + g_{\theta\phi,\theta} - g_{\theta\theta,\phi}) = 0 \quad (26)$$

$$\Gamma_{\theta\phi}^{\phi} = \frac{1}{2}g^{\phi\phi}(g_{\theta\phi,\phi} + g_{\phi\phi,\theta} - g_{\theta\phi,\phi}) = \frac{1}{2}g^{\phi\phi}g_{\phi\phi,\theta} = \frac{\cos\theta}{\sin\theta} \quad (27)$$

$$\Gamma_{\phi\theta}^{\phi} = \frac{1}{2}g^{\phi\phi}(g_{\phi\phi,\theta} + g_{\phi\theta,\phi} - g_{\phi\theta,\phi}) = \frac{1}{2}g^{\phi\phi}g_{\phi\phi,\theta} = \frac{\cos\theta}{\sin\theta} \quad (28)$$

$$\Gamma_{\phi\phi}^{\phi} = \frac{1}{2}g^{\phi\phi}(g_{\phi\phi,\phi} + g_{\phi\phi,\phi} - g_{\phi\phi,\phi}) = 0 \quad (29)$$

Problem 8: Riemann Curvature

We shall calculate only $R_{\theta\phi\theta\phi}$ for calculating sectional curvature. It is straight-forwards but tedious to show that only the permutations of this tensor are non zero. We will use equation 19.

$$R_{\theta\phi\theta\phi} = g_{\theta\theta}(\partial_{\theta}\Gamma_{\phi\phi}^{\theta} - \partial_{\phi}\Gamma_{\phi\theta}^{\theta} + \Gamma_{\theta\theta}^{\theta}\Gamma_{\phi\phi}^{\theta} + \Gamma_{\theta\phi}^{\theta}\Gamma_{\phi\phi}^{\phi} - \Gamma_{\phi\theta}^{\theta}\Gamma_{\phi\theta}^{\theta} - \Gamma_{\phi\phi}^{\theta}\Gamma_{\phi\theta}^{\phi}) \quad (30)$$

$$= R^2(\sin^2 \theta - \cos^2 \theta - 0 + 0 + 0 - 0 - (-\sin \theta \cos \theta)(\frac{\cos \theta}{\sin \theta})) \quad (31)$$

$$= R^2 \sin^2 \theta \quad (32)$$

Problem 8: Sectional Curvature

So far we have

$$R_{\theta\phi\theta\phi} = R^2 \sin^2 \theta \quad (33)$$

Also, we have the formula for sectional curvature

$$K(v, w) = \frac{\langle R(v, w)w, v \rangle}{|v \wedge w|^2} = \frac{R_{vwvw}}{\det(g)} \quad (34)$$

Hence, we get the sectional curvature of the sphere to be

$$K(\theta, \phi) = \frac{R_{\theta\phi\theta\phi}}{\det(g)} = \frac{R^2 \sin^2 \theta}{R^4 \sin^2 \theta} = \frac{1}{R^2} \quad (35)$$

Problem 8: QED

Thus, for a unit sphere, curvature is constant ($= 1$) at all points, irrespective of the values of θ , ϕ .

Problem 8: QED

Thus, for a unit sphere, curvature is constant ($= 1$) at all points, irrespective of the values of θ , ϕ .

Bonus

What surface has constant negative curvature (say, -1)?

Riemann Curvature from Sectional Curvature

Riemann Curvature from Sectional Curvature

We can deduce the Riemann curvature at a point in the manifold given the sectional curvature of the surface.

Motivation

We will define some basic assumptions that we will use in our proof.

- Define

$$R(x, y, z, w) = \langle R(x, y)z, w \rangle \quad (36)$$

- Skew symmetry

$$R(x, y, z, w) = -R(y, x, z, w) \quad (37)$$

$$R(x, y, z, w) = -R(x, y, w, z) \quad (38)$$

- Symmetry

$$R(x, y, z, w) = R(z, w, x, y) \quad (39)$$

- Bianchi Identity

$$\sum_{\pi(x,y,z)} R(x, y, z, w) = 0 \quad (40)$$

Setting Up

Now, define a polynomial

$$f(t) = R(x + tw, y + tz, y + tz, x + tw) \quad (42)$$

$$- t^2(R(x, z, z, x) + R(w, y, y, w)) \quad (43)$$

Since all terms are symmetric, $f(t)$ can be expressed in terms of the sectional curvature and norm of wedge products. Thus, the RHS of equation 42 is a polynomial in t whose coefficients are the sectional curvatures.

Setting Up

Now, define a polynomial

$$f(t) = R(x + tw, y + tz, y + tz, x + tw) \quad (42)$$

$$- t^2(R(x, z, z, x) + R(w, y, y, w)) \quad (43)$$

Since all terms are symmetric, $f(t)$ can be expressed in terms of the sectional curvature and norm of wedge products. Thus, the RHS of equation 42 is a polynomial in t whose coefficients are the sectional curvatures.

Consider the coefficient of t^2 in equation 42. Using the multilinearity property of R tensor, we can write it as

$$f''(0) = R(x, y, z, w) + R(x, z, y, w) + R(w, z, y, x) + R(w, y, z, x) \quad (44)$$

Some Simple Manipulations

Using equations 37, 38, 39, we obtain

$$f''(0) = 2R(x, y, z, w) - 2R(z, x, y, w) \quad (45)$$

Now, exchange x and y and define a new polynomial

$$g(t) = R(y + tw, x + tz, x + tz, y + tw) \quad (46)$$

$$- t^2(R(y, z, z, y) + R(w, x, x, w)) \quad (47)$$

Since all terms are symmetric, $g(t)$ can again be expressed in terms of the sectional curvature and norm of wedge products. Following a similar procedure, we obtain

$$g''(0) = -2R(x, y, z, w) + 2R(y, z, x, w) \quad (48)$$

Some Simple Manipulations

Using equations 37, 38, 39, we obtain

$$f''(0) = 2R(x, y, z, w) - 2R(z, x, y, w) \quad (45)$$

Now, exchange x and y and define a new polynomial

$$g(t) = R(y + tw, x + tz, x + tz, y + tw) \quad (46)$$

$$- t^2(R(y, z, z, y) + R(w, x, x, w)) \quad (47)$$

Since all terms are symmetric, $g(t)$ can again be expressed in terms of the sectional curvature and norm of wedge products. Following a similar procedure, we obtain

$$g''(0) = -2R(x, y, z, w) + 2R(y, z, x, w) \quad (48)$$

Now, 45-48 gives

$$f''(0) - g''(0) = 4R(x, y, z, w) - 2R(z, x, y, w) - 2R(y, z, x, w) \quad (49)$$

Completion of Proof

Using 40,

$$R(z, x, y, w) + R(y, z, x, w) = -R(x, y, z, w) \quad (50)$$

Substituting 50 in 49,

$$f''(0) - g''(0) = 6R(x, y, z, w) \quad (51)$$

$$\Rightarrow R(x, y, z, w) = \frac{f''(0) - g''(0)}{6} = \frac{(f - g)''(0)}{6} \quad (52)$$

Final Result

We need the polynomials f and g in terms of sectional curvature. Observe the form of f in 42. Using the formula for sectional curvature,

$$f(t) = K(x + tw, y + tz) \|(x + tw) \wedge (y + tz)\|^2 \quad (53)$$

$$- t^2(K(x, z) \|x \wedge z\|^2 + K(w, y) \|w \wedge y\|^2) \quad (54)$$

Likewise

$$g(t) = K(y + tw, x + tz) \|(y + tw) \wedge (x + tz)\|^2 \quad (55)$$

$$- t^2(K(y, z) \|y \wedge z\|^2 + K(w, x) \|w \wedge x\|^2) \quad (56)$$

Halloween Special: A Scary Expression

Complete evaluation of equations 51, 53, and 55 gives an explicit formula for $R(x, y, z, w)$.

Halloween Special: A Scary Expression

Complete evaluation of equations 51, 53, and 55 gives an explicit formula for $R(x, y, z, w)$.

$$\begin{aligned}
 6 \langle R(x, y)z, w \rangle = & K(x + w, y + z) \|(x + w) \wedge (y + z)\|^2 \\
 & - K(y + w, x + z) \|(y + w) \wedge (x + z)\|^2 \\
 & - K(x, y + z) \|x \wedge (y + z)\|^2 - K(y, x + w) \|y \wedge (x + w)\|^2 \\
 & - K(z, x + w) \|z \wedge (x + w)\|^2 - K(w, y + z) \|w \wedge (y + z)\|^2 \\
 & + K(x, y + w) \|x \wedge (y + w)\|^2 + K(y, z + w) \|y \wedge (z + w)\|^2 \\
 & + K(z, y + w) \|z \wedge (y + w)\|^2 + K(w, x + z) \|w \wedge (x + z)\|^2 \\
 & + K(x, z) \|x \wedge z\|^2 + K(y, w) \|y \wedge w\|^2 \\
 & - K(x, w) \|x \wedge w\|^2 - K(y, z) \|y \wedge z\|^2
 \end{aligned}$$

Quite scary, isn't it?

THANK YOU!

References

- ① Notes, MA 556, Prof. Swapneel Mahajan, IIT Bombay, Autumn 2021
- ② Manfredo Perdigão do Carmo, *Riemannian Geometry*, 2nd edition, Birkhäuser, 1992
- ③ Jeff Cheeger and David G. Ebin, *Comparison Theorems in Riemannian Geometry*, AMS Chelsea Publishing, vol. 365, 2008
- ④ https://www.maths.usyd.edu.au/PMH7/r/rg_emma_notes15.pdf
- ⑤ https://www.youtube.com/watch?v=ZhDNijOEwOY&ab_channel=eigenchris