

Quiz 6

Yuyan Zhou

April 10, 2020

1. Initialize θ
2. Repeat
3. E-step: Use \mathbf{X} and current θ to calculate $P(\mathbf{Z}|\mathbf{X}, \theta)$
4. M-step: Replace current θ by

$$\theta \leftarrow \arg \max_{\theta'} Q(\theta'|\theta) + \log P(\theta')$$

where $Q(\theta'|\theta) = E_{P(\mathbf{Z}|\mathbf{X}, \theta)}[\log P(\mathbf{X}, \mathbf{Z}|\theta')]$

5. Until convergence

Only M-step is changed, because in MAP, we have

$$\begin{aligned} & E_{P(\mathbf{Z}|\mathbf{X}, \theta)}[\log(P(\mathbf{X}, \mathbf{Z}|\theta')P(\theta'))] \\ &= E_{P(\mathbf{Z}|\mathbf{X}, \theta)}[\log P(\mathbf{X}, \mathbf{Z}|\theta')] + E_{P(\mathbf{Z}|\mathbf{X}, \theta)}[\log P(\theta')] \\ &= E_{P(\mathbf{Z}|\mathbf{X}, \theta)}[\log P(\mathbf{X}, \mathbf{Z}|\theta')] + \log P(\theta') \\ &= Q(\theta'|\theta) + \log P(\theta') \end{aligned}$$