

# JEE ADVANCED

ee24btech11004 - ANKIT JAINAR

## I. MCQs WITH ONE OR MORE THAN ONE CORRECT

- 1) The minimum value of expression  $\sin \alpha + \sin \beta + \sin \gamma$ , where  $(\alpha, \beta, \gamma)$  are real numbers satisfying  $(\alpha + \beta + \gamma) = \pi$  is (1995)
  - a) positive
  - b) 0
  - c) negative
  - d) -3
- 2) The number of values of  $x$  in the interval  $[0, 5\pi]$  satisfying equation  $3 \sin(x^2) - 7 \sin x + 2 = 0$  (1998-2 Marks)
  - a) 0
  - b) 5
  - c) 6
  - d) 10
- 3) Which of the following number(s) is/are rational? (1998-2 Marks)
  - a)  $\sin 15^\circ$
  - b)  $\cos 15^\circ$
  - c)  $\sin 15^\circ \cos 15^\circ$
  - d)  $\sin 15^\circ \cos 75^\circ$
- 4) For a positive integer  $n$ , let  $f_n(\theta) = (\tan \frac{\theta}{2})(1 + \sec \theta)(1 + \sec 2\theta)(1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$ . Then (1999 - 3Marks)
  - a)  $f_2(\frac{\pi}{16}) = 1$
  - b)  $f_3(\frac{\pi}{32}) = 1$
  - c)  $f_4(\frac{\pi}{64}) = 1$
  - d)  $f_5(\frac{\pi}{128}) = 1$
- 5) If  $\frac{\sin^4 x}{2} + \frac{\cos^4 x}{3} = \frac{1}{5}$ , Then (2009)
  - a)  $\tan^2 x = \frac{2}{3}$
  - b)  $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{1}{125}$
  - c)  $\tan^2 x = \frac{1}{3}$
  - d)  $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{2}{125}$
- 6) For  $0 < \theta < \frac{\pi}{2}$ , the solution(s) of  $\sum_{m=1}^6 \operatorname{cosec}(\theta + \frac{(m-1)\pi}{4}) \operatorname{cosec}(\theta + \frac{m\pi}{4}) = 4\sqrt{2}$  is(are) (2009)
  - a)  $\frac{\pi}{4}$
  - b)  $\frac{\pi}{6}$
  - c)  $\frac{\pi}{12}$
  - d)  $\frac{5\pi}{12}$
- 7) Let  $\theta, \varphi \in [0, 2\pi]$  be such that  $2 \cos(\theta(1 - \sin \varphi)) = \sin^2(\theta(\tan \frac{\theta}{2} + \cot \frac{\theta}{2})) \cos \varphi - 1, \tan(2\pi - \theta) > 0$  and  $-1 < \sin \theta < -\frac{\sqrt{3}}{2}$ , then  $\varphi$  cannot satisfy (2012)
  - a)  $0 < \varphi < \frac{\pi}{2}$
  - b)  $\frac{\pi}{2} < \varphi < \frac{4\pi}{3}$
  - c)  $\frac{4\pi}{3} < \varphi < \frac{3\pi}{2}$
  - d)  $\frac{3\pi}{2} < \varphi < 2\pi$
- 8) The number of points in  $(-\infty, \infty)$ , for which  $x - x \sin x - \cos x = 0$ , is (JEE Adv.2013)
  - a) 6
  - b) 4
  - c) 2
  - d) 0
- 9) Let  $f(x) = x \sin \pi x$ ,  $x > 0$ . Then for all natural numbers  $n$ ,  $f'(x)$  vanishes at (JEE Adv. 2013)
  - a) A unique point in the interval  $(n, n + \frac{1}{2})$
  - b) A unique point in the interval  $(n + \frac{1}{2}, n + 1)$
  - c) A unique point in the interval  $(n, n + 1)$
  - d) Two points in the interval  $(n, n + 1)$
- 10) Let  $\alpha$  and  $\beta$  be non-zero real numbers such that  $2(\cos \beta - \cos \alpha) + \cos \alpha \cos \beta = 1$ . Then which of the following is/are true? (JEE Adv.2017)
  - a)  $\tan(\frac{\alpha}{2}) + \sqrt{3} \tan(\frac{\beta}{2}) = 0$
  - b)  $\sqrt{3}(\tan \frac{\alpha}{2}) + \tan(\frac{\beta}{2}) = 0$
  - c)  $\tan(\frac{\alpha}{2}) - \tan(\frac{\beta}{2}) = 0$
  - d)  $\sqrt{3} \tan(\frac{\alpha}{2}) - \tan(\frac{\beta}{2}) = 0$

## II. SUBJECTIVE PROBLEMS

- 11) If  $\tan \alpha = \frac{m}{m+1}$  and  $\tan \beta = \frac{1}{2m+1}$ , find the possible values of  $(\alpha + \beta)$  (1978)
- 12) a) Draw the graph of  $y = \frac{1}{\sqrt{2}} (\sin x + \cos x)$  from  $x = -\frac{\pi}{2}$  to  $x = \frac{\pi}{2}$   
 b) If  $\cos(\alpha + \beta) = \frac{4}{5}, \sin(\alpha - \beta) = \frac{5}{13}$ , and  $\alpha, \beta$  lies between 0 and  $\frac{\pi}{4}$ , find  $\tan 2\alpha$  (1979)
- 13) Given  $\alpha + \beta - \gamma = \pi$ , prove that  $\sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma = 2 \sin \alpha \sin \beta \cos \gamma$  (1980)
- 14) Given  $A = \{x : \frac{\pi}{6} \leq x \leq \frac{\pi}{3}\}$  and  $f(x) = \cos x - x(1 + x)$ ; find  $f(A)$  (1980)

- 15) For all  $\theta$  in  $\left[0, \frac{\pi}{2}\right]$  show that,  $\cos(\sin \theta) \geq \sin(\cos \theta)$ .  
(1981-4 Marks)