

QnA's

1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

CRITICAL OBSERVATIONS

The demand of bike is less in the season 1 ie spring

The demand bike increased in the year 2019 wrt year 2018.

Month Jun - Sep is the period when bike demand is high. The Months Jan-Feb have lowest demand.

Bike demand is less in holidays .

2. Why is it important to use `drop_first=True` during dummy variable creation?

`drop_first=True` is important to use, as it helps in reducing the extra column created during dummy variable creation. Hence it reduces the correlations created among dummy variables.

3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

'temp' and 'atemp'

4. How did you validate the assumptions of Linear Regression after building the model on the training set?

Assumption of Regression Model :

- Linearity: The relationship between dependent and independent variables should be linear.
- Homoscedasticity: Constant variance of the errors should be maintained.

- Multivariate normality: Multiple Regression assumes that the residuals are normally distributed.
- Lack of Multicollinearity: It is assumed that there is little or no multicollinearity in the data.

5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

Top picks would be:

'temp'

'yr'

'weathersit'

1. Explain the linear regression algorithm in detail.

Linear Regression is a machine learning algorithm based on supervised learning. It performs a regression task. Regression models a target prediction value based on independent variables. It is mostly used for finding out the relationship between variables and forecasting. Different regression models differ based on – the kind of relationship between dependent and independent variables they are considering, and the number of independent variables getting used.

Linear regression performs the task to predict a dependent variable value (y) based on a given independent variable (x). So, this regression technique finds out a linear relationship between x (input) and y(output). Hence, the name is Linear Regression.

In the figure above, X (input) is the work experience and Y (output) is the salary of a person. The regression line is the best fit line for our model.

2. Explain the Anscombe's quartet in detail.

Anscombe's quartet tells us about the importance of visualizing data before applying various algorithms to build models. This suggests the data features must be plotted to see the distribution of the samples that can help you identify the various anomalies present in the data (outliers, diversity of the data, linear separability of the data, etc.). Moreover, the linear regression can only be considered a fit for the data with linear relationships and is incapable of handling any other kind of data set.

3. What is Pearson's R?

In statistics, the Pearson correlation coefficient (PCC), also referred to as Pearson's r , the Pearson product-moment correlation coefficient (PPMCC), or the bivariate correlation, is a measure of linear correlation between two sets of data. It is the covariance of two variables, divided by the product of their standard deviations; thus it is essentially a normalised measurement of the covariance, such that the result always has a value between -1 and 1 .

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

It is a step of data Pre-Processing which is applied to independent variables to normalize the data within a particular range. It also helps in speeding up the calculations in an algorithm.

Most of the times, collected data set contains features highly varying in magnitudes, units and range. If scaling is not done then algorithm only takes magnitude in account and not units hence incorrect modelling. To solve this issue, we have to do scaling to bring all the variables to the same level of magnitude.

It is important to note that scaling just affects the coefficients and none of the other parameters like t-statistic, F-statistic, p-values, R-squared, etc.

Normalization/Min-Max Scaling:

It brings all of the data in the range of 0 and 1. `sklearn.preprocessing.MinMaxScaler` helps to implement normalization in python.

Standardization Scaling:

Standardization replaces the values by their Z scores. It brings all of the data into a standard normal distribution which has mean (μ) zero and standard deviation one (σ).

`sklearn.preprocessing.scale` helps to implement standardization in python.

One disadvantage of normalization over standardization is that it loses some information in the data, especially about outliers.

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

An infinite value of VIF for a given independent variable indicates that it can be perfectly predicted by other variables in the model.

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

Quantile-Quantile (Q-Q) plot, is a graphical tool to help us assess if a set of data plausibly came from some theoretical distribution such as a Normal, exponential or Uniform distribution. Also, it helps to determine if two data sets come from populations with a common distribution.

This helps in a scenario of linear regression when we have training and test data set received separately and then we can confirm using Q-Q plot that both the data sets are from populations with same distributions.

Few advantages:

a) It can be used with sample sizes also

b) Many distributional aspects like shifts in location, shifts in scale, changes in symmetry, and the presence of outliers can all be detected from this plot.