

# SQL – Subqueries in WHERE Clause and Quantifiers

---

**CS 4750**  
**Database Systems**

[A. Silberschatz, H. F. Korth, S. Sudarshan, Database System Concepts, Ch.5.3]

# Subqueries in WHERE

Return a **single value** or a **relation** that can be compared to another value in a WHERE clause

Find the name(s) of the employee(s) who earn the highest salary for each job

HW\_emp(empno, ename, job, sal)

```
SELECT E1.ename
FROM HW_emp E1
WHERE E1.sal = (SELECT MAX(E2.sal)
                FROM HW_emp E2
                WHERE E1.job = E2.job);
```

**"Correlated" query**

[Refer to slide 14, <https://www.cs.virginia.edu/~up3f/cs4750/slides/4750meet16-17-SQL-subqueries.pdf> ]

# Subqueries in WHERE

Return a **single value** or a **relation** that can be compared to another value in a WHERE clause

Find the name(s) of the instructor(s) who earn the highest salary

Instructor (dept\_name, ID, name, salary)

```
SELECT name
FROM   instructor
WHERE  salary = (SELECT MAX(salary)
                  FROM instructor);
```

**“Uncorrelated”** query

# Subqueries in WHERE

Can be used to evaluate existential or universal quantifiers

At least one element

All elements

Any relational operators

Values of a column

ANY ( $\exists$ )

```
SELECT ... WHERE constant > ANY (subquery);
```

ALL ( $\forall$ )

```
SELECT ... WHERE constant > ALL (subquery);
```

IN

```
SELECT ... WHERE attribute IN (subquery);
```

NOT IN

```
SELECT ... WHERE attribute NOT IN (subquery);
```

EXISTS

```
SELECT ... WHERE EXISTS (subquery);
```

NOT EXISTS

```
SELECT ... WHERE NOT EXISTS (subquery);
```

Rows

# Existential Quantifier (EXISTS)

```
account (account_number, branch_name, balance)
borrower (customer_name, loan_number)
branch (branch_name, branch_city, assets)
depositor (customer_name, account_number)
loan (loan_number, branch_name, amount)
```

Find the names of the branches that have **some** customers who have both loan(s) and account(s) from the bank

```
SELECT T1.branch_name
FROM account T1
WHERE EXISTS
    (SELECT *
     FROM depositor D NATURAL JOIN borrower B
     WHERE D.account_number = T1.account_number);
```

tuple(s)

Correlated

# Existential Quantifier (IN)

```
account (account_number, branch_name, balance)
borrower (customer_name, loan_number)
branch (branch_name, branch_city, assets)
depositor (customer_name, account_number)
loan (loan_number, branch_name, amount)
```

Find the names of the branches that have **some** customers who have both loan(s) and account(s) from the bank

```
SELECT T1.branch_name
FROM account T1
WHERE T1.account_number IN
      (SELECT D.account_number
       FROM depositor D NATURAL JOIN borrower B);
```

attribute

attribute

Decorrelated

# IN and Set Membership

takes (ID, course\_id, sec\_id, semester, year, grade)

I need to know who takes the following courses:  
CS-101, CS-315, BIO-101

```
SELECT *  
FROM takes  
WHERE course_id IN  
      ('CS-101', 'CS-315', 'BIO-101');
```

attribute

Set of literal values



```
SELECT *  
FROM takes  
WHERE course_id='CS-101' OR course_id='CS-315' OR  
      course_id='BIO-101';
```

# Existential Quantifier

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase\_date, pid, custId, quantity, price)

Find the names of the customers who made **some** purchases that are > \$1000

```
SELECT DISTINCT T1.name
FROM Customer T1 NATURAL JOIN purchase T2
WHERE 1000 < T2.price
```



# Existential Quantifier (ANY)

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase date, pid, custId, quantity, price)

Find the names of the customers who made **some** purchases that are > \$1000

```
SELECT DISTINCT T1.name
```

```
FROM Customer T1
```

```
WHERE 1000 < ANY
```

```
(SELECT price  
FROM purchase T2
```

```
WHERE T1.custId = T2.custId)
```

constant

Correlated

Note: another way to solve (with subquery)

# Universal Quantifier

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase\_date, pid, custId, quantity, price)

Find the names of the customers who made purchases that are  
> \$1000 **only**



Find the names of the customers such that **all** their purchases  
are > \$1000



There does **not exist** any purchases the customer made where  
price <= \$1000

# Universal Quantifier (NOT IN)

Solution #1

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase\_date, pid, custId, quantity, price)

Find the names of the customers who made purchases that are  
> \$1000 **only**

Step 1: Find the customers who make **some** purchases  $\leq 1000$

```
SELECT DISTINCT T1.name
FROM Customer T1
WHERE T1.custId IN (SELECT T2.custId
                    FROM Purchase T2
                    WHERE T2.price <= 1000)
```

# Universal Quantifier (NOT IN)

Find the names of the customers who made purchases that are > \$1000 **only**

Step 1: Find the customers who make **some** purchases  $\leq 1000$

```
SELECT DISTINCT T1.name
FROM Customer T1
WHERE T1.custId IN (SELECT T2.custId
                    FROM Purchase T2
                    WHERE T2.price <= 1000)
```

Step 2: Find **all** customers who make purchase > 1000

```
SELECT DISTINCT T1.name
FROM Customer T1 NATURAL JOIN Purchase P
WHERE T1.custId NOT IN (SELECT T2.custId
                       FROM Purchase T2
                       WHERE T2.price <= 1000)
```

# Universal Quantifier (NOT EXISTS)

Solution #2

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase\_date, pid, custId, quantity, price)

Find the names of the customers who made purchases that are  
> \$1000 **only**

```
SELECT DISTINCT T1.name
FROM Customer T1 NATURAL JOIN purchase
WHERE NOT EXISTS (SELECT *
                  FROM purchase T2
                  WHERE T1.custId = T2.custId AND
                        T2.price <= 1000)
```

# Universal Quantifier (ALL)

Solution #3

Product(pid, name, cid)

Company(cid, cname, city)

Customer(custId, name, city)

Purchase(purchase\_date, pid, custId, quantity, price)

Find the names of the customers who made purchases that are  
> \$1000 **only**

```
SELECT DISTINCT T1.name
FROM Customer T1 NATURAL JOIN Purchase
WHERE 1000 < ALL (SELECT T2.price
                  FROM purchase T2
                  WHERE T1.custId = T2.custId)
```

# Let's Try 1: NOT EXISTS

```
account (account_number, branch_name, balance)
borrower (customer_name, loan_number)
branch (branch_name, branch_city, assets)
depositor (customer_name, account_number)
loan (loan_number, branch_name, amount)
```

Find the name(s) of the customer(s) who has a loan but does not have an account. Do not repeat the customer name.

```
SELECT DISTINCT B.customer_name
FROM borrower AS B
WHERE NOT EXISTS tuple(s)
    ( SELECT *
      FROM depositor AS D
      WHERE B.customer_name = D.customer_name );
```

Correlated

# Let's Try 2: NOT IN

account (account\_number, branch\_name, balance)  
borrower (customer\_name, loan\_number)  
branch (branch\_name, branch\_city, assets)  
depositor (customer\_name, account\_number)  
loan (loan\_number, branch\_name, amount)

Find the name(s) of the customer(s) who has a loan but does not have an account. Do not repeat the customer name.

```
SELECT DISTINCT B.customer_name  
FROM borrower AS B  
WHERE B.customer_name NOT IN  
(SELECT D.customer_name  
FROM depositor AS D);
```

attribute

attribute

Decorrelated

Note: another way to solve (equivalent to previous slide)



# Wrap-Up

- Subqueries in WHERE
- Internal interpretation of nested queries
- Many ways to express queries

## **Note:**

- Avoid nested queries – if aiming for speed
- Be careful of semantics of nested queries
  - Correlated vs. Uncorrelated

## **What's next?**

- Advanced SQL