

# **CQE, MSE, ITCZ, or all three? Theories for past and future monsoon rainfall changes**

Spencer A. Hill<sup>1,2</sup>

1. California Institute of Technology Division of Geological and Planetary Sciences, Pasadena, CA
2. UCLA Department of Earth, Planetary, and Space Sciences, Los Angeles, CA

*Degrees: B.S., Ph.D.*

*Email: shill@gps.caltech.edu*

Keywords:

1. monsoons
2. convective quasi-equilibrium
3. moist static energy
4. Intertropical convergence zone
5. paleoclimate
6. climate change

## Abstract

*Purpose of review:* Biases in simulations of past and present climate states and climate model disagreement even in sign of future monsoon rainfall changes bely limitations in our theoretical understanding.

*Recent findings:* The dominant theoretical paradigms for understanding monsoon rainfall --- convective-quasi equilibrium (CQE), moist static energy (MSE) budget, and monsoons as local Intertropical Convergence Zone (ITCZ) shifts --- all jettison the traditional “land-sea breeze” paradigm. Summer monsoon precipitation falls when the assumptions of CQE are most satisfied but those of the ITCZ shifts framework are least satisfied. Zonal asymmetries, changes in ITCZ width and strength, hydrology-vegetation-CO<sub>2</sub> coupling, and timescale-dependent responses complicate inferences of monsoon rainfall from paleoclimate proxy records. The MSE budget framework combined with careful simulation design can illuminate key mechanisms underlying monsoon responses to external forcings, presenting a path toward falsifying model projections.

*Summary:* Sustained, rapid progress in monsoon rainfall theory is urgently needed by society and seems attainable based on exciting recent advances.

## 1. Introduction

Earth's monsoons generate rainfall relied upon by billions of people and countless ecosystems. This makes the societal and ecological implications of any changes in the monsoon potentially enormous; the Indian finance minister once deemed the Indian monsoon the country's "real finance minister" (1). Yet general circulation models (GCMs) exhibit pronounced biases in monsoon rainfall compared to modern observations in simulations of present day (2) and compared to proxy records in simulations of past paleoclimate states (3,4). And projections of future rainfall change in monsoon regions differ starkly across GCMs, as model- and region-dependent circulation changes (5) amplify or counteract simple thermodynamic scaling-paced changes driving the zonal-mean tropical precipitation response (6). At a time when society demands actionable information to guide climate adaptation efforts, we are still unable to constrain even the sign of future precipitation change in any of Earth's monsoon regions, at least from GCM simulations alone.

Computing power that makes global, convection-resolving simulations routine --- and thus problematic cumulus parameterizations obsolete --- may one day greatly reduce model biases and future uncertainty (7), but such an era remains years to decades away (8). Some means of falsifying model projections is needed in the interim. Past warm states in Earth's history may provide useful lessons, but uncertainties inherent to proxies of continental hydroclimate encumber the paleoclimate record's ability to provide quantitative constraints, at least at our current levels of understanding. As such, I argue, improved theoretical understanding of monsoon rainfall responses to external forcings vital. Only with such knowledge can we improve GCMs' representations of monsoons through better-informed model development and, given the models' current state, use physically based arguments such as emergent observational constraints (9) to narrow probability distributions for the future beyond whatever emerges from the CMIP archive.

Fortunately, our theoretical understanding of region-mean monsoon precipitation has improved markedly in recent decades. Three primary frameworks have emerged: convective quasi-equilibrium (CQE) (10–12), the moist static energy (MSE) budget (13–16), and monsoons as local manifestations of energetically-forced shifts in the Intertropical Convergence Zone (ITCZ) (17). The ITCZ shifts framework has become particularly popular, both for the zonal-mean and increasingly as a basis for understanding zonally confined monsoons (17). Successfully adapting from the ITCZ to individual monsoons would be a tremendous accomplishment in terms of importance but also, I will argue, difficulty. Meanwhile, the pace of progress in recent years within the CQE and MSE frameworks has been no less brisk.

Here, I review recent advances in our theoretical understanding of forced monsoon rainfall changes based on these three frameworks. For brevity and biased by my own expertise, I largely restrict attention to the region- and wet-season-mean scales --- considering transients only insofar as they contribute to the mean state --- at the expense of promising recent work on the theory of synoptic-scale monsoon phenomena (18,19). For each of the CQE, MSE, and ITCZ frameworks (Sections 2–4, respectively), I present their foundational ideas, recent success stories, key limitations, and, for each limitation, encouraging results from recent studies and/or ideas for future work that could help resolve it. Section 3 also describes an example of the MSE budget being used toward developing an emergent observational constraint on future rainfall change in the West African Monsoon. I then discuss the implications of recent studies on inferring past monsoon rainfall changes from paleoclimate proxy records and the constraints that they can place on future anthropogenically driven changes (Section 5).

Finally, a parting discussion reflects on how to make the theoretical community's work more societally useful (Section 6).

Two prefatory points remain. First, none of the theories are consistent with the traditional view of monsoons as giant land-sea breezes fundamentally driven by land-sea contrasts in heat capacity and thus seasonal surface temperature evolution (20). Monsoon-like overturning circulations can emerge in aquaplanet simulations with uniform surface heat capacity (21). Surface thermal gradients usefully predict monsoon rainfall insofar as they are correlated with near-surface MSE (or, nearly equivalently, moist entropy or equivalent potential temperature). When relative humidity gradients are large, the dry and moist thermodynamic tracers decouple, and monsoon rainfall continues to track the moist tracer (12,22). Interannually, mean continental precipitation and surface temperature in each of Earth's major monsoon regions are *anti-correlated*: moisture delivered to the soil by rainfall promotes evapotranspiration at the expense of sensible heat flux, cooling the surface (23). [Over oceans, SST gradients are known to generate boundary layer convergence that drives convection (24–26), but the relevance of these circulations to continental monsoon rainfall is not obvious.]

Second, our understanding of the dynamics (rather than thermodynamics or energetics) of monsoons has also improved markedly in the past decades (21,27,28). While the classical angular momentum conserving model of the Hadley cells (29–31) and zonally confined monsoons (11) usefully characterizes the core of the cross-equatorial, zonal-mean overturning cells, otherwise the cells are strongly influenced by eddy stresses and do not homogenize angular momentum (28,32,33). During monsoon onset, the monsoonal cell rapidly transitions from the eddy-driven toward the classical regime, mediated by feedbacks from the mean flow and the rapid cessation of eddy stresses in the cell core (21,27,34,35). Outstanding challenges for this framework include that cells regularly fall in an intermediate regime, neither purely angular momentum conserving nor purely eddy-dominated (35), and that zonally oriented mass circulations can alter the leading-order balances within zonally confined monsoons (36). As regards monsoon rainfall, in what follows I show several useful ways that these free tropospheric zonal momentum considerations have been combined with the other frameworks.

## **2. CQE: convection-driven control of the column by the subcloud layer**

CQE is the state in which moist convection is sufficiently frequent and vigorous as to prohibit large values of convectively available potential energy (CAPE) in the time mean, by generating a nearly time-mean moist adiabatic stratification. To a good approximation, a column's thermodynamic structure is then set entirely by its near-surface MSE (strictly speaking, its subcloud moist entropy, an unimportant distinction in the present context) (10), a vertical truncation that has become fundamental to our understanding of myriad phenomena of the tropical atmosphere (37). Most notably for monsoons, it requires that (provided vertical shear is weak locally) the summer edge of a cross-equatorial monsoonal cell be co-located with a local maximum in near-surface MSE (38), placing the monsoon rainbelt at or just equatorward thereof. Observational and reanalysis data confirm that the core summer monsoon rainbelts coincide with local maxima in boundary layer equivalent potential temperature and upper tropospheric temperature (12). In a dry form, CQE also accounts for the shallow, dry circulations embedded (and acting to inhibit rainfall) within the deep, moist circulations in several monsoon regions, most notably the West African Monsoon (12,39,40).

Combining this general CQE assumption with the angular momentum conserving theory discussed above (10,11) leads to the existence of a “critical” near-surface MSE field, i.e. that in gradient wind balance with an angular momentum conserving circulation (provided that subsidence in the descending branch does not decouple the boundary layer and free troposphere). Insofar as the time-mean MSE field cannot strongly violate this criticality condition (11), the “supercriticality” of the radiative-convective equilibrium (RCE) state that would exist absent a large-scale overturning circulation then controls the strength of the overturning, as has been showed in idealized models (41,42).

Angular momentum is never fully homogenized throughout a monsoonal cell, however, although it is to good approximation along any of the cell’s individual streamlines within the free troposphere. This, combined with the core CQE assumption, implies that moist isentropes, angular momentum contours, and Hadley cell streamlines are all parallel in the free troposphere --- slantwise convective neutrality (43). Recently, Singh (44) used these arguments to derive a scaling for the latitude of the cross-equatorial solstitial cell’s edge in the summer hemisphere. It generalizes the aforementioned diagnostic (38) to cases with appreciable shear and is quantitatively accurate when applied to solstitial cross-equatorial Hadley cells in an idealized aquaplanet GCM. It will be interesting to see if it retains quantitative accuracy when applied to Earth’s monsoons and Hadley cells.

The scaling’s major drawback is that it is diagnostic, requiring knowledge of the equilibrated near-surface MSE field throughout the cell. The criticality arguments provide a prognostic scaling, requiring knowledge only of the RCE state (29,44). However, its assumption of true angular momentum homogenization causes it to generally be only qualitatively useful. It would be useful, therefore, to apply analytically (45) or empirically derived near-surface MSE profiles that more accurately reflect typical monsoon angular momentum distributions but that remain prognostic.

### **3. The MSE budget: leading-order balances to identify key processes limiting monsoon rainfall**

#### *3.1 The framework, its successes, and its limitations*

The thermodynamic equation of an atmospheric column can be usefully approximated as the column-integrated budget of MSE,  $c_p T + gz + Lq$ , the sum of sensible heat, gravitational potential energy, and latent heat. Conceptually, the budget states that the time tendency of internal energy depends on the balance between net energetic input via top-of-atmosphere radiative fluxes, surface radiative fluxes, and surface latent and sensible heat fluxes on the one hand, and the flux divergence of MSE by the atmospheric circulation on the other. Though not a coherent theory for the monsoons in itself (it is an expression of physical laws that must be satisfied whether or not a monsoon is present), its value stems from how the relative magnitudes of its terms succinctly characterize the nature of local circulations. For example, it has been used to identify key mechanisms underpinning the response of tropical precipitation to El Niño (46), global warming (46,47), and anthropogenic aerosols (48)

For monsoons, it has been used to identify key mechanisms limiting the poleward extent of monsoon rainfall (14–16). Specifically, monsoon extent can be limited by high surface albedo that prevents large net energetic input into the column needed to drive deep moist convection, or by “ventilation” --- the horizontal advection of low-MSE air into the monsoon region. In response to warming, monsoon rains are influenced by the famous “rich-get-richer” mechanism but also by the

“upped-ante” mechanism, wherein prevailing moisture and MSE gradients are enhanced by the mean warming, which combined with prevailing inflow into convective regions directed up these gradients leads to precipitation being suppressed at convective margins (47).


One of the key limitations of the MSE budget framework is a technical one: diagnosing the various budget terms from model output or reanalysis data *post hoc* typically results in residuals larger than many of the actual budget terms, precluding quantitative analyses. And existing adjustment methods that empirically correct for these residuals can be difficult to implement and require onerous amounts of data (7,49,50). As has been known for over a decade (51), this could be largely solved by models diagnosing the budget terms as they run, with column-integrated terms outputted as standard. Fortunately, recent model development (52) and intercomparison (53) efforts have taken this to heart, hopefully spurring others to do the same.

It is important to distinguish this framework from the energetic framework for ITCZ shifts to be discussed next. The MSE budget characterizes energetics only locally and is typically expressed in terms of MSE flux divergences. The ITCZ shifts framework considers planetary-scale energy flows and integrates the column MSE budget zonally and meridionally in order to examine meridional MSE fluxes. Similarly, though the previous section described the CQE framework primarily in terms of MSE, the CQE assumptions are not made in MSE budget diagnoses generally, and CQE was originally derived and is most properly expressed in terms of moist entropy.

### *3.2 Case study: rainfall change in the Sahel driven by mean SST warming*

This subsection details a recent success story of the MSE budget framework, wherein the framework is applied to targeted SST perturbation simulations in an AGCM in order to better understand anthropogenically forced rainfall changes in the Sahel, the northernmost region that receives appreciable rainfall from the West African Monsoon (7,54).

Though coupled atmosphere-ocean GCMs forced with likely changes to radiative forcing agents are ultimately our best tool for projecting the climate’s overall evolution in the coming century, inferring mechanisms of any regional changes in such simulations is hindered by the myriad, confounding processes triggered by those forcings. Prescribed SST simulations can disentangle those factors by imposing only one at a time. The key forcings include those due to the coupled model’s present-day SST biases (55,56), the direct influence of altered radiative forcing (57), changes in the global-mean SST value (6), and changes in the spatial pattern of SSTs (58–61). [This procedure can be extended even further to isolate e.g. the effect of plant physiological responses to CO<sub>2</sub> (62).]



*Figure 1: July-August-September precipitation response in (left) AM2.1, (center) AM3, and (right) HiRAM to the (top row) full, (middle row) mean, and (bottom row) spatial pattern components of the SST anomalies due to historical anthropogenic aerosol emissions. Precipitation is normalized by the control simulation Sahel region-mean JAS value in each model, with the region boundaries shown as the blue box.*

Using the MSE budget and uniform SST warming simulations, Hill et al. (7) demonstrate that uniform SST warming essentially triggers the upped-ante mechanism for the Sahel by enhancing the meridional MSE gradient spanning from the Sahara Desert to the Gulf of Guinea; acted upon by prevailing northerlies, this inhibits Sahelian precipitation. This mechanism also acts in reverse for global-mean cooling, yielding increased Sahel rainfall. But in either case, the magnitude of the anomalous dry air advection and its downstream impacts on Sahel precipitation are sensitive to the convective parameterization, and as such vary in strength across AGCMs (54).

The large amounts of sulfate aerosols emitted by the industrialized nations of the Northern Hemisphere in the 20<sup>th</sup> century acted to cool the globe overall and the northern hemisphere relative to the southern hemisphere. A previous study (63) computed SST anomalies driven by this historical aerosol anthropogenic aerosol forcing as simulated by the NOAA Geophysical Fluid Dynamics Laboratory (GFDL) AGCM AM2.1 (64,65) coupled with a slab ocean (66), and then re-ran the AGCM with those SST anomalies added to an observational climatological SST field (preventing contamination from any SST biases) in three different ways: without modification; with their tropical mean value (-1.1 K) applied at every ocean gridpoint; or with the tropical mean value subtracted from the full anomalies at each gridpoint (leaving the spatial pattern intact but with little mean change). Here I analyze the precipitation response in July-August-September over northern Africa in the original experiments and repeated in two additional GFDL AGCMs, AM3 (67) and HiRAM (68).

Figure 1 shows the precipitation responses in each simulation. In all three models, the spatial pattern component of relative Northern Hemisphere cooling and Southern Hemisphere warming acts to draw the Atlantic ITCZ and West African monsoon rainfall southward, resulting in widespread drying over the Sahel, consistent with expectations from the ITCZ shifts framework to be discussed in the next

section (69). Mean SST cooling, meanwhile, triggers the (reversed) upped-ante mechanism, but its differing strength across the models results in widely varying precipitation responses, from strong Sahel wettening in AM2.1 to little coherent response in HiRAM (54,70). As such, in response to the full aerosol SST anomalies, only in AM2.1 does the mean-driven wettening win out over the pattern-driven drying, so that Sahel precipitation actually increases appreciably, adjacent to strong drying in the Atlantic ITCZ.

In short, with knowledge of the Sahara-driven mechanism for Sahel wettening with mean cooling (which is ill-suited for interpretation via the energetic shifts framework), its differing magnitude across models, and the SST spatial pattern's drying influence (which, conversely, does naturally adhere to the shifts framework), the disparate model responses to the full aerosol SST pattern are readily understood.

The upped-ante mechanism of Sahel drying with mean SST warming has the potential to generate an emergent observational constraint, as follows. If the enhancement of dry Saharan air advection scales with the climatological convective depth in the Sahel (on the grounds that deeper convection will more effectively moisten and warm the region in the column integral relative to the Sahara than would shallow convection), then we should discount those models with convective depths much shallower or much deeper than the observational value. Initial work in this vein indicated that, indeed, AM2.1 has a very top-heavy ascent profile in the Sahel compared to observations and most other AGCMs (54). Although various complications precluded developing a definitive formal emergent constraint in this case (9), this seems like a useful template for additional future work toward developing emergent constraints for other regions and/or based on other physical mechanisms.

#### **4. ITCZ shifts: monsoons as regional manifestation of energetically driven rainbelt migrations**

The precipitation in the zonal-mean ITCZ is the hydrological imprint of the ascending branch of the zonal mean Hadley cells. In the simplest picture, the ITCZ is co-located with the zero crossing of the total atmospheric energy transport as well as the shared inner edge of the two Hadley cells. That ascending branch, and with it ITCZ rainfall, can migrate north and south in response to meridionally asymmetric forcings, such as when the ITCZ “follows the sun” over the annual cycle. In the annual mean, forcings with appreciable meridional structure can also move the ITCZ, as has been posited for, e.g., historical anthropogenic aerosol emissions and the growth and retreat of high-latitude ice sheets (71–75).

More recently, Bischoff and Schneider have developed a formalism that yields a quantitative prediction for ITCZ shifts and double ITCZ states based on anomalous cross-equatorial energy transport and the net energy input at the equator (76–79). Other authors have documented the often leading-order role played by processes the framework initially neglected, including Ekman-driven coupling with shallow ocean heat transports (80–82), interactions with the deep ocean meridional overturning circulation (83), radiative feedbacks from clouds (84) and water vapor (85) that attend ITCZ shifts, and the propensity for changes in ITCZ width and strength in addition to position (86). Recent work has even implicated the ITCZ's position (87) and width (88) as influencing the poleward extent of the Hadley cells' *descending* branch based on the notion that the cells nearly conserve angular momentum until they are truncated by the onset of baroclinic instability (89,90). Finally, the framework has recently been used to identify of several otherwise-unintuitive predictors of the



infamous “double ITCZ” bias (2), include biases in land surface albedo (91) and temperature (92), biases in cloud cover over the Southern Ocean (93), and values of tropical surface energy fluxes in prescribed SST AMIP simulations (94).

These ITCZ arguments bear on individual monsoons insofar as the local precipitation closely follows that of the zonal-mean, but this is not always the case. Pairs of GCM simulations exhibiting identical changes in cross-equatorial energy transport or in the position of the ITCZ can exhibit drastically different precipitation responses within different monsoon sectors (66). And the Atlantic ITCZ and adjacent West African Monsoon precipitation can respond with opposite sign to precessional orbital forcing (95,96). As such, the energetic shifts framework must incorporate zonally asymmetric processes. This has led to recent work incorporating zonally confined meridional MSE transports and precipitation shifts (79) and zonally oriented transports and precipitation shifts (97) into the framework.

As is often true of the original, zonal-mean version, results from these initial studies largely indicate qualitative, but not quantitative, utility of the theory. Accordingly, the remainder of this section considers several possible refinements. To be of value for monsoon rainfall, those couched in terms of the zonal-mean framework will ultimately need to be investigated as applied to zonally confined monsoons as well.

#### *4.1 Seasonality mismatch between ITCZ energetic framework accuracy and monsoon rainfall*

The ITCZ, energy flux equator, and the Hadley cells’ shared inner edge are all nearly co-located under equinoctial and annual-mean conditions, but they separate during solstitial seasons, with the core ITCZ precipitation moving equatorward of both the energy flux equator and the boundary between the cross-equatorial winter cell and the weak summer cell (98). Meanwhile, monsoon rain falls predominantly during local summer --- i.e. precisely when the assumptions underlying the ITCZ shifts framework are least satisfied! A pressing need, therefore, is to determine how useful the shifts framework is in a perturbative sense (which only requires that the various metrics be well correlated, rather than coincident) despite its relative inadequacy for the unperturbed solstitial state.

To address this, one could start by applying hemispherically asymmetric extratropical forcing as standard (99) in simulations with insolation set to either its annual mean, perpetual solstice, or the full annual cycle. In perpetual solstice simulations, though insolation and (on aquaplanets) MSE both maximize at the summer pole, dynamical constraints prevent the tropical rainbelts from extending much beyond their maximal extent with a real seasonal cycle and removes the complications stemming from the circulation’s time dependence (100,29,44). [Note in this case that any linearizations of the ITCZ position should be performed about the unperturbed energy flux equator, not the geographic equator (101).]

The ITCZ energetic framework is agnostic regarding the Hadley cells’ dynamical regime: whether the cells are angular momentum conserving or eddy-dominated is immaterial so long as the ITCZ, energy flux equator, and cell edge are well correlated. At the same time, Singh’s slantwise neutrality metric described above (44) and other recent work invoking angular momentum-related arguments places limits on the poleward edge of the Hadley cells in the summer hemisphere (29,44,100). Any such constraint on the cell’s overall poleward edge in turn places an upper bound on the poleward extent of the ITCZ (the ITCZ cannot occur poleward of where the Hadley cells

terminate). For example, relative warming of the summer hemisphere may shift the ITCZ less far poleward than identical but opposite signed forcing shifts the ITCZ equatorward. This could also be informed by recent work on the boundary layer momentum dynamics of ITCZ excursions (100,102,103).

### *3.2 On the context dependence and relationship to seasonality of the ITCZ-cross equatorial energy flux slope*

The regression of the zonal-mean ITCZ position on cross-equatorial atmospheric energy fluxes has been diagnosed in several contexts including the seasonal cycle to be approximately  $3^\circ \text{PW}^{-1}$  (77), leading to two claims that increasingly appear problematic (98). The first is that this  $3^\circ \text{PW}^{-1}$  slope is effectively invariant across climate states. However, large [i.e.  $O(1)$ ] differences between the slope during the seasonal cycle and in the annual mean response to external forcings have recently been reported in GCMs (104). Moreover, the aforementioned Bischoff and Schneider formalism --- arguably the closest we have to a closed theory for the ITCZ position --- suggests that this slope varies inversely with the equatorial net energetic input (76), which is known to vary on various timescales, e.g. interannually between El Nino and La Nina years (78).

The second claim is that the slope is fundamentally determined by the seasonal cycle: because the annual mean ITCZ position reflects the residual of large seasonal swings into either hemisphere, annual mean shifts also must follow this slope (98). However, as just noted, the annual cycle and annual mean slopes can in fact differ appreciably in GCMs. And ITCZ shifts in simulations perturbed with asymmetric forcings but lacking a seasonal cycle --- which arguably constitutes the bulk of theoretical studies on the topic to date --- do not obviously differ in this regard from those with seasonal cycles.

Nevertheless, a related claim --- that future annual-mean ITCZ shifts are bound to be weak (i.e.  $<1^\circ$ ) --- is indeed plausible (98). Barring some forcing or radiative feedback process that is far more hemispherically asymmetric than anticipated or is not currently represented, anomalous low-latitude energy fluxes will likely be much less than 1 PW in the coming century. Meanwhile, although the  $3^\circ \text{PW}^{-1}$  value can vary, there is no argument or simulation suggesting that it will increase in magnitude by several multiples. This may not usefully constrain shifts of any individual monsoon rainbelt, however, as the solstitial, zonally confined circulation may have a different slope than the annual- and zonal-mean.

### *3.3 Role of gross moist stability and eddy MSE fluxes in ITCZ shifts framework*

The coupling of movements in the ITCZ location, energy flux equator, and Hadley cell edge hinges on anomalous energy fluxes by the Hadley cells dominating over anomalous eddy energy fluxes, as well as on the anomalous Hadley cell energy fluxes in the vicinity of the ITCZ being effected predominantly by changes in the mass overturning strength rather than in the efficiency of energy transport per unit mass overturning, i.e. the Hadley cell gross moist stability (GMS) (63,105,106). Otherwise, the ITCZ and energy flux equator can separate, as can occur during monsoon retreat with the two features sitting in opposite hemispheres (107). And simulations of the mid-Holocene and other states exist in which, even absent the complications of ocean dynamical coupling, the zonal-mean ITCZ

responds to asymmetric forcing counter to expectations due either to changes in GMS (95,106) or eddy MSE fluxes (108).

Though GMS is often usefully approximated as the difference between upper- and lower-level MSE values (under the assumption that flow is concentrated into one narrow layer with each branch, each with a single representative MSE value), the actual GMS value can be sensitive to the vertical profile of the cell's meridional flow, for example decreasing the more the upper-branch flow is spread toward the mid-troposphere where MSE in the tropics is generally smallest (107). As such, a potentially useful approach could be a three layer conceptual model in which the upper branch flow is partitioned between upper- and mid-troposphere layers according to e.g. the meridional Laplacian of SSTs, motivated by the propensity for sharp SST gradients to generate shallow circulations embedded within the larger cell (107,24,25,109).

The roles of GMS and eddies are in fact related, with all else equal the eddy energy fluxes increasing in magnitude the smaller GMS is. Annual-mean GMS decreases in the deep tropics in an idealized aquaplanet GCM when the convective parameterization is made less active (110); it becomes negative when the parameterization is turned off entirely, meaning that the Hadley cells *converge* energy into the equatorial region, to be transported away by eddies. This modulation of GMS through the convection scheme could be exploited via experiments imposing the canonical asymmetric extratropical forcings (99) to mean states with GMS values from strongly positive to negative, to see how strongly the ITCZ response is affected.

Perhaps the simplest approach to incorporating transient eddy MSE fluxes into the energetic framework would be through a simple down-gradient diffusive approximation as is standard in other contexts and appears to be reasonably accurate in reanalysis data for various monsoon regions (39). Separately, one could apply the well-known phase speed-wavenumber spectral decomposition of equatorial variability (111) to meridional energy transports; knowing e.g. the relative contributions of equatorial Rossby waves vs. the MJO to energy export out of the deep tropics (especially during solstitial seasons) could lead to improved mechanistic understanding of the mean state, differences across models [consider that the widely used Frierson idealized GCM's equatorial wave spectra is, unlike observations, dominated by Kelvin waves (112)], and the response to forcing, the latter perhaps akin to how changes in the phase-speed spectra of extratropical waves has been linked to Hadley cell expansion with warming (113). For the monsoons, this could also be informed by the recent theoretical studies of monsoon intraseasonal waves mentioned in the Introduction (18,19).

## **5. On past monsoon rainfall behavior and interpretations of paleoclimate proxy records**

This section summarizes considers challenges in inferring monsoon rainfall changes from paleoclimate proxy records and using them to constrain future changes. Some stem directly from the monsoon theoretical frameworks; others are more general, but their relevance to monsoons, I contend, bears discussing.

### *5.1 Distinguishing among local monsoon behavior and zonal-mean change modes*

On the one hand, the aforementioned results of Roberts et al. (104) and Smyth et al. (95) imply that the rainfall signal recorded by any individual point proxy record in a monsoon region need not

reflect the zonal-mean ITCZ behavior. And, even if no zonal asymmetries existed, the propensity for the ITCZ to vary in width and strength in addition to position (86) complicates interpretations of proxy records. For example, simultaneous changes in ITCZ position, width, and intensity could largely cancel one another in terms of the local precipitation change recorded at a single point proxy record. Or, conversely, they could all be of the same sign, which if interpreted purely as a shift would lead to an overestimate of the shift magnitude. Separately, ITCZ shifts are often computed as changes in the centroid or some other median-like quantity of the zonal-mean precipitation distribution. This requires knowledge of precipitation over the entire tropics, a tall order for past paleoclimate states (104).

On the other hand, a recent simple model of zonal-mean tropical precipitation responses to orbital variations based on the ITCZ shifts framework is remarkably accurate at the obliquity and precessional timescales in reproducing rainfall proxy records at various individual tropical sites (114). It would be useful to compare this model to proxies for sites at the same latitude but different longitudes, with mutually shared signals likely reflecting zonally symmetric influences and discrepancies reflecting zonally asymmetric processes. Similarly, it could be compared to precipitation at different sites in long-running simulations in intermediate complexity GCMs, removing the uncertainties stemming from the proxy records themselves and thus enabling more direct analysis of the physical processes. Finally, the model could be readily modified in order to account for seasonal ITCZ width variations (which it currently neglects) and the ITCZ's tendency (at least in Earth's recent history) to be north of the equator because of northward ocean heat transport (83).

## *5.2 Interpreting abrupt shifts in point records given sharp spatial rainfall gradients*

Many published interpretations of paleoclimate monsoon rainfall records conclude that past rainfall changes have been abrupt, often with the implication that future changes will be as well. However, as has been pointed out, a locally abrupt change in precipitation could result from gradual changes in a monsoon rainbelt, given the latter's often sharp spatial gradients (115). And a recent study demonstrates that "tipping points" and other nonlinear responses of monsoon rainfall inferred from simple theoretical models are based on questionable assumptions and do not emerge in GCM simulations subject to a wide range of forcings (116).

A potentially useful approach would be to construct a toy model of paleoclimate proxy records, starting with e.g. a single point measurement that perfectly records annual mean precipitation subject to a local rainbelt of fixed shape and sinusoidal seasonal migrations. The complicating factors of noise, a spatially aggregated signal, sampling bias, multiple records, more complex annual cycles of the rainbelt position, and secular changes representing e.g. orbital forcing or zonal migrations could be progressively introduced, and the ability to detect the rapidity of changes in monsoon rainfall could be assessed. (This could also assist with the aforementioned need to disentangle changes in width, strength, and position.)

## *5.3 Coupling among hydrology, CO<sub>2</sub>, vegetation, temperature, and albedo on land*

Over ocean, the saturated surface makes standard bulk aerodynamic formulas adequate for computing evaporation and the sensible heat flux; over land, a complex interplay exists among the supply of moisture by precipitation, the soil's ability to absorb that water, the partitioning of the

resulting soil moisture into direct evaporation and into transpiration by plants, the vertical distribution of water within the soil, and the surface energy balance. Moreover, stomatal closure by plants (which acts to decrease transpiration) with increasing CO<sub>2</sub> can lead to divergent responses of soil moisture and vegetation growth under CO<sub>2</sub>-driven climate change (117). As summarized by a recent series of literature reviews (118–120), this can lead to offline diagnostics such as the aridity index, Palmer Drought Severity Index, and the Penman-Monteith computation of potential evapotranspiration falsely indicating drought when in fact photosynthesis and runoff are enhanced. Fortunately, simple time-mean precipitation tends to be a useful predictor of the more directly impacts-related fields of runoff and photosynthesis (117). Nevertheless, the monsoon community should recognize this potential for hydroclimatic indicators to diverge, particularly under large CO<sub>2</sub> changes such as glacial-interglacial variations and future anthropogenic warming.

A further complication comes from the coupling of vegetation cover with surface albedo. Plants tend to be darker than the soil they grow over, such that more insolation is absorbed the more plants grow. This is thought to have figured centrally in the Green Sahara paleoclimate state of the Holocene, wherein the West African Monsoon rainfall expanded far north into what is now the Sahara desert. In this view, orbital precession enhanced boreal summer insolation, pushing the monsoon rains slightly farther northward, bringing with them enhanced vegetation growth, increasing the column shortwave radiation, and thus further enhancing the monsoon (along with other, non-albedo-mediated positive feedbacks from soil hydrology) (3). However, GCMs with dynamic vegetation models typically strongly underestimate the northward expansion of vegetation cover and of West African Monsoon rainfall into the Sahara. As such, one potential means of progress would be to introduce simple vegetation-dependent surface albedo formulations in simpler models (121), including idealized GCMs and diffusive moist energy balance models that include simple parameterizations of moisture fluxes by meridional overturning cells (122).

#### *5.4 Interpretation of paleoclimate analogs given long equilibration times of deep ocean*

Past warm states in Earth's history such as the Pliocene have become increasingly of interest as potential “analogs” to future warming. However, some paleoclimate proxies record near-equilibrium conditions, which includes the deep ocean's response that takes millennia and need not even be temporally monotonic: surface warming initially slows the Atlantic Meridional Overturning Circulation (AMOC), but after multiple millennia the AMOC can ultimately be enhanced (123), and/or a Pacific Meridional Overturning Circulation (PMOC) generated (124). PMOC or no, the concurrent slow release of heat from the deep ocean increases polar amplification of the initial warming signal. Both the AMOC (125) and polar amplified warming (126) have been argued to influence West African Monsoon rainfall, and other monsoons may well be impacted also. Therefore, past climate states as recorded in proxies may have only limited relevance as analogs to the transient component of anthropogenic warming with respect to monsoon behavior (127).

## **6. Conclusions**

Recent work at the nexus of social science and climate science suggests that the standard probabilistic framework to future uncertainty may not provide much actionable information for stakeholders; this could be ameliorated via a “storylines” approach in which a discrete number plausible outcomes are analyzed in detail (128). By no means should all theorists be compelled to

engage directly with the public in this way. But by being more reflective about our motivations and about the kinds of research that most effectively addresses those motivations, we can ultimately make our work all the more impactful.

A theorist espousing the societal value of theory may arouse skepticism, but already the advances in understanding I have described are being used to improve CMIP-class GCMs: aforementioned predictors of the double ITCZ bias based on the ITCZ shifts framework have informed model development and tuning of the NOAA GFDL AM4 AGCM, leading to substantial reductions in the double ITCZ bias and Southern Ocean cloud albedo biases in AMIP simulations (129).

As such, we should be encouraged by the tremendous improvements in recent years in our theoretical understanding of monsoons and their responses to climate changes past and future, advances that can and should form the foundation for further creative inquiries in years to come.

### **Acknowledgments**

I thank Yi Ming, Simona Bordoni, Jonathan Mitchell, Isaac Held, Ming Zhao, Natalie Burls, Martin Singh, Dargan Frierson, Bill Boos, Sarah Kang, Alex Gonzalez, Sean Faulk, David Neelin, Tim Merlis, Ho-Hsuan Wei, and Jack Scheff for conversations over the past few years that have shaped my thinking on these topics. I also thank Michael Byrne for an insightful review of the manuscript. This work was supported by a Caltech Foster and Coco Stanback Postdoctoral Fellowship.

### **Conflict of interest statement**

On behalf of all authors, the corresponding author states that there is no conflict of interest.

### **Works cited**

1. ....Indo-Asian News Service. India cheers as monsoon arrives; hopes of better farm output raised. Hindustan Times [Internet]. 2010 May 31 [cited 2015 Dec 31]; Available from: <http://www.hindustantimes.com/india/india-cheers-as-monsoon-arrives-hopes-of-better-farm-output-raised/story-Og0hZJ0ULuibRu4y7CVFpO.html>
2. ....Adam O, Schneider T, Brient F, Bischoff T. Relation of the double-ITCZ bias to the atmospheric energy budget in climate models. *Geophys Res Lett*. 2016 Jul 28;43(14):2016GL069465.
3. ..Tierney JE, Pausata FSR, deMenocal PB. Rainfall regimes of the Green Sahara. *Science Advances*. 2017 Jan 1;3(1):e1601503.
4. ....Haywood AM, Dowsett HJ, Dolan AM, Rowley D, Abe-Ouchi A, Otto-Bliesner B, et al. The Pliocene Model Intercomparison Project (PlioMIP) Phase 2: scientific objectives and experimental design. *Clim Past*. 2016 Mar 16;12(3):663–75.
5. ....Shepherd TG. Atmospheric circulation as a source of uncertainty in climate change projections. *Nature Geosci*. 2014 Oct;7(10):703–8.
6. ....Held IM, Soden BJ. Robust Responses of the Hydrological Cycle to Global Warming. *J Climate*. 2006 Nov 1;19(21):5686–99.

7. ....Hill SA, Ming Y, Held IM, Zhao M. A Moist Static Energy Budget–Based Analysis of the Sahel Rainfall Response to Uniform Oceanic Warming. *J Climate*. 2017 Apr 11;30(15):5637–60. **Uses MSE budget to identify simple upped-ante-like mechanism of enhanced Saharan air advection that dries the Sahel under mean SST warming.**
8. ....Schneider T, Teixeira J, Bretherton CS, Brient F, Pressel KG, Schär C, et al. Climate goals and computing the future of clouds. *Nature Clim Change*. 2017 Jan;7(1):3–5.
9. ....Klein SA, Hall A. Emergent Constraints for Cloud Feedbacks. *Curr Clim Change Rep*. 2015 Dec 1;1(4):276–87.
10. ....Emanuel KA, David Neelin J, Bretherton CS. On large-scale circulations in convecting atmospheres. *QJR Meteorol Soc*. 1994 Jul 1;120(519):1111–43.
11. ....Emanuel KA. On Thermally Direct Circulations in Moist Atmospheres. *J Atmos Sci*. 1995 May 1;52(9):1529–34.
12. ...Nie J, Boos WR, Kuang Z. Observational Evaluation of a Convective Quasi-Equilibrium View of Monsoons. *Journal of Climate*. 2010 Aug;23(16):4416–28.
13. ....Neelin JD, Held IM. Modeling Tropical Convergence Based on the Moist Static Energy Budget. *Mon Wea Rev*. 1987 Jan 1;115(1):3–12.
14. ....Chou C, Neelin JD, Su H. Ocean-atmosphere-land feedbacks in an idealized monsoon. *QJR Meteorol Soc*. 2001 Jul 1;127(576):1869–91.
15. ....Chou C, Neelin JD. Mechanisms limiting the southward extent of the South American Summer Monsoon. *Geophys Res Lett*. 2001 Jun 15;28(12):2433–6.
16. ....Chou C, Neelin JD. Mechanisms Limiting the Northward Extent of the Northern Summer Monsoons over North America, Asia, and Africa. *J Climate*. 2003 Feb 1;16(3):406–25.
17. ...Biasutti M, Voigt A, Boos WR, Braconnot P, Hargreaves JC, Harrison SP, et al. Global energetics and local physics as drivers of past, present and future monsoons. *Nature Geoscience*. 2018 Jun;11(6):392–400. **Reviews the ITCZ energetic framework and other conceptual frameworks for monsoon precipitation, emphasizing potential of adapting the ITCZ shifts framework to zonally confined monsoons.**
18. ....Diaz M, Boos WR. Barotropic Growth of Monsoon Depressions. *Quarterly Journal of the Royal Meteorological Society* [Internet]. 2019 [cited 2019 Jan 2];0(ja). Available from: <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3467>
19. ....Adames ÁF, Ming Y. Interactions between Water Vapor and Potential Vorticity in Synoptic-Scale Monsoonal Disturbances: Moisture Vortex Instability. *J Atmos Sci*. 2018 Mar 26;75(6):2083–106.
20. ....Gadgil S. The monsoon system: Land–sea breeze or the ITCZ? *J Earth Syst Sci*. 2018 Jan 27;127(1):1.
21. .Bordoni S, Schneider T. Monsoons as eddy-mediated regime transitions of the tropical overturning circulation. *Nature Geosci*. 2008 Aug;1(8):515–9.
22. ....Zhou W, Xie S-P. A Hierarchy of Idealized Monsoons in an Intermediate GCM. *J Climate*. 2018 Sep 10;31(22):9021–36. **Derives analytical model for monsoon rainfall based on surface equivalent potential temperature incorporating ventilation by subtropical westerly winds and other important processes.**
23. ....Hurley JV, Boos WR. Interannual Variability of Monsoon Precipitation and Local Subcloud Equivalent Potential Temperature. *J Climate*. 2013 Aug 1;26(23):9507–27.
24. ....Lindzen RS, Nigam S. On the Role of Sea Surface Temperature Gradients in Forcing Low-Level Winds and Convergence in the Tropics. *J Atmos Sci*. 1987 Sep 1;44(17):2418–36.
25. Back LE, Bretherton CS. On the Relationship between SST Gradients, Boundary Layer Winds, and Convergence over the Tropical Oceans. *J Climate*. 2009 Aug 1;22(15):4182–96.

26. . Sobel AH. Simple models of ensemble-averaged tropical precipitation and surface wind, given the sea surface temperature. In: The Global Circulation of the Atmosphere. Princeton University Press; 2007. p. 219–51.
27. ....Schneider T, Bordoni S. Eddy-Mediated Regime Transitions in the Seasonal Cycle of a Hadley Circulation and Implications for Monsoon Dynamics. *J Atmos Sci*. 2008 Mar 1;65(3):915–34.
28. ....Walker CC, Schneider T. Eddy Influences on Hadley Circulations: Simulations with an Idealized GCM. *J Atmos Sci*. 2006 Dec 1;63(12):3333–50.
29. ..Hill SA, Bordoni S, Mitchell JL. Axisymmetric constraints on cross-equatorial Hadley cell extent. *Journal of the Atmospheric Sciences* [Internet]. 2019 [cited 2018 Oct 30];Accepted. Available from: <https://arxiv.org/abs/1810.11105>
30. ....Schneider EK. Axially Symmetric Steady-State Models of the Basic State for Instability and Climate Studies. Part II. Nonlinear Calculations. *J Atmos Sci*. 1977 Feb 1;34(2):280–96.
31. ....Held IM, Hou AY. Nonlinear Axially Symmetric Circulations in a Nearly Inviscid Atmosphere. *J Atmos Sci*. 1980 Mar 1;37(3):515–33.
32. ..Walker CC, Schneider T. Response of idealized Hadley circulations to seasonally varying heating. *Geophys Res Lett*. 2005 Mar 1;32(6):L06813.
33. ...Schneider T. The general circulation of the atmosphere. *Annu Rev Earth Planet Sci*. 2006;34:655–688.
34. ...Bordoni S, Schneider T. Regime Transitions of Steady and Time-Dependent Hadley Circulations: Comparison of Axisymmetric and Eddy-Permitting Simulations. *J Atmos Sci*. 2010 May 1;67(5):1643–54.
35. ..Geen R, Lambert FH, Vallis GK. Regime Change Behavior during Asian Monsoon Onset. *Journal of Climate*. 2018 Apr;31(8):3327–48.
36. ....Zhai J, Boos W. Regime Transitions of Cross-Equatorial Hadley Circulations with Zonally Asymmetric Thermal Forcings. *J Atmos Sci*. 2015 Jul 23;72(10):3800–18.
37. ....Emanuel K. Inferences from Simple Models of Slow, Convectively Coupled Processes. *J Atmos Sci*. 2019 Jan;76(1):195–208.
38. Privé NC, Plumb RA. Monsoon Dynamics with Interactive Forcing. Part I: Axisymmetric Studies. *J Atmos Sci*. 2007 May 1;64(5):1417–30.
39. ....Zhai J, Boos WR. The drying tendency of shallow meridional circulations in monsoons. *QJR Meteorol Soc*. 2017 Oct 1;143(708):2655–64.
40. ....Shekhar R, Boos WR. Weakening and Shifting of the Saharan Shallow Meridional Circulation during Wet Years of the West African Monsoon. *J Climate*. 2017 Jun 22;30(18):7399–422.
41. ....Singh MS, Kuang Z. Exploring the Role of Eddy Momentum Fluxes in Determining the Characteristics of the Equinoctial Hadley Circulation: Fixed-SST Simulations. *J Atmos Sci*. 2016 Mar 25;73(6):2427–44.
42. ..Singh MS, Kuang Z, Tian Y. Eddy Influences on the Strength of the Hadley Circulation: Dynamic and Thermodynamic Perspectives. *J Atmos Sci*. 2017 Feb;74(2):467–86.
43. .Emanuel KA. The Lagrangian Parcel Dynamics of Moist Symmetric Instability. *J Atmos Sci*. 1983 Oct 1;40(10):2368–76.
44. ..Singh MS. Limits on the extent of the solstitial Hadley cell: the role of planetary rotation. *Journal of the Atmospheric Sciences*. 2019;Submitted. **Uses slantwise convective neutrality to generalize the Prive and Plumb 2007 diagnostic for the monsoon rainbelt position to cases with nonzero vertical zonal wind shear.**
45. ..Adam O, Paldor N. Global Circulation in an Axially Symmetric Shallow-Water Model, Forced by Off-Equatorial Differential Heating. *J Atmos Sci*. 2010 Apr 1;67(4):1275–86.



46. Neelin JD, Chou C, Su H. Tropical drought regions in global warming and El Niño teleconnections. *Geophys Res Lett*. 2003 Dec 1;30(24):2275.
47. .Chou C, Neelin JD. Mechanisms of Global Warming Impacts on Regional Tropical Precipitation. *J Climate*. 2004 Jul 1;17(13):2688–701.
48. Chou C, Neelin JD, Lohmann U, Feichter J. Local and Remote Impacts of Aerosol Climate Forcing on Tropical Precipitation. *Journal of climate* [Internet]. 2005 [cited 2014 Apr 30];18(22). Available from: <http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=08948755&AN=19883071&h=uNI9092WFmofs5dY6nUpKqBCsGJU7xMZGFcPjThRJzihU5UQloJF4KIKg52KQsi9m65d9hKrM9pXha9tgO9BGw%3D%3D&crl=c>
49. ....Trenberth KE. Climate Diagnostics from Global Analyses: Conservation of Mass in ECMWF Analyses. *J Climate*. 1991 Jul 1;4(7):707–22.
50. ....Peters ME, Kuang Z, Walker CC. Analysis of Atmospheric Energy Transport in ERA-40 and Implications for Simple Models of the Mean Tropical Circulation. *J Climate*. 2008 Oct 1;21(20):5229–41.
51. ..Neelin JD. Moist dynamics of tropical convection zones in monsoons, teleconnections, and global warming. In: *The Global Circulation of the Atmosphere* [Internet]. Princeton University Press; 2007 [cited 2014 Apr 30]. p. 267–301. Available from: <http://www.atmos.ucla.edu/~csi/REF/pdfs/gencircrev.pdf>
52. ....Zhao M. An Analysis of Global Climate Model Simulated Madden-Julian Oscillation with Fully Closed Moist Static Energy Budget. In *AMS*; 2018 [cited 2019 Mar 11]. Available from: <https://ams.confex.com/ams/33HURRICANE/webprogram/Paper338683.html>
53. ....Wing AA, Reed KA, Satoh M, Stevens B, Bony S, Ohno T. Radiative–convective equilibrium model intercomparison project. *Geosci Model Dev*. 2018 Mar 2;11(2):793–813.
54. ....Hill SA, Ming Y, Zhao M. Robust Responses of the Sahelian Hydrological Cycle to Global Warming. *J Climate*. 2018 Dec 15;31(24):9793–814. **Shows that enhanced Saharan air advection into the Sahel is robust across AGCMs under uniform SST warming and attempts to generate an emergent observational constraint based on this mechanism.**
55. ....He J, Soden BJ. The impact of SST biases on projections of anthropogenic climate change: A greater role for atmosphere-only models? *Geophys Res Lett*. 2016 Jul 28;43(14):2016GL069803.
56. ..Pascale S, Boos WR, Bordoni S, Delworth TL, Kapnick SB, Murakami H, et al. Weakening of the North American monsoon with global warming. *Nature Climate Change*. 2017 Nov;7(11):806–12.
57. ....Bony S, Bellon G, Klocke D, Sherwood S, Fermepin S, Denvil S. Robust direct effect of carbon dioxide on tropical circulation and regional precipitation. *Nature Geosci*. 2013 Jun;6(6):447–51.
58. ...Ma J, Xie S-P. Regional Patterns of Sea Surface Temperature Change: A Source of Uncertainty in Future Projections of Precipitation and Atmospheric Circulation. *J Climate*. 2012 Sep 17;26(8):2482–501.
59. Chadwick R, Boutle I, Martin G. Spatial Patterns of Precipitation Change in CMIP5: Why the Rich Do Not Get Richer in the Tropics. *J Climate*. 2013 Jun 1;26(11):3803–22.
60. Chadwick R, Good P, Andrews T, Martin G. Surface warming patterns drive tropical rainfall pattern responses to CO2 forcing on all timescales. *Geophys Res Lett*. 2014 Jan 28;41(2):610–5.
61. ....Chadwick R. Which Aspects of CO2 Forcing and SST Warming Cause Most Uncertainty in Projections of Tropical Rainfall Change over Land and Ocean? *J Climate*. 2016 Feb 3;29(7):2493–509.
62. ....Chadwick R, Douville H, Skinner CB. Timeslice experiments for understanding regional climate projections: applications to the tropical hydrological cycle and European winter circulation. *Clim Dyn*. 2017 Nov 1;49(9–10):3011–29.

63. ....Hill SA, Ming Y, Held IM. Mechanisms of Forced Tropical Meridional Energy Flux Change. *J Climate*. 2015;28(5):1725–42.
64. ....GFDL Atmospheric Model Development Team. The New GFDL Global Atmosphere and Land Model AM2–LM2: Evaluation with Prescribed SST Simulations. *J Climate*. 2004 Dec 1;17(24):4641–73.
65. .Delworth TL, Broccoli AJ, Rosati A, Stouffer RJ, Balaji V, Beesley JA, et al. GFDL’s CM2 Global Coupled Climate Models. Part I: Formulation and Simulation Characteristics. *J Climate*. 2006 Mar 1;19(5):643–74.
66. ....Ming Y, Ramaswamy V. Nonlinear Climate and Hydrological Responses to Aerosol Effects. *J Climate*. 2009 Mar 1;22(6):1329–39.
67. ....Donner LJ, Wyman BL, Hemler RS, Horowitz LW, Ming Y, Zhao M, et al. The Dynamical Core, Physical Parameterizations, and Basic Simulation Characteristics of the Atmospheric Component AM3 of the GFDL Global Coupled Model CM3. *J Climate*. 2011 Mar 2;24(13):3484–519.
68. ..Zhao M, Held IM, Lin S-J, Vecchi GA. Simulations of Global Hurricane Climatology, Interannual Variability, and Response to Global Warming Using a 50-km Resolution GCM. *J Climate*. 2009 Dec 1;22(24):6653–78.
69. ....Hwang Y-T, Frierson DMW, Kang SM. Anthropogenic sulfate aerosol and the southward shift of tropical precipitation in the late 20th century. *Geophys Res Lett*. 2013 Jun 16;40(11):2845–50.
70. Held IM, Delworth TL, Lu J, Findell KL, Knutson TR. Simulation of Sahel drought in the 20th and 21st centuries. *PNAS*. 2005 Dec 13;102(50):17891–6.
71. ....Kraus EB. The Seasonal Excursion of the Intertropical Convergence Zone. *Mon Wea Rev*. 1977 Aug 1;105(8):1052–5.
72. Kraus EB. Subtropical Droughts and Cross-Equatorial Energy Transports. *Mon Wea Rev*. 1977 Aug 1;105(8):1009–18.
73. ....Broccoli AJ, Dahl KA, Stouffer RJ. Response of the ITCZ to Northern Hemisphere cooling. *Geophys Res Lett*. 2006 Jan 16;33(1):L01702.
74. Kang SM, Frierson DMW, Held IM. The Tropical Response to Extratropical Thermal Forcing in an Idealized GCM: The Importance of Radiative Feedbacks and Convective Parameterization. *J Atmos Sci*. 2009 Sep 1;66(9):2812–27.
75. Schneider T, Bischoff T, Haug GH. Migrations and dynamics of the intertropical convergence zone. *Nature*. 2014 Sep 4;513(7516):45–53.
76. ....Bischoff T, Schneider T. Energetic Constraints on the Position of the Intertropical Convergence Zone. *J Climate*. 2014 Jul 1;27(13):4937–51.
77. ....Bischoff T, Schneider T. The Equatorial Energy Balance, ITCZ Position, and Double-ITCZ Bifurcations. *J Climate*. 2016 Apr 15;29(8):2997–3013.
78. .Adam O, Bischoff T, Schneider T. Seasonal and Interannual Variations of the Energy Flux Equator and ITCZ. Part I: Zonally Averaged ITCZ Position. *J Climate*. 2016 Feb 19;29(9):3219–30.
79. .Adam O, Bischoff T, Schneider T. Seasonal and Interannual Variations of the Energy Flux Equator and ITCZ. Part II: Zonally Varying Shifts of the ITCZ. *J Climate*. 2016 Jul 19;29(20):7281–93.
80. ....Green B, Marshall J. Coupling of Trade Winds with Ocean Circulation Damps ITCZ Shifts. *J Climate*. 2017 Mar 8;30(12):4395–411.
81. ..Schneider T. Feedback of Atmosphere-Ocean Coupling on Shifts of the Intertropical Convergence Zone. *Geophys Res Lett*. 2017 Nov 28;44(22):2017GL075817.
82. ....Kang SM, Shin Y, Xie S-P. Extratropical forcing and tropical rainfall distribution: energetics framework and ocean Ekman advection. *npj Climate and Atmospheric Science*. 2018 Jan 9;1(1):2.

83. ....Frierson DMW, Hwang Y-T, Fučkar NS, Seager R, Kang SM, Donohoe A, et al. Contribution of ocean overturning circulation to tropical rainfall peak in the Northern Hemisphere. *Nature Geoscience*. 2013 Oct 20;6(11):940–4.
84. ....Harrop BE, Hartmann DL. The Role of Cloud Radiative Heating in Determining the Location of the ITCZ in Aquaplanet Simulations. *J Climate*. 2016 Jan 29;29(8):2741–63.
85. ....Clark SK, Ming Y, Held IM, Philipps PJ. The role of the water vapor feedback in the ITCZ response to hemispherically asymmetric forcings. *Journal of Climate* [Internet]. 2018 Feb 21 [cited 2018 Apr 11]; Available from: <http://journals.ametsoc.org/doi/10.1175/JCLI-D-17-0723.1>
86. ....Byrne MP, Pendergrass AG, Rapp AD, Wodzicki KR. Response of the Intertropical Convergence Zone to Climate Change: Location, Width, and Strength. *Curr Clim Change Rep*. 2018 Dec 1;4(4):355–70.
87. ....Hilgenbrink CC, Hartmann DL. The Response of Hadley Circulation Extent to an Idealized Representation of Poleward Ocean Heat Transport in an Aquaplanet GCM. *J Climate*. 2018 Oct 16;31(23):9753–70.
88. ..Watt-Meyer O, Frierson DMW. ITCZ Width Controls on Hadley Cell Extent and Eddy-Driven Jet Position and Their Response to Warming. *J Climate*. 2019 Feb 15;32(4):1151–66.
89. ....Held IM. The General Circulation of the Atmosphere. In: *The General Circulation of the Atmosphere: 2000 Program in Geophysical Fluid Dynamics*. Woods Hole Oceanographic Institution; 2000. p. 1–54. (Woods Hole Oceanog. Inst. Tech. Rept.).
90. ....Kang SM, Lu J. Expansion of the Hadley Cell under Global Warming: Winter versus Summer. *J Climate*. 2012 Dec 1;25(24):8387–93.
91. ..Levine XJ, Boos WR. Land surface albedo bias in climate models and its association with tropical rainfall. *Geophys Res Lett*. 2017 Jun 28;44(12):2017GL072510.
92. Zhou W, Xie S-P. Intermodel spread of the double-ITCZ bias in coupled GCMs tied to land surface temperature in AMIP GCMs. *Geophys Res Lett*. 2017 Aug 16;44(15):2017GL074377.
93. Hwang Y-T, Frierson DMW. Link between the double-Intertropical Convergence Zone problem and cloud biases over the Southern Ocean. *PNAS*. 2013 Mar 26;110(13):4935–40.
94. Xiang B, Zhao M, Held IM, Golaz J-C. Predicting the severity of spurious “double ITCZ” problem in CMIP5 coupled models from AMIP simulations. *Geophys Res Lett*. 2017 Feb 16;44(3):2016GL071992.
95. ...Smyth JE, Hill SA, Ming Y. Simulated Responses of the West African Monsoon and Zonal-Mean Tropical Precipitation to Early Holocene Orbital Forcing. *Geophysical Research Letters*. 2018 Nov 16;45(21):12,049–12,057.
96. ..Liu X, Battisti DS, Donohoe A. Tropical Precipitation and Cross-Equatorial Ocean Heat Transport during the Mid-Holocene. *J Climate*. 2017 Jan 23;30(10):3529–47.
97. ....Boos WR, Korty RL. Regional energy budget control of the intertropical convergence zone and application to mid-Holocene rainfall. *Nature Geosci* [Internet]. 2016 Nov 21 [cited 2016 Nov 28];advance online publication. Available from: <http://www.nature.com/ngeo/journal/vaop/ncurrent/full/ngeo2833.html?cookies=accepted>
98. Donohoe A, Voigt A. Why Future Shifts in Tropical Precipitation Will Likely Be Small. In: *Climate Extremes* [Internet]. American Geophysical Union (AGU); 2017 [cited 2019 Jan 9]. p. 115–37. Available from: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/9781119068020.ch8>
99. ..Kang SM, Held IM, Frierson DMW, Zhao M. The Response of the ITCZ to Extratropical Thermal Forcing: Idealized Slab-Ocean Experiments with a GCM. *J Climate*. 2008 Jul 1;21(14):3521–32.
100. Faulk S, Mitchell J, Bordoni S. Effects of Rotation Rate and Seasonal Forcing on the ITCZ Extent in Planetary Atmospheres. *J Atmos Sci*. 2017 Mar;74(3):665–78.

101. . .Shekhar R, Boos WR. Improving Energy-Based Estimates of Monsoon Location in the Presence of Proximal Deserts. *J Climate*. 2016 Apr 4;29(13):4741–61.
102. ....Gonzalez AO, Slocum CJ, Taft RK, Schubert WH. Dynamics of the ITCZ Boundary Layer. *J Atmos Sci*. 2015 Dec 31;73(4):1577–92.
103. ...Gonzalez AO, Schubert WH. Violation of Ekman balance in the Eastern Pacific ITCZ boundary layer. *Journal of the Atmospheric Sciences*. 2019;Submitted.
104. Roberts WHG, Valdes PJ, Singarayer JS. Can energy fluxes be used to interpret glacial/interglacial precipitation changes in the tropics? *Geophys Res Lett*. 2017 Jun 28;44(12):2017GL073103.  
**Demonstrates that relationships between zonal-mean ITCZ and energy transports can differ between seasonal cycle and forced, annual-mean changes and very weakly constrains local precipitation change for most tropical regions.**
105. ....Held IM. The Partitioning of the Poleward Energy Transport between the Tropical Ocean and Atmosphere. *J Atmos Sci*. 2001 Apr 1;58(8):943–8.
106. ....Merlis TM, Schneider T, Bordoni S, Eisenman I. Hadley Circulation Response to Orbital Precession. Part I: Aquaplanets. *J Climate*. 2013;26(3):740–53.
107. . Wei H-H, Bordoni S. Energetic Constraints on the ITCZ Position in Idealized Simulations With a Seasonal Cycle. *Journal of Advances in Modeling Earth Systems*. 2018;10(7):1708–25.
108. ....Xiang B, Zhao M, Ming Y, Yu W, Kang SM. Contrasting Impacts of Radiative Forcing in the Southern Ocean versus Southern Tropics on ITCZ Position and Energy Transport in One GFDL Climate Model. *J Climate*. 2018 Apr 19;31(14):5609–28.
109. ....Inoue K, Back LE. Gross Moist Stability Assessment during TOGA COARE: Various Interpretations of Gross Moist Stability. *J Atmos Sci*. 2015 Aug 17;72(11):4148–66.
110. . Frierson DMW. The Dynamics of Idealized Convection Schemes and Their Effect on the Zonally Averaged Tropical Circulation. *J Atmos Sci*. 2007 Jun 1;64(6):1959–76.
111. ....Wheeler M, Kiladis GN. Convectively coupled equatorial waves: Analysis of clouds and temperature in the wavenumber-frequency domain. *Journal of the Atmospheric Sciences*. 1999;56(3):374–399.
112. ....Frierson DMW. Convectively Coupled Kelvin Waves in an Idealized Moist General Circulation Model. *J Atmos Sci*. 2007 Jun 1;64(6):2076–90.
113. ....Chen G, Held IM. Phase speed spectra and the recent poleward shift of Southern Hemisphere surface westerlies. *Geophysical Research Letters* [Internet]. 2007 Nov 3 [cited 2014 Apr 30];34(21). Available from: <http://doi.wiley.com/10.1029/2007GL031200>
114. .Bischoff T, Schneider T, Meckler AN. A Conceptual Model for the Response of Tropical Rainfall to Orbital Variations. *J Climate*. 2017 Jun 23;30(20):8375–91. **Presents conceptual model for tropical precipitation responses to orbital precession and obliquity forcing based in part on ITCZ shifts framework that captures many features of paleoclimate records at various individual sites.**
115. . .Shanahan TM, McKay NP, Hughen KA, Overpeck JT, Otto-Bliesner B, Heil CW, et al. The time-transgressive termination of the African Humid Period. *Nature Geosci*. 2015 Feb;8(2):140–4.
116. ....Boos WR, Storelvmo T. Near-linear response of mean monsoon strength to a broad range of radiative forcings. *PNAS*. 2016 Feb 9;113(6):1510–5. **Identifies fundamental errors in previous simple theoretical models projecting monsoon “tipping points”, and demonstrates that monsoon responses in a GCM to widely varying forcings are very nearly linear.**
117. Scheff J, Seager R, Liu H, Coats S. Are Glacials Dry? Consequences for Paleoclimatology and for Greenhouse Warming. *J Climate*. 2017 May 22;30(17):6593–609. **Shows that, for land, no single catch-all measure of “wetness” exists, and that plant stomatal responses to CO2 can cause commonly used indices of dryness to err.**

118. ....Scheff J. Drought Indices, Drought Impacts, CO<sub>2</sub>, and Warming: a Historical and Geologic Perspective. *Curr Clim Change Rep.* 2018 Jun 1;4(2):202–9.
119. ....Swann ALS. Plants and Drought in a Changing Climate. *Curr Clim Change Rep.* 2018 Jun 1;4(2):192–201.
120. ....Berg A, Sheffield J. Climate Change and Drought: the Soil Moisture Perspective. *Curr Clim Change Rep.* 2018 Jun 1;4(2):180–91.
121. ..Timm O, Köhler P, Timmermann A, Menviel L. Mechanisms for the Onset of the African Humid Period and Sahara Greening 14.5–11 ka BP. *J Climate.* 2010 Jan 26;23(10):2612–33.
122. Siler N, Roe GH, Armour KC. Insights into the Zonal-Mean Response of the Hydrologic Cycle to Global Warming from a Diffusive Energy Balance Model. *J Climate.* 2018 Jun 19;31(18):7481–93.
123. ....Jansen MF, Nadeau L-P, Merlis TM. Transient versus Equilibrium Response of the Ocean's Overturning Circulation to Warming. *J Climate.* 2018 Apr 2;31(13):5147–63.
124. ....Burls NJ, Fedorov AV, Sigman DM, Jaccard SL, Tiedemann R, Haug GH. Active Pacific meridional overturning circulation (PMOC) during the warm Pliocene. *Science Advances.* 2017 Sep 1;3(9):e1700156.
125. ....Zhang R, Delworth TL. Impact of Atlantic multidecadal oscillations on India/Sahel rainfall and Atlantic hurricanes. *Geophys Res Lett.* 2006 Sep 1;33(17):L17712.
126. ....Park J-Y, Bader J, Matei D. Northern-hemispheric differential warming is the key to understanding the discrepancies in the projected Sahel rainfall. *Nat Commun.* 2015 Jan 21;6:5985.
127. ....Seth A, Giannini A, Rojas M, Rauscher SA, Bordoni S, Singh D, et al. Monsoon Responses to Climate Changes: Connecting Past, Present and Future. *Current Climate Change Reports.* 2019;In review.
128. ....Shepherd TG, Boyd E, Calel RA, Chapman SC, Dessai S, Dima-West IM, et al. Storylines: an alternative approach to representing uncertainty in physical aspects of climate change. *Climatic Change.* 2018 Dec 1;151(3):555–71.
129. ....Zhao M, Golaz J-C, Held IM, Guo H, Balaji V, Benson R, et al. The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 1. Simulation Characteristics With Prescribed SSTs. *Journal of Advances in Modeling Earth Systems.* 2018;10(3):691–734.