

Cook Is.: information on national emissions, population and GDP, and mitigation targets

September 24, 2020

Authors:

Annika Guenther¹

Johannes Guetschow¹

Affiliations:

1. Potsdam Institute for Climate Impact Research, Germany

DOI: [to be added]

TODO

- Table with info on target (main and reclass; emissions from NDC; target quantis + plot).
- GWP: NDC emissions converted from AR2 to AR4 by national conversion factor (2010–2017, PRIMAP-hist v2.1).
- References!

1 Emissions and socio-economic data

With national emissions of 103.0 kt CO₂eq, Cook Is. contributed 0.0002% to global emissions in 2017, while in 2030 its share is estimated to decrease to 0.0001% (Table ??). The estimates for 2030 are based on the downscaled SSP2 Middle of the Road marker scenario (dmSSP2), in which Cook Is. is estimated to emit 92.2 kt CO₂eq in 2030. That change in emissions would constitute a substantial decrease of -10.5% compared to 2017. The pathways dmSSP1–5 show a range of 92.2–92.2 kt CO₂eq in 2030, and 92.2–92.2 kt CO₂eq in 2050. The country’s global rank in terms of total per-capita emissions was 90 in 2017 (103 in 2030). In terms of accumulated historical emissions, Cook Is. contributed to the global 1850–2017 emissions by 0.0001%. When only accounting for the years 1990–2017, its contribution stays the same to 0.0001%. All of the emissions are presented following GWP AR4, and exclude emissions from LULUCF (exclLU), and bunkers fuels emissions (exclBunkers).

Table 1: National emissions (dmSSP2), GDP and population for Cook Is., together with the emissions per unit of GDP and per capita emissions (all for 2017 and 2030). Additionally, the global share and its rank are displayed.

	Year	Total	Unit	Glob. share	Rank
Emissions	2017	103.0	kt CO ₂ eq	0.0002%	200
	2030	92.2	kt CO ₂ eq	0.0001%	200
GDP	2017	–	2011 GK\$	–	–
	2030	–	2011 GK\$	–	–
Emissions per GDP	2017	–	t CO ₂ eq / 2011 GK\$	–	–
	2030	–	t CO ₂ eq / 2011 GK\$	–	–
Population	2017	17.5	Thousand Pers	0.0002%	199
	2030	17.7	Thousand Pers	0.0002%	198
Emissions per capita	2017	5.9	t CO ₂ eq / Pers	0.3%	90
	2030	5.2	t CO ₂ eq / Pers	0.3%	103

For Cook Is., in 2017 the main emissions share on sectoral level (Fig. ??) came from the Energy sector (83.1%), followed by Agriculture (15.8%) The Kyoto GHG with the highest emissions in 2017 was CO₂, constituting as much as 82.9% of the national emissions. Second largest contributor was CH₄ (13.9%) The total of F-gasesonly represented 0.0%. The total CO₂ emissions are expected to be 81.4% of the national Kyoto GHG emissions in 2030 (dmSSP2).




Figure 1: 'Stacked' timeseries of national emissions (exclLU) per main-sector (a) and Kyoto GHG (b). No information available on the sectoral contributions after 2017.




Figure 2: Timeseries of national GDP (a) and population (c), and Kyoto GHG emissions (exclLU, exclBunkers) per unit of GDP (b) or per capita (d).

The population decreased, while the per capita emissions dropped. Cook Is.'s population is assumed to grow after 2017 but to diminish again before 2050, and the per capita emissions are expected to decline towards 2050. For Cook Is., no GDP data are available from dmSSPs.

LULUCF emissions data for Cook Is. are available from the following sources (Fig. ??): FAO (2019) / 28, with the number of available data points in 1990–2017 displayed additionally. Based on data from FAO (2019), for the year 2017, LULUCF is estimated to be a net sink of -5.2 kt CO₂eq, which in absolute terms is lower than the non-LULUCF emissions of 103.0 kt CO₂eq. The emissions range for FAO (2019) and 1990–2017 is -73.3–0.0 kt CO₂eq.




Figure 3: Timeseries of emissions from LULUCF (CO₂ plus CH₄ and N₂O) as available from different data-sources. Indicated in pink are the 'chosen' data, as used in our assessment of Cook Is.'s NDC (if needed). The pink timeseries was inter- and / or extrapolated (interpolation: linear, extrapolation: constant).

2 Mitigation targets (NDC)

Give the %cov for the base and target year (and 2017).

Global share for 2030 for the mitigated pathways and % reduction relative to 1990 and 2017.

Table with the 'input' data and the resulting targets (like `ndcs_targets.csv`).

Cook Is. has an NDC, with a GHG mitigation target of the type RBY (base year target; main target type). The reclassified target type is ABS (absolute emissions target).

Table 2: Information on Cook Is.'s GHG mitigation target(s).

type	condi.	range	value	refYr	tarYr	LU
ABS	condi.	best	0.050 Mt CO ₂ eq	2006	2030	exclLU
RBY	condi.	best	-34.9%	2006	2030	exclLU




Figure 4: Timeseries of Cook Is.'s national emissions (exclLU) and the share of emissions that is assumed to be covered by Cook Is.'s mitigation target.

Table 3: Information on covered sectors and gases as retrieved from NDC and adapted ('Adap.': used to calculate %cov), and their shares in Cook Is.'s 2017 emissions (exclLU, exclBunkers; total 103.0 kt CO₂eq). If either the sector or gas is assessed as 'not-covered', the emissions from this sector-gas combination are counted as not-covered (-). Else the emissions are counted as covered (+; covered shares given in bold). (/) means that no information is available. LULUCF: NDC '/' and adapted '-' (estimated as a net sink of -5.2 kt CO₂eq in 2017; based on the 'chosen' LULUCF emissions).

	NDCs	Adap.	CO ₂	CH ₄	N ₂ O	HFCs	PFCs	SF ₆	NF ₃	Total
NDCs			/	/	/	/	/	/	/	
Adap.			+	+	+	-	-	-	-	
Energy	+	+	82.9%	0.06%	0.1%	/	/	/	/	83.1%
IPPU	/	-	0.0%	/	0.1%	/	/	/	/	0.1%
Agri.	/	-	/	13.0%	2.9%	/	/	/	/	15.8%
Waste	/	-	/	0.8%	/	/	/	/	/	0.8%
Other	/	-	/	/	0.1%	/	/	/	/	0.1%
Total			82.9%	13.9%	3.2%	/	/	/	/	100.0%




Figure 5: Quantified mitigation targets (based on different input data and calculation options). Vertical lines: conditionality / range; colour coded: dmSSP1–5; first / second set of six: prio NDCs / SSPs; set of six: coverage 100, lulucf unfccc, lulucf fao, bl uncondi, const emi, estimated coverage.

3 Data sources, additional information and references

PRIMAP-hist v2.1: emissions from PRIMAP-hist are data from the country reported data priority scenario (HISTCR).

dmSSPs: emissions, population and GDP data are PMSSPBIE data for the five marker scenarios.

SSPs Shared Socio-economic Pathways.

Narratives and challenges to mitigation and adaptation:

SSP1: Sustainability, Taking the Green Road (low / low);

SSP2: Middle of the Road (medium / medium);

SSP3: Regional Rivalry, A Rocky Road (high / high);

SSP4: Inequality, A Road Divided (low / high); and

SSP5: Fossil-fuelled Development, Taking the Highway (high / low).

GDP Gross Domestic Product.

Throughout this document the GDP is given as GDP PPP, with PPP being the Purchasing Power Parity.

GWP Global Warming Potential.

We use GWP values from the IPCC 4th Assessment Report (AR4). They reflect the forcing potential of one kilogram of a gas' emissions in comparison to one kilogram of CO₂ (GWP_{CO2} = 1). The GWPs correspond to a 100-yr period and are for CH₄: 25, for N₂O: 298, for SF₆: 22800, and for NF₃: 17200. For the basket of HFC-gases the GWPs from AR4 are in the range 4–14800, and for PFCs 7190–12200. To assess emissions of several GHGs, their emissions are weighted by their respective GWPs and presented in CO₂ equivalents (CO₂eq).

LULUCF Land Use, Land-Use Change and Forestry.

Emissions from LULUCF are excluded throughout the document, unless stated otherwise.

Bunkers fuels Emissions from international aviation and shipping.

Kyoto GHG Kyoto GHG (Greenhouse Gas) basket.

Carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulfur hexafluoride (SF₆), and nitrogen trifluoride (NF₃).

F-gases Fluorinated gases.

Basket of HFCs, PFCs, and the gases SF₆ and NF₃. Some F-gases have very long atmospheric lifetimes and high Global Warming Potentials.

Target reclassification When a country has, e.g., an RBU target (relative reduction compared to Business-As-Usual), and the BAU emissions are provided, it can be quantified based on the given emissions, and is reclassified from type_main RBU to type_reclass ABS (absolute emissions target). Additionally, 'NGT' targets can be reclassified as 'ABU' (absolute reduction compared to Business-As-Usual) if absolute mitigation effects due to planned policies and measures are provided.

Quantification options Different quantification options were tested.

dmSSP1–5: down-scaled SSP marker scenarios;

type_reclass: external data prioritised (PRIMAP-hist, dmSSPs);

type_main: emissions data from within NDCs were prioritised;

100% coverage & estimated coverage;

constant emi: constant emissions after last target year (instead of constant relative difference to baseline);
baseline uncondi: baseline emissions as uncond. pathways for Parties without uncond. targets, even if baseline is better than cond. targets (instead of cond. pathway as uncond. pathways in these cases).

Links to additional information:

- [CLIMATEWATCH](#)
- [CarbonBrief: Clear on Climate](#)
- [China's carbon neutral pledge could curb global warming by 0.3°C - researchers](#) (23 September 2020)
- [Climate Action Tracker](#)
- [Coronavirus: Climate action cannot be another Covid victim - PM](#) (23 September 2020)
- [Countries' Climate Plans \(NDCs\) Are Missing a Big Opportunity: Reducing Food Loss and Waste](#) (3 July 2019)
- [Country resolved combined emission and socio-economic pathways based on the RCP and SSP scenarios](#) (February 2020)
- [Few countries living up to Covid 'green recovery' pledges - analysis](#) (23 September 2020)
- [Guest post: Calculating the true climate impact of aviation emissions](#) (September 2020)
- [IGES NDC Database](#)
- [IPCC \(The Intergovernmental Panel on Climate Change\)](#)
- [IPCC Special Report: Global Warming of 1.5°](#) (2018)
- [ISIMIP / ISIpedia](#)
- [Melting Antarctic ice will raise sea level by 2.5 metres - even if Paris climate goals are met, study finds](#) (23 September 2020)
- [NDC Explorer](#)
- [NDC PARTNERSHIP](#)
- [PBL Climate Pledge NDC tool](#)
- [SSP Database \(Shared Socioeconomic Pathways\) - Version 2.0](#) (December 2018)
- [The PRIMAP-hist national historical emissions time series \(1850-2017\)](#) (2019)
- [UNFCCC \(United Nations Framework Convention on Climate Change\)](#)
- [WORLD RESOURCES INSTITUTE](#)
- [Why the 2020 Atlantic hurricane season has spun out of control: Extra-warm ocean waters, boosted by climate change, and La Niña are key drivers in historic season.](#) (September 2020)
- [World's richest 1% cause double CO₂ emissions of poorest 50%, says Oxfam](#)
- [#showyourbudgets](#)