

# Argentina: information on national emissions, population and GDP, and mitigation targets

September 24, 2020

## Authors:

Annika Guenther<sup>1</sup>

Johannes Guetschow<sup>1</sup>

## Affiliations:

1. Potsdam Institute for Climate Impact Research, Germany

**DOI:** [to be added]

## TODO

- Table with info on target (main and reclass; emissions from NDC; target quantis + plot).
- GWP: NDC emissions converted from AR2 to AR4 by national conversion factor (2010–2017, PRIMAP-hist v2.1).
- References!

# 1 Emissions and socio-economic data

With national emissions of 348.2 Mt CO<sub>2</sub>eq, Argentina contributed 0.7% to global emissions in 2017, and in 2030 its share is estimated to stay at a similar level (Table ??). The estimates for 2030 are based on the downscaled SSP2 Middle of the Road marker scenario (dmSSP2), in which Argentina is estimated to emit 403.7 Mt CO<sub>2</sub>eq in 2030. That change in emissions would constitute a substantial increase of 15.9% compared to 2017. The pathways dmSSP1–5 show a range of 384.9–486.7 Mt CO<sub>2</sub>eq in 2030, and 414.0–570.6 Mt CO<sub>2</sub>eq in 2050. The country’s global rank in terms of total emissions per unit of GDP was 87 in 2017, and 59 regarding the per-capita emissions (84 and 52 in 2030). In terms of accumulated historical emissions, Argentina contributed to the global 1850–2017 emissions by 0.7%. When only accounting for the years 1990–2017, its contribution increases to 0.8%. All of the emissions are presented following GWP AR4, and exclude emissions from LULUCF (exclLU), and bunkers fuels emissions (exclBunkers).

Table 1: National emissions (dmSSP2), GDP and population for Argentina, together with the emissions per unit of GDP and per capita emissions (all for 2017 and 2030). Additionally, the global share and its rank are displayed.

|                             | Year | Total | Unit                                      | Glob. share | Rank |
|-----------------------------|------|-------|---|-------------|------|
| <b>Emissions</b>            | 2017 | 348.2 | Mt CO <sub>2</sub> eq                     | 0.7%        | 26   |
|                             | 2030 | 403.7 | Mt CO <sub>2</sub> eq                     | 0.7%        | 28   |
| <b>GDP</b>                  | 2017 | 0.7   | Trillion 2011 GK\$                        | 0.7%        | 29   |
|                             | 2030 | 1.2   | Trillion 2011 GK\$                        | 0.6%        | 31   |
| <b>Emissions per GDP</b>    | 2017 | 0.4   | t CO <sub>2</sub> eq / Thousand 2011 GK\$ | 0.4%        | 87   |
|                             | 2030 | 0.3   | t CO <sub>2</sub> eq / Thousand 2011 GK\$ | 0.4%        | 84   |
| <b>Population</b>           | 2017 | 43.9  | Million Pers                              | 0.6%        | 32   |
|                             | 2030 | 47.5  | Million Pers                              | 0.6%        | 35   |
| <b>Emissions per capita</b> | 2017 | 7.9   | t CO <sub>2</sub> eq / Pers               | 0.5%        | 59   |
|                             | 2030 | 8.5   | t CO <sub>2</sub> eq / Pers               | 0.6%        | 52   |

For Argentina, in 2017 the main emissions share on sectoral level (Fig. ??) came from the Energy sector (56.7%), followed by Agriculture (32.8%) The Kyoto GHG with the highest emissions in 2017 was CO<sub>2</sub>, constituting 58.7% of the national emissions. Second largest contributor was CH<sub>4</sub> (27.8%) The total of F-gases only represented 0.2%. The trend in total emissions is mostly driven by CH<sub>4</sub> and N<sub>2</sub>O emissions from the Agriculture sector, which contributed 20.4% and 12.2% to Argentina’s 2017 emissions.<sup>1</sup> The total CO<sub>2</sub> emissions are expected to be 63.1% of the national Kyoto GHG emissions in 2030 (dmSSP2).

<sup>1</sup>Analysis based on the correlations between total national emissions (exclLU) versus the emissions of the combinations of main-sectors & the gases CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and F-gases. Only data from 2010 to 2017 are assessed. The (up to) three gas & sector combinations are chosen for which the slope of the regression line to the correlated values exceeds 0.2.




Figure 1: 'Stacked' timeseries of national emissions (exclLU) per main-sector (a) and Kyoto GHG (b). No information available on the sectoral contributions after 2017.




Figure 2: Timeseries of national GDP (a) and population (c), and Kyoto GHG emissions (exclLU, exclBunkers) per unit of GDP (b) or per capita (d).

The national GDP increased in recent years, and the emissions per unit of GDP had a similar trend (Fig. ??). The population increased, and the per capita emissions were on the rise. Following dmSSP2, the GDP is projected to increase towards 2050. The emissions per GDP are estimated to decrease towards 2050. Argentina's population is assumed to grow towards 2050, and the per capita emissions are expected to increase after 2017 but to decline again before 2050.

LULUCF emissions data for Argentina are available from the following sources (Fig. ??): BUR (2<sup>nd</sup>) / 25, UNFCCC (2019) / 6, FAO (2019) / 28, with the number of available data points in 1990–2017 displayed additionally. Based on data from UNFCCC (2019), for the year 2017, LULUCF is estimated to be a net source of 70.4 Mt CO<sub>2</sub>eq, which in absolute terms is lower than the non-LULUCF emissions of 348.2 Mt CO<sub>2</sub>eq. The emissions range for UNFCCC (2019) and 1990–2017 is 45.7–104.7 Mt CO<sub>2</sub>eq.



Figure 3: Timeseries of emissions from LULUCF (CO<sub>2</sub> plus CH<sub>4</sub> and N<sub>2</sub>O) as available from different data-sources. Indicated in pink are the 'chosen' data, as used in our assessment of Argentina's NDC (if needed). The pink timeseries was inter- and / or extrapolated (interpolation: linear, extrapolation: constant).

## 2 Mitigation targets (NDC)

Give the %cov for the base and target year (and 2017).

Global share for 2030 for the mitigated pathways and % reduction relative to 1990 and 2017.

Table with the 'input' data and the resulting targets (like `ndcs_targets.csv`).

Argentina has an NDC, with a GHG mitigation target of the type ABS (absolute emissions target; main target type). The reclassified target type equals the main target type.

Table 2: Information on Argentina's GHG mitigation target(s).

| type | condi.   | range | value                         | tarYr | LU     |
|------|----------|-------|-------------------------------|-------|--------|
| ABS  | uncondi. | best  | 483 Mt CO <sub>2</sub> eq SAR | 2030  | inclLU |
| ABS  | condi.   | best  | 369 Mt CO <sub>2</sub> eq SAR | 2030  | inclLU |




Figure 4: Timeseries of Argentina's national emissions (exclLU) and the share of emissions that is assumed to be covered by Argentina's mitigation target.

Table 3: Information on covered sectors and gases as retrieved from NDC and adapted ('Adap.': used to calculate %cov), and their shares in Argentina's 2017 emissions (exclLU, exclBunkers; total 348.2 Mt CO<sub>2</sub>eq). If either the sector or gas is assessed as 'not-covered', the emissions from this sector-gas combination are counted as not-covered (-). Else the emissions are counted as covered (+; covered shares given in bold). (/) means that no information is available. LULUCF: NDC '+' and adapted '+' (estimated as a net source of 70.4 Mt CO<sub>2</sub>eq in 2017; based on the 'chosen' LULUCF emissions).

|        | NDCs | Adap. | CO <sub>2</sub> | CH <sub>4</sub> | N <sub>2</sub> O | HFCs        | PFCs         | SF <sub>6</sub> | NF <sub>3</sub> | Total  |
|--------|------|-------|-----------------|-----------------|------------------|-------------|--------------|-----------------|-----------------|--------|
| NDCs   |      |       | +               | +               | +                | +           | +            | +               | /               |        |
| Adap.  |      |       | +               | +               | +                | +           | +            | +               | -               |        |
| Energy | +    | +     | <b>53.6%</b>    | <b>2.6%</b>     | <b>0.5%</b>      | /           | /            | /               | /               | 56.7%  |
| IPPU   | +    | +     | <b>4.8%</b>     | <b>0.02%</b>    | <b>0.03%</b>     | <b>0.2%</b> | <b>0.05%</b> | <b>0.0%</b>     | /               | 5.2%   |
| Agri.  | +    | +     | <b>0.2%</b>     | <b>20.4%</b>    | <b>12.2%</b>     | /           | /            | /               | /               | 32.8%  |
| Waste  | +    | +     | <b>0.0%</b>     | <b>4.8%</b>     | <b>0.2%</b>      | /           | /            | /               | /               | 5.0%   |
| Other  | /    | +     | /               | /               | <b>0.3%</b>      | /           | /            | /               | /               | 0.3%   |
| Total  |      |       | 58.7%           | 27.8%           | 13.3%            | 0.2%        | 0.05%        | 0.0%            | /               | 100.0% |




Figure 5: Quantified mitigation targets (based on different input data and calculation options). Vertical lines: conditionality / range; colour coded: dmSSP1–5; first / second set of six: prio NDCs / SSPs; set of six: coverage 100, lulucf unfccc, lulucf fao, bl uncondi, const emi, estimated coverage.

### 3 Data sources, additional information and references

**PRIMAP-hist v2.1:** emissions from PRIMAP-hist are data from the country reported data priority scenario (HISTCR).

**dmSSPs:** emissions, population and GDP data are PMSSPBIE data for the five marker scenarios.

**SSPs** Shared Socio-economic Pathways.

Narratives and challenges to mitigation and adaptation:

*SSP1*: Sustainability, Taking the Green Road (low / low);

*SSP2*: Middle of the Road (medium / medium);

*SSP3*: Regional Rivalry, A Rocky Road (high / high);

*SSP4*: Inequality, A Road Divided (low / high); and

*SSP5*: Fossil-fuelled Development, Taking the Highway (high / low).

**GDP** Gross Domestic Product.

Throughout this document the GDP is given as GDP PPP, with PPP being the Purchasing Power Parity.

**GWP** Global Warming Potential.

We use GWP values from the IPCC 4<sup>th</sup> Assessment Report (AR4). They reflect the forcing potential of one kilogram of a gas' emissions in comparison to one kilogram of CO<sub>2</sub> (GWP<sub>CO2</sub> = 1). The GWPs correspond to a 100-yr period and are for CH<sub>4</sub>: 25, for N<sub>2</sub>O: 298, for SF<sub>6</sub>: 22800, and for NF<sub>3</sub>: 17200. For the basket of HFC-gases the GWPs from AR4 are in the range 4–14800, and for PFCs 7190–12200. To assess emissions of several GHGs, their emissions are weighted by their respective GWPs and presented in CO<sub>2</sub> equivalents (CO<sub>2</sub>eq).

**LULUCF** Land Use, Land-Use Change and Forestry.

Emissions from LULUCF are excluded throughout the document, unless stated otherwise.

**Bunkers fuels** Emissions from international aviation and shipping.

**Kyoto GHG** Kyoto GHG (Greenhouse Gas) basket.

Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulfur hexafluoride (SF<sub>6</sub>), and nitrogen trifluoride (NF<sub>3</sub>).

**F-gases** Fluorinated gases.

Basket of HFCs, PFCs, and the gases SF<sub>6</sub> and NF<sub>3</sub>. Some F-gases have very long atmospheric lifetimes and high Global Warming Potentials.

**Target reclassification** When a country has, e.g., an RBU target (relative reduction compared to Business-As-Usual), and the BAU emissions are provided, it can be quantified based on the given emissions, and is reclassified from type\_main RBU to type\_reclass ABS (absolute emissions target). Additionally, 'NGT' targets can be reclassified as 'ABU' (absolute reduction compared to Business-As-Usual) if absolute mitigation effects due to planned policies and measures are provided.

**Quantification options** Different quantification options were tested.

*dmSSP1–5*: down-scaled SSP marker scenarios;

*type\_reclass*: external data prioritised (PRIMAP-hist, dmSSPs);

*type\_main*: emissions data from within NDCs were prioritised;

*100% coverage & estimated coverage*;



*constant emi*: constant emissions after last target year (instead of constant relative difference to baseline);  
*baseline uncondi*: baseline emissions as uncond. pathways for Parties without uncond. targets, even if baseline is better than cond. targets (instead of cond. pathway as uncond. pathways in these cases).

### Links to additional information:

- [CLIMATEWATCH](#)
- [CarbonBrief: Clear on Climate](#)
- [China's carbon neutral pledge could curb global warming by 0.3°C - researchers](#) (23 September 2020)
- [Climate Action Tracker](#)
- [Coronavirus: Climate action cannot be another Covid victim - PM](#) (23 September 2020)
- [Countries' Climate Plans \(NDCs\) Are Missing a Big Opportunity: Reducing Food Loss and Waste](#) (3 July 2019)
- [Country resolved combined emission and socio-economic pathways based on the RCP and SSP scenarios](#) (February 2020)
- [Few countries living up to Covid 'green recovery' pledges - analysis](#) (23 September 2020)
- [Guest post: Calculating the true climate impact of aviation emissions](#) (September 2020)
- [IGES NDC Database](#)
- [IPCC \(The Intergovernmental Panel on Climate Change\)](#)
- [IPCC Special Report: Global Warming of 1.5°](#) (2018)
- [ISIMIP / ISIpedia](#)
- [Melting Antarctic ice will raise sea level by 2.5 metres - even if Paris climate goals are met, study finds](#) (23 September 2020)
- [NDC Explorer](#)
- [NDC PARTNERSHIP](#)
- [PBL Climate Pledge NDC tool](#)
- [SSP Database \(Shared Socioeconomic Pathways\) - Version 2.0](#) (December 2018)
- [The PRIMAP-hist national historical emissions time series \(1850-2017\)](#) (2019)
- [UNFCCC \(United Nations Framework Convention on Climate Change\)](#)
- [WORLD RESOURCES INSTITUTE](#)
- [Why the 2020 Atlantic hurricane season has spun out of control: Extra-warm ocean waters, boosted by climate change, and La Niña are key drivers in historic season.](#) (September 2020)
- [World's richest 1% cause double CO<sub>2</sub> emissions of poorest 50%, says Oxfam](#)
- [#showyourbudgets](#)