Eq. Guinea: information on national emissions, population and GDP, and mitigation targets

September 23, 2020

Authors:

Annika Guenther¹ Johannes Guetschow¹

Affiliations:

1. Potsdam Institute for Climate Impact Research, Germany

DOI: [to be added]

TODO

- Table with info on target (main and reclass; emissions from NDC; target quantis + plot).
- GWP: NDC emissions coverted from AR2 to AR4 by national conversion factor (2010–2017, PRIMAP-hist v2.1).
- References!

1 Non-LULUCF emissions and socio-economic data

With national emissions of 13.0 Mt CO₂eq, Eq. Guinea contributed 0.02% to global emissions in 2017, and in 2030 its share is estimated to stay at a similar level (Table ??). The estimates for 2030 are based on the downscaled SSP2 Middle of the Road marker scenario (dmSSP2), in which Eq. Guinea is estimated to emit 16.6 Mt CO₂eq in 2030. That change in emissions would constitute a substantial increase of 27.4% compared to 2017. The pathways dmSSP1–5 show a range of 14.1–18.9 Mt CO₂eq in 2030, and 11.1–24.9 Mt CO₂eq in 2050. The country's global rank in terms of total emissions per unit of GDP was 59 in 2017, and 37 regarding the per-capita emissions (55 and 39 in 2030). In terms of accumulated historical emissions, Eq. Guinea contributed to the global 1850–2017 emissions by 0.01%. When only accounting for the years 1990–2017, its contribution increases to 0.02%. All of the emissions are presented following GWP AR4, and exclude emissions from LULUCF (exclLU), and bunkers fuels emissions (exclBunkers).

Table 1: National emissions (dmSSP2), GDP and population for Eq. Guinea, together with the emissions per unit of GDP and per capita emissions (all for 2017 and 2030). Additionally, the global share and its rank are displayed.

	Year	Total	Unit	Glob. share	Rank
Emissions	2017	13.0	Mt CO ₂ eq	0.02%	136
	2030	16.6	$Mt CO_2eq$	0.02%	132
GDP	2017	20.2	Billion 2011 GK\$	0.01%	144
	2030	38.3	Billion 2011 GK\$	0.02%	143
Emissions	2017	0.6	t CO ₂ eq / Thousand 2011 GK\$	0.6%	59
per GDP	2030	0.4	t CO_2eq / Thousand 2011 GK\$	0.6%	55
Population	2017	1.3	Million Pers	0.01%	154
	2030	1.6	Million Pers	0.01%	151
Emissions	2017	10.3	t CO ₂ eq / Pers	0.7%	37
per capita	2030	10.1	t CO ₂ eq / Pers	0.7%	39

For Eq. Guinea, in 2017 the main emissions share on sectoral level (Fig. ??) came from the Energy sector (87.8%), followed by IPPU (9.3%) The Kyoto GHG with the highest emissions in 2017 was CO₂, constituting 55.4% of the national emissions. Second largest contributor was CH₄ (44.3%) The total of F-gasesonly represented 0.0%. The trend in total emissions is mostly driven by CH₄ emissions from the Energy sector, which contributed 40.9% to Eq. Guinea's 2017 emissions.¹ The total CO₂ emissions are expected to be 59.7% of the national Kyoto GHG emissions in 2030 (dmSSP2).

¹Analysis based on the correlations between total national emissions (exclLU) versus the emissions of the combinations of main-sectors & the gases CO₂, CH₄, N₂O and F-gases. Only data from 2010 to 2017 are assessed. The (up to) three gas & sector combinations are chosen for which the slope of the regression line to the correlated values exceeds 0.2.




Figure 1: 'Stacked' timeseries of national emissions (exclLU) per main-sector (a) and Kyoto GHG (b). No information available on the sectoral contributions after 2017.




Figure 2: Timeseries of national GDP (a) and population (c), and Kyoto GHG emissions (exclLU, exclBunkers) per unit of GDP (b) or per capita (d).

The national GDP decreased in recent years, and the emissions per unit of GDP had a similar trend (Fig. ??). The population increased, while the per capita emissions were on the rise. Following dmSSP2, the GDP is projected to increase towards 2050. The emissions per GDP are estimated to decrease towards 2050. Eq. Guinea's population is assumed to grow towards 2050, and the per capita emissions are expected to increase after 2017 but to decline again before 2050.

2 LULUCF emissions

LULUCF emissions data for Eq. Guinea are available from the following sources (Fig. ??): FAO (2019).

High fluctuations? Data gaps? Difference between sources?




Figure 3: Timeseries of emissions from LULUCF (CO_2 plus CH_4 and N_2O) as available from different data-sources. Indicated in pink are the 'chosen' data, as used in our assessment of Eq. Guinea's NDC (if needed). The pink timeseries was inter- and / or extrapolated (interpolation: linear, extrapolation: constant).

3 Mitigation targets (NDC)

Give the %cov for the base and target year (and 2017). Global share for 2030 for the mitigated pathways and % reduction relative to 1990 and 2017. Table with the 'input' data and the resulting targets (like ndcs_targets.csv). Eq. Guinea has an NDC, with a GHG mitigation target of the type RBY (base year target; main target type). The reclassified target type equals the main target type.

Table 2: Information on Eq. Guinea's GHG mitigation target(s).

\mathbf{type}	condi.	range	value	refYr	tarYr	LU
RBY	condi.	best	-20%	2010	2030	inclLU
RBY	condi.	best	-50%	2010	2050	inclLU




Figure 4: Timeseries of Eq. Guinea's national emissions (exclLU) and the share of emissions that is assumed to be covered by Eq. Guinea's mitigation target.

Table 3: Information on covered sectors and gases as retrieved from NDC and adapted ('Adap.': used to calculate %cov), and their shares in Eq. Guinea's 2017 emissions (exclLU, exclBunkers; total 13.0 Mt CO₂eq). If either the sector or gas is assessed as 'not-covered', the emissions from this sector-gas combination are counted as not-covered (–). Else the emissions are counted as covered (+; covered shares given in bold). (/) means that no information is available. LULUCF: NDC '+' and adapted '+' (estimated as a net source of 5.4 Mt CO₂eq in 2017; based on the 'chosen' LULUCF emissions).

	NDCs	Adap.	\mathbf{CO}_2	\mathbf{CH}_4	N_2O	\mathbf{HFCs}	PFCs	\mathbf{SF}_6	\mathbf{NF}_3	Total
NDCs			/	/	/	/	/	/	/	
Adap.			+	+	+	_	_	_	_	
Energy	+	+	46.8%	40.9%	0.1%	/	/	/	/	87.8%
\mathbf{IPPU}	+	+	8.6%	0.7%	0.0%	/	/	/	/	9.3%
Agri.	+	+	/	0.1%	0.07%	/	/	/	/	0.1%
Waste	+	+	/	2.7%	0.0%	/	/	/	/	2.7%
\mathbf{Other}	/	+	/	/	0.03%	/	/	/	/	0.03%
Total			55.4%	44.3%	0.2%	/	/	/	/	100.0%




Figure 5: Quantified mitigation targets (based on different input data and calculation options). Vertical lines: conditionality / range; colour coded: dmSSP1-5; first / second set of six: prio NDCs / SSPs; set of six: coverage 100, lulucf unfccc, lulucf fao, bl uncondi, const emi, estimated coverage.

4 Data sources, additional information and references

PRIMAP-hist v2.1: emissions from PRIMAP-hist are data from the country reported data priority scenario (HISTCR).

dmSSPs: emissions, population and GDP data are PMSSPBIE data for the five marker scenarios.

- SSPs Shared Socio-economic Pathways. Narratives and challenges to mitigation and adaptation: SSP1: Sustainability, Taking the Green Road (low / low); SSP2: Middle of the Road (medium / medium); SSP3: Regional Rivalry, A Rocky Road (high / high); SSP4: Inequality, A Road Divided (low / high); and SSP5: Fossil-fuelled Development, Taking the Highway (high / low).
- **GDP** Gross Domestic Product. Throughout this document the GDP is given as GDP PPP, with PPP being the Purchasing Power Parity.
- **GWP** Global Warming Potential. we use GWP values from the IPCC 4^{th} Assessment Report (AR4). They reflect the forcing potential of one kilogram of a gas' emissions in comparison to one kilogram of CO_2 (GWP_{CO2} = 1). The GWPs correspond to a 100-yr period and are for CH₄: 25, for N₂O: 298, for SF₆: 22800, and for NF₃: 17200. For the basket of HFC-gases the GWPs from AR4 are in the range 4–14800, and for PFCs 7190–12200. To assess emissions of several GHGs, their emissions are weighted by their respective GWPs and presented in CO_2 equivalents (CO_2 eq).
- **LULUCF** Land Use, Land-Use Change and Forestry. Emissions from LULUCF are excluded throughout the document, unless stated otherwise.

Bunkers fuels Emissions from international aviation and shipping.

- **Kyoto GHG** Kyoto GHG (Greenhouse Gas) basket: carbon dioxide (CO_2) , methane (CH_4) , nitrous oxide (N_2O) , hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulfur hexafluoride (SF_6) , and nitrogen trifluoride (NF_3) .
- **F-gases** Fluorinated gases.Basket of HFCs, PFCs, and the gases SF₆ and NF₃. Some F-gases have very long atmospheric lifetimes and high Global Warming Potentials.
- Target reclassification When a country has, e.g., an RBU target (relative reduction compared to Business-As-Usual), and the BAU emissions are provided, it can be quantified based on the given emissions, and is reclassified from type_main RBU to type_reclass ABS (absolute emissions target). Additionally, 'NGT' targets can be reclassified as 'ABU' (absolute reduction compared to Business-As-Usual) if absolute mitigation effects due to planned policies and measures are provided.

Links to additional information:

- CLIMATEWATCH
- CarbonBrief: Clear on Climate
- Climate Action Tracker
- Country resolved combined emission and socio-economic pathways based on the RCP and SSP scenarios (February 2020)
- Guest post: Calculating the true climate impact of aviation emissions (September 2020)
- IGES NDC Database

- IPCC (The Intergovernmental Panel on Climate Change)
- IPCC Special Report: Global Warming of 1.5° (2018)
- ISIMIP / ISIpedia
- NDC Explorer
- NDC PARTNERSHIP
- PBL Climate Pledge NDC tool
- SSP Database (Shared Socioeconomic Pathways) Version 2.0 (December 2018)
- The PRIMAP-hist national historical emissions time series (1850-2017) (2019)
- UNFCCC (United Nations Framework Convention on Climate Change)
- World's richest 1% cause double CO₂ emissions of poorest 50%, says Oxfam
- #showyourbudgets